

All pneumoconioses: Number of deaths, death rates (per million population), and years of potential life lost (YPLL) by state, U.S. residents age 15 and over, 1995–2004

State	No. of Deaths		Death Rate				YPLL to Life Expectancy			
			Crude	Rank	Age-Adjusted	Rank	Total	Rank	YPLL/death	Rank
Alabama	671	12	19.00	8	18.85	8	7,740	12	11.5	20
Alaska	28	49	5.95	35	14.82	12	292	49	10.4	44
Arizona	221	28	5.56	39	5.48	39	2,375	28	10.7	37
Arkansas	162	32	7.65	28	7.03	30	1,989	32	12.3	7
California	1,308	4	5.00	41	5.80	37	14,600	5	11.2	24
Colorado	301	23	8.89	20	11.65	17	3,248	23	10.8	35
Connecticut	201	30	7.47	29	6.77	31	2,332	29	11.6	17
Delaware	150	33	23.89	5	24.43	5	1,669	33	11.1	28
District of Columbia	10	51	2.09	51	2.28	51	156	51	15.6	2
Florida	1,070	8	8.29	23	6.25	34	11,744	8	11.0	30
Georgia	253	25	3.96	47	5.15	42	3,039	25	12.0	11
Hawaii	30	48	3.09	50	3.03	49	410	46	13.7	3
Idaho	81	40	8.15	27	8.80	25	906	40	11.2	24
Illinois	619	13	6.35	33	6.55	33	6,853	14	11.1	28
Indiana	308	22	6.45	32	6.57	32	3,455	21	11.2	24
Iowa	128	35	5.46	40	4.51	44	1,279	36	10.0	47
Kansas	96	39	4.54	43	4.35	45	1,254	38	13.1	5
Kentucky	1,237	6	37.97	3	39.71	3	14,486	6	11.7	14
Louisiana	332	20	9.53	18	10.32	20	4,420	20	13.3	4
Maine	134	34	12.87	14	11.71	16	1,446	34	10.8	35
Maryland	568	15	13.61	12	15.13	11	6,562	16	11.6	17
Massachusetts	422	19	8.22	25	7.70	26	4,430	19	10.5	43
Michigan	454	18	5.81	36	5.98	35	5,231	18	11.5	20
Minnesota	225	27	5.79	38	5.92	36	2,486	27	11.0	30
Mississippi	462	16	20.82	6	21.61	7	7,975	11	17.3	1
Missouri	198	31	4.46	44	4.22	46	2,313	30	11.7	14
Montana	101	38	13.97	11	13.27	14	1,076	39	10.7	37
Nebraska	50	45	3.71	48	3.37	48	464	45	9.3	50
Nevada	68	43	4.37	45	5.16	41	844	41	12.4	6
New Hampshire	66	44	6.71	31	7.14	29	736	43	11.2	24
New Jersey	977	9	14.65	9	14.01	13	10,315	9	10.6	41
New Mexico	125	36	8.88	21	9.83	23	1,262	37	10.1	45
New York	711	10	4.70	42	4.65	43	8,006	10	11.3	23
North Carolina	597	14	9.34	19	10.00	22	7,063	13	11.8	13
North Dakota	32	47	6.18	34	5.38	40	342	48	10.7	37
Ohio	1,182	7	13.17	13	12.67	15	12,834	7	10.9	33
Oklahoma	116	37	4.25	46	4.12	47	1,414	35	12.2	9
Oregon	313	21	11.49	16	11.20	18	3,354	22	10.7	37
Pennsylvania	6,182	1	62.50	2	51.00	2	58,404	1	9.4	49
Rhode Island	70	42	8.26	24	7.28	28	763	42	10.9	33
South Carolina	270	24	8.42	22	9.00	24	3,213	24	11.9	12
South Dakota	19	50	3.20	49	2.77	50	230	50	12.1	10
Tennessee	458	17	10.08	17	10.56	19	5,307	17	11.6	17
Texas	1,308	4	8.19	26	10.14	21	16,095	4	12.3	7
Utah	203	29	12.27	15	17.31	9	2,025	31	10.0	47
Vermont	36	46	7.30	30	7.42	27	383	47	10.6	41
Virginia	1,754	3	31.08	4	35.73	4	19,984	3	11.4	22
Washington	675	11	14.49	10	15.95	10	6,828	15	10.1	45
West Virginia	2,921	2	197.26	1	172.40	1	32,244	2	11.0	30
Wisconsin	247	26	5.80	37	5.61	38	2,894	26	11.7	14
Wyoming	79	41	20.10	7	22.48	6	723	44	9.2	51

NOTE: See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Census Bureau.