

Hypersensitivity pneumonitis: Number of deaths, death rates (per million population), and years of potential life lost (YPLL) by state, U.S. residents age 15 and over, 1995–2004

State	No. of Deaths		Death Rate				YPLL to Life Expectancy			
	Deaths	Rank	Crude	Rank	Age-Adjusted	Rank	Total	Rank	YPLL/death	Rank
Alabama	4	32	0.11	41	0.11	41	60	33	15.1	22
Alaska	-	-	-	-	-	-	-	-	-	-
Arizona	7	21	0.18	27	0.17	30	152	16	21.7	5
Arkansas	1	43	0.05	46	0.04	47	15	43	14.6	26
California	55	1	0.21	22	0.23	19	1,005	1	18.3	12
Colorado	8	17	0.24	17	0.27	16	149	17	18.7	11
Connecticut	6	25	0.22	20	0.21	23	127	21	21.2	7
Delaware	-	-	-	-	-	-	-	-	-	-
District of Columbia	1	43	0.21	22	0.23	19	9	46	8.6	46
Florida	19	10	0.15	32	0.12	38	276	10	14.5	27
Georgia	1	43	0.02	48	0.02	48	9	46	8.6	46
Hawaii	-	-	-	-	-	-	-	-	-	-
Idaho	9	14	0.91	5	0.97	4	129	20	14.3	29
Illinois	18	11	0.18	27	0.19	27	242	11	13.4	35
Indiana	8	17	0.17	30	0.17	30	114	24	14.2	30
Iowa	25	6	1.07	2	0.91	5	284	9	11.4	39
Kansas	9	14	0.43	13	0.43	12	187	13	20.8	8
Kentucky	5	29	0.15	32	0.15	34	115	23	23.1	3
Louisiana	4	32	0.11	41	0.12	38	96	27	24.1	2
Maine	5	29	0.48	9	0.44	11	61	32	12.2	37
Maryland	7	21	0.17	30	0.18	28	116	22	16.6	18
Massachusetts	7	21	0.14	36	0.13	36	72	30	10.2	44
Michigan	24	8	0.31	14	0.31	14	337	7	14.0	33
Minnesota	25	6	0.64	7	0.64	6	300	8	12.0	38
Mississippi	2	39	0.09	43	0.09	43	45	38	22.7	4
Missouri	9	14	0.20	24	0.20	25	139	19	15.4	21
Montana	1	43	0.14	36	0.13	36	8	48	8.4	48
Nebraska	6	25	0.44	12	0.42	13	82	28	13.7	34
Nevada	2	39	0.13	39	0.17	30	56	35	28.1	1
New Hampshire	2	39	0.20	24	0.21	23	28	41	14.2	30
New Jersey	6	25	0.09	43	0.09	43	100	26	16.7	17
New Mexico	1	43	0.07	45	0.07	45	14	45	14.4	28
New York	36	3	0.24	17	0.23	19	512	5	14.2	30
North Carolina	18	11	0.28	16	0.30	15	242	11	13.4	35
North Dakota	3	37	0.58	8	0.45	10	27	42	9.0	45
Ohio	21	9	0.23	19	0.23	19	378	6	18.0	13
Oklahoma	4	32	0.15	32	0.14	35	68	31	17.1	16
Oregon	13	13	0.48	9	0.46	9	145	18	11.2	40
Pennsylvania	30	5	0.30	15	0.27	16	538	4	17.9	14
Rhode Island	1	43	0.12	40	0.11	41	15	43	14.9	23
South Carolina	6	25	0.19	26	0.20	25	105	25	17.5	15
South Dakota	4	32	0.67	6	0.60	8	60	33	14.9	23
Tennessee	8	17	0.18	27	0.18	28	166	14	20.8	8
Texas	35	4	0.22	20	0.26	18	573	3	16.4	19
Utah	8	17	0.48	9	0.63	7	159	15	19.9	10
Vermont	5	29	1.01	4	1.03	3	54	36	10.8	43
Virginia	3	37	0.05	46	0.06	46	47	37	15.7	20
Washington	7	21	0.15	32	0.16	33	78	29	11.1	41
West Virginia	2	39	0.14	36	0.12	38	43	40	21.6	6
Wisconsin	55	1	1.29	1	1.25	1	810	2	14.7	25
Wyoming	4	32	1.02	3	1.12	2	44	39	11.0	42

- indicates no deaths listed. NOTE: See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Census Bureau.