

Idaho, 2005

Primary and secondary syphilis

- **CASES:** Idaho reported 20 cases in 2005.
- **RATES:** Idaho ranked 30 among 50 states + DC + 3 territories: 1.4 cases per 100,000 population compared to the U.S. rate of 3.0 cases per 100,000 population.

Male-female rates of primary and secondary syphilis

- In Idaho, the rate among males was 2.0 per 100,000 population compared to the U.S. male rate of 5.1 per 100,000.
- The rate among females was 0.9 per 100,000 compared to the U.S. female rate of 0.9 per 100,000.

Racial disparities of primary and secondary syphilis


In Idaho, the race/ethnicity adjusted rates per 100,000 population were 0.3 among whites, 0.0 among blacks, 12.9 among Hispanics, 0.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.

- The rate among Hispanics was 43 times that of whites.

Congenital syphilis

- **CASES:** Idaho reported 0 cases in 2005.
- **RATES:** Idaho had a rate of 0.0 cases per 100,000 live births compared to the U.S. rate of 8.0 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Idaho, 1996-2005



Total P&S Syphilis Rates by County, Idaho, 2005

