Illinois, 2005

Primary and secondary syphilis

- CASES: Illinois reported 525 cases in 2005.
- RATES: Illinois ranked 9 among 50 states + DC + 3 territories: 4.1 cases per 100,000 population compared to the U.S. rate of 3.0 cases per 100,000 population.

Male-female rates of primary and secondary syphilis

- In Illinois, the rate among males was 7.7 per 100,000 population compared to the U.S. male rate of 5.1 per 100,000.
- The rate among females was 0.7 per 100,000 compared to the U.S. female rate of 0.9 per 100,000.

Racial disparities of primary and secondary syphllis


In Illinois, the race/ethnicity adjusted rates per 100,000 population were 2.9 among whites, 11.2 among blacks, 3.3 among Hispanics, 1.0 among Asian/Pacific Islanders, and 0.0 among American Indians/Alaska Natives.

- The rate among blacks was 3.9 times that of whites.
- The rate among Hispanics was 1.1 times that of whites.

Congenital syphilis

- CASES: Illinois reported 23 cases in 2005.
- RATES: Illinois ranked 12 among 28 states + 2 territories reporting at least one congenital syphilis case: 12.6 cases per 100,000 live births compared to the U.S. rate of 8.0 cases per 100,000.

Syphilis: P&S Rates and Congenital Cases, Illinois, 1996-2005



Total P&S Syphilis Rates by County, Illinois, 2005



Rate (per 100,000 population)

