Average Length of Stay in Short-Stay Hospitals:

Demographic, Diagnostic, and Surgical Statistics United States, 1977

Statistics are presented on the utilization of short-stay hospitals based on data collected by means of the National Hospital Discharge Survey from a national sample of hospital records of discharged patients. Average length of stay is presented by patient characteristics of age, sex, marital status, color, and diagnostic and surgical status in conjunction with hospital characteristics of size, ownership or control, and geographic region in which located.

PORPTY OF THE PUSHICATIONS APATION DO LOS TONS APATIONS APATION OF THE PUBLICATION OF THE

DHEW Publication No. (PHS) 81-1711

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service
Office of Health Research, Statistics, and Technology
National Center for Health Statistics
Hyattsville, Md. January 1981



Library of Congress Cataloging in Publication Data

Lewis, W. Frank

Average length of stay in short-stay hospitals, United States, 1977.

(Vital and health statistics: Series 13, Data from the National Health Survey; no. 50) (DHHS publication; (PHS) 81-1711)

Includes bibliographical references.

1: Hospital utilization—United States—Length of stay—Statistics. 2. United States—Statistics, Medical. I. Title. II. Series: United States. National Center for Health Statistics. Vital and health statistics: Series 13, Data from the National Health Survey, Data from the hospital discharge survey; no. 50. III. Series: United States. Dept. of Health and Human Services. DHHS publication; no. (PHS) 81-1711.

RA407.3.A349

no. 50

[RA981.A2]

362.1′1′0973s

ISBN 0-8406-0198-0

[362.1'1'0973]

80-29464

NATIONAL CENTER FOR HEALTH STATISTICS

DOROTHY P. RICE, Director

ROBERT A. ISRAEL, Deputy Director

JACOB J. FELDMAN, Ph.D., Acting Associate Director for Analysis and Epidemiology
GAIL F. FISHER, Ph.D., Acting Associate Director for the Cooperative Health Statistics System
GARRIE J. LOSEE, Acting Associate Director for Data Processing and Services
ALVAN O. ZARATE, Assistant Director for International Statistics
E. EARL BRYANT, Acting Associate Director for Interview and Examination Statistics
ROBERT C. HUBER, Acting Associate Director for Management
MONROE G. SIRKEN, Ph.D., Acting Associate Director for Research and Methodology
PETER L. HURLEY, Acting Associate Director for Vital and Health Care Statistics
ALICE HAYWOOD, Information Officer

DIVISION OF HEALTH CARE STATISTICS

W. EDWARD BACON, Ph.D., Acting Director

W. EDWARD BACON, Ph.D., Acting Chief, Hospital Care Statistics Branch
JAMES E. DELOZIER, Chief, Ambulatory Care Statistics Branch
MANOOCHEHR K. NOZARY, Chief, Technical Services Branch
STEWART C. RICE, Chief, Family Planning Statistics Branch
JOSEPH P. BARBANO, Chief, Health Professions Statistics Branch
EVELYN S. MATHIS, Chief, Long-Term Care Statistics Branch

COOPERATION OF THE U.S. BUREAU OF THE CENSUS

Under the legislation establishing the National Health Survey, the Public Health Service is authorized to use, insofar as possible, the services or facilities of other Federal, State, or private agencies.

In accordance with specifications established by the National Center for Health Statistics, the Bureau of the Census, under a contractual arrangement, participated in planning the survey and collecting the data.

Vital and Health Statistics-Series 13-No. 50

DHEW Publication No. (PHS) 81-1711
Library of Congress Catalog Card Number 80-29464

CONTENTS

	roduction	1 1
	Scope and Plan of Report	1
	erage Length of Stay	2 2 5 7
	Prage Length of Stay by Geographic Region	7 7 7
Rat	tio Comparison by Selected Characteristics	9
Con	nclusion	10
Ref	ferences	11
List	t of Detailed Tables	12
	pendixes I. Technical Notes on Methods II. Definitions of Certain Terms Used in This Report	35 42
	LIST OF TEXT FIGURES	
1.	Percent distribution of patients discharged from short-stay hospitals with first-listed diagnoses, single-listed diagnoses, and single-listed surgery, by age, color, and sex: United States, 1977	4
2.	Average length of stay for patients discharged from short-stay hospitals with first-listed diagnoses, single-listed diagnoses, and single-listed surgery, by age, color, and sex: United States, 1977	e
3.	Percent of patients discharged from short-stay hospitals, by type of ownership of hospital and geographic region: United States, 1977	8
	LIST OF TEXT TABLES	
Α.	Number of discharges and average length of stay for patients discharged from short-stay hospitals with first-listed diagnoses, single-listed diagnoses, and single-listed surgery, by sex and age: United States, 1977	5
В.	Average length of stay for patients discharged from short-stay hospitals with first-listed diagnoses, single-listed diagnoses, and single-listed surgery, by geographic region, bed size of hospital, age, and sex: United States, 1977	1
C.	Percent distribution of patients discharged from short-stay hospitals with first-listed diagnoses, single-listed diagnoses, and single-listed surgery, by geographic region, bed size of hospital, age,	

AVERAGE LENGTH OF STAY IN SHORT-STAY HOSPITALS: DEMOGRAPHIC, DIAGNOSTIC, AND SURGICAL STATISTICS

W. Frank Lewis, M.A., Division of Health Care Statistics

INTRODUCTION

Background of Study

This report presents a special analysis of the average length of stay in short-stay hospitals by using data collected in the National Hospital Discharge Survey, a continuous nationwide survey conducted by the National Center for Health Statistics. Data were abstracted from approximately 224,000 hospital records of inpatients discharged from the 423 hospitals that participated in the survey in 1977.

Data from this survey provide input for three basic types of reports—nonmedical, diagnostic, and surgical—that are published both in Series 13 of Vital and Health Statistics and as selected supplements of Monthly Vital Statistics Reports. Nonmedical data have been published for 1965-78; diagnostic data for 1965-68 and 1971-78; and surgical data on operations performed in 1965, 1968, and 1971-78. In addition, one report has been published on a study of charges of care in short-stay hospitals that was based on data collected by the National Hospital Discharge Survey in 1968-70.

Information on hospitalization is collected by means of the National Health Interview Survey, another program of the National Center for Health Statistics. The estimates provided by the National Health Interview Survey are usually lower for the number of discharges and greater for the average length of stay than the National Hospital Discharge Survey estimates because of differences in collection procedures, definitions, and population sampled. The National Health Interview Survey data are published in Series 10 reports. Data used in this report include a classification of discharges by age, sex, race, marital status, bed size of hospital and hospital ownership, and geographic region. Diagnostic and surgical data include only those discharges with first-listed diagnosis (all patients), single-listed diagnosis, and single-listed surgery.

Definitions of the terms used in this report relating to hospitalization and characteristics of the inpatients are given in appendix II. Because several terms used in this report have specialized meanings, familiarity with these definitions will be useful in interpreting the data. Appendix I provides a description of the survey design, data collection procedures, and the estimation process. A detailed report on the design of the National Hospital Discharge Survey has previously been published.¹

Scope and Plan of Report

Average length of stay is an important means by which comparisons can be made in hospital utilization. Differences in length of stay between hospitals can be related to their size, ownership, or geographical location because these variables reflect differences in availability of services. In addition to these factors, differences among hospitals can be related to the characteristics of the patients who are discharged from them. Finally, length-of-stay differences among hospitals can be related to the number and kinds of diagnoses and to the surgical procedures that these hospitals employ.

Prior studies by the National Hospital Discharge Survey (NHDS) have shown that the average length of stay is influenced by these factors in a predictable and consistent manner. Specifically, NHDS data have indicated the following about length of stay:

- 1. Length of stay is lowest for patients under 15 years of age and increases steadily with age.
- 2. Length of stay for males, overall, is higher than that for females, except for patients 65 years of age and over where it is higher for females.
- 3. White patients have a shorter average length of stay than patients in the category classified as "all other" have.
- 4. Patients discharged from small hospitals generally have the shortest length of stay, and patients discharged from the largest hospitals have the longest length of stay.
- 5. Voluntary nonprofit hospitals have the longest average length of stay, and proprietary hospitals have the shortest.
- 6. Unmarried patients over 15 years of age have longer lengths of stay than married patients have.
- 7. Average length of stay is consistently highest in the Northeast Region and lowest in the West Region.
- 8. Patients discharged with single-listed surgery have shorter length of stays than those with multiple surgery.
- 9. Patients discharged with a single-listed diagnosis have shorter stays than those with multiple diagnoses.

Interviews conducted during the National Health Interview Survey (NHIS) have shown similar length of stay patterns for age, sex, color, type of hospital, marital status, and region. No satisfactory answer has been given as to why the average length of stay in the West Region is always lower than that in the Northeast, North Central, or South Regions. This report describes the singular effects of patient, hospital, diagnostic, and surgical characteristics on the length of stay and analyzes their cumulative effect on length of stay, especially as they may affect regional differences.

AVERAGE LENGTH OF STAY

٠,

Age, Sex, and Color

The average length of stay for the 35.9 million patients discharged from short-stay hospitals with a first-listed diagnosis during 1977 was 7.3 days. Average length of stay increased with each successive age group from a low of 4.2 days for patients under 15 years of age to a high of 11.1 days for patients 65 years and over (table A). Males had an average length of stay of 7.8 days and females, including deliveries, had an average length of stay of 7.0 days. This difference is primarily due to the longer length of stay for males in the age group 15-44 years (6.3 days for males and 4.9 for females). Excluding deliveries, the length of stay for females increased to 7.6 days for all ages and 5.4 days for the age group 15-44 years. This increase is caused by the removal of relatively short stays associated with deliveries, 99 percent of which occur in the age group 15-44 years.

There was no significant overall difference in the average length of stay by color, with white patients averaging 7.3 days and all other patients averaging 7.6 days per stay (table 1). The average length of stay for the total of both color groups was about the same because the white discharged patients included a larger percent of older patients with longer hospital stays than the all other group did (figure 1). However, for every age and sex group, except females aged 15-44 years, the average stay for white patients was shorter than the average stay for all other patients. Regardless of color, males overall had longer stays than females had.

All other males in the age group 45-64 years had an average length of stay that was 3.1 days

Table A. Number of discharges and average length of stay for patients discharged from short-stay hospitals with first-listed diagnoses, single-listed diagnoses, and single-listed surgery, by sex and age: United States, 1977

•	D		Fer	male		
Type of diagnosis or surgery and age	Both sexes	Male	Including deliveries	Excluding deliveries		
First-listed diagnosis	Number of discharges in thousands					
All ages	35,902	14,385	21,518	18,186		
Under 15 years	3,775 15,180 8,604 8,344	2,137 4,553 4,042 3,653	1,638 10,627 4,562 4,690	1,624 7,315 4,557 4,690		
<i>,</i>	ļ <i>f</i>	Average le	ngth of stay in	n days		
All ages	7.3	7.8	. 7.0	7.6		
Under 15 years	4.2 5.3 8.5 11.1	4.3 6.3 8.6 10.8	4.1 4.9 8.4 11.3	4.1 5.4 8.4 11.3		
Single-listed diagnosis	Nu	mber of d	ischarges in th	ousands		
All ages	18,627	7,025	11,602	8,733		
Under 15 years	2,474 10,208 3,652 2,293	1,404 2,822 1,765 1,034	1,070 7,386 1,886 1,260	1,058 4,533 1,882 1,260		
	Δ	verage ler	ngth of stay in	days		
All ages	5.4	5.9	5.1	5.5		
Under 15 years	3.7 4.5 6.9 8.7	3.9 5.4 6.9 8.2	3.6 4.1 7.0 9.1	3.6 4.4 7.0 9.1		
Single-listed surgery	Nun	nber of di	scharges in the	ousands		
All ages	10,227	4,093	6,133	5,280		
Under 15 years	1,202 4,847 2,350 1,827	682 1,441 1,115 855	520 3,406 1,235 972	516 2,558 1,233 972		
	A A	verage len	gth of stay in	days		
All ages	6.6	7.2	6.2	6.4		
Under 15 years	3.7 4.7 8.0	3.8 5.6 8.4	3.4 4.4 7.7	3.4 4.2 7.7		
65 years and over	11.7	11.1	12.2	12.2		




Figure. 1. Percent distribution of patients discharged from short-stay hospitals with first-listed diagnoses, single-listed diagnoses, and single-listed surgery, by age, color, and sex: United States, 1977

longer than that for white males in the same age group. The difference between the length of stay of all other females and white females in this age group was only 1.3 days.

The percent distribution of white males discharged in the age groups over 14 years was almost evenly distributed among the age groups 15-44 years, 45-64 years, and 65 years and over—30.8, 28.8, and 26.2 percent, respectively (figure 1). The proportion of male discharges in the all other group was largest for ages 15-44 (37.9 percent), followed by 23.8 for ages 15-64 and 18.2 for ages 65 and over. Females in the all other group accounted for proportionately more discharges at ages 15-44 than white females of the same ages (64.0 compared with 47.0 percent) and proportionately less at ages over 44.

Patients discharged with only a single-listed diagnosis totaled 18.6 million and had an average length of stay of 5.4 days. The average length of stay for males was longer than that for females at all ages and longest at ages 15-44 (table A). In figure 2, the length of stay is shown by sex and color. Males with a single-listed diagnosis in both color categories, showed an increase in the length of stay with each succeeding age group. However, female patients in the age group 15-44 years did not have a significantly longer length of stay than those under 15 years of age, primarily because of the inclusion of deliveries. The average length of stay for both white and all other patients was 5.4 days. The all other group showed longer stays for each age group except ages 15-44 years, primarily because the length of stay for both all other and white females was 4.1 days (table 2).

Patients discharged with single-listed surgery totaled 10.2 million and had an average length of stay of 6.6 days (table A). This was less than that for all discharges (7.3 days), but more than that for discharges with a single-listed diagnosis (5.4 days). The average stay for males (7.2 days) was longer than that for females (6.2 days) for the total of all ages. Stays for females aged 15-44 years were significantly greater than for those under 15 years of age, because deliveries are not counted as operations. Patients with single-listed surgery had 47.4 percent of their discharges in the age group 15-44 years (figure 1). For white males, the proportion of discharges in the age

group under 15 years was 15.9 percent, and for all other males, it was 22.4 percent.

Table 3 shows the average length of stay for discharges with a single-listed surgery. For both color groups the single-listed surgery discharges in all age and sex groups included a smaller percent in the age group 65 years and over than those with a first-listed diagnosis did, but included a larger percent than those with a single-listed diagnosis did (figure 1). Females in the all other category with single-listed surgery had a greater percent of discharges (71.2 percent) in the age group 15-44 years than those discharged with a first-listed diagnosis (64.0 percent).

Bed Size of Hospital and Hospital Ownership

The average length of stay increased as age or hospital bed size increased, ranging from 6.0 days in the smallest hospitals to 8.4 days in the largest hospitals and from 4.2 days in the youngest group to 12.4 days for the age group 65 years and over who were discharged from hospitals with 500 beds or more (table 4). Male discharges from hospitals with 6-99 beds averaged 6.1 days per stay. The stay increased to 9.3 days in hospitals with 500 beds or more. Females showed a significantly smaller increase in stay by bed size of hospital, from 6.0 days (6-99 beds) to 7.8 days (500 beds or more).

Length of stay for single-listed diagnosis showed differences between the youngest and oldest age group of only 5.0 days (3.7 and 8.7 days, respectively) and between the smallest and the largest hospitals of 1.8 days (table 5). Discharges with only a single-listed surgery showed a length of stay for the age group 65 years and over to be 3.2 times greater than that for the age group under 15 years (table 6).

The average length of stay was shorter in government and proprietary hospitals than in voluntary nonprofit hospitals (table 7). This difference was due to the longer stays for males (8.0 days) and for the age groups 45 years and over discharged from nonprofit hospitals. In government hospitals, for the age group 15-44 years the average length of stay for males was 2.0 days longer than it was for females.

For patients discharged with a single-listed




Figure 2. Average length of stay for patients discharged from short-stay hospitals with first-listed diagnoses, single-listed diagnoses, and and single-listed surgery, by age, color, and sex: United States, 1977

diagnosis, voluntary, government, and proprietary hospitals showed stays that were not significantly different (table 8). Females discharges from proprietary hospitals had a length of stay of 5.1 days, which was not significantly different from male discharges (5.5 days). Table 9 shows persons discharged with a single-listed surgery.

Marital Status

For all discharges 15 years of age and over, unmarried persons had longer stays (8.7 days) than married persons (7.1 days) (table 10). However, for discharges aged 15-44 years, there was little significant difference between married and unmarried discharged persons. In hospitals with less than 100 beds, no significant difference existed for this age group. In government and voluntary hospitals, the average length of stay was 1.6 and 1.7 days longer, respectively, for unmarried discharged persons than for married persons (table 11). In voluntary hospitals, the average length of stay of 8.6 days for discharged married males exceeded that for married females (6.4 days) by 2.2 days. However, unmarried females discharged from voluntary hospitals had stays similar to those of unmarried males. Regardless of hospital ownership, unmarried females had longer stays than married females.

Table 12 shows the average length of stay for persons discharged with a single-listed diagnosis by marital status. The difference between the length of stay of married and unmarried discharges 15 years and over was 1.0 days. However, for persons aged 15-44 years discharged from hospitals with less than 100 beds, there was no significant difference between the married and unmarried.

The difference between stays of married and unmarried discharges with a single-listed surgery was 1.1 days (table 13). For ages 15-44 years, length of stay was about the same regardless of marital status or hospital size.

AVERAGE LENGTH OF STAY BY GEOGRAPHIC REGION

Age and Sex

Average length of stay was highest in the Northeast Region and lowest in the West Region,

8.6 days and 6.0 days, respectively (table 14). This regional pattern was consistent for all age groups, with the difference between the Northeast and the West Regions being greatest in the age group 65 years and over, 13.6 and 8.9 days, respectively, or a difference of 4.7 days. Length of stay for both sexes was longest in the Northeast Region for the age group 65 years and over, and shortest in the West Region for the age group under 15 years. The average length of stay for males was longer than that for females in each of the four Regions.

For patients discharged with a single-listed diagnosis, the Northeast Region had a length of stay of 6.1 days, or 1.7 days longer than that for discharges in the West Region (4.4 days). Length of stay did not differ significantly among the Northeast, North Central, and South Regions for persons discharged under 15 years of age with a single-listed diagnosis (table 15) or with a single-listed surgery (table 16).

Bed Size of Hospital and Hospital Ownership

Average length of stay by bed size of hospital, region, age, and sex is shown in table 17. For each region, length of stay increased with bed size and age; it was shortest in hospitals with less than 100 beds for discharges under 15 years of age, and longest in hospitals with 500 beds or more for discharges 65 years of age and over. Average stays were longest in the Northeast Region for persons 65 years of age and over discharged from hospitals with 500 beds or more. The West Region consistently had a shorter average length of stay for each hospital size and age group than the Northeast Region did. Length of stay for hospitals with 500 beds or more in the Northeast Region was only 28.9 percent longer than that for hospitals with less than 100 beds; however, in the West Region, the comparative difference was 40.8 percent.

Patients discharged with a single-listed diagnosis showed length of stay for hospitals with 500 beds or more (6.7 days) in the Northeast Region to be only 13.6 percent greater than that for hospitals with less than 100 beds (5.9 days) (table B). In the West Region, the difference between the largest and smallest hospitals was 43.2 percent (5.3 and 3.7 days, respectively). Females discharged under 15 years of age from

Table B. Average length of stay for patients discharged from short-stay hospitals with first-listed diagnoses, single-listed diagnoses, and single-listed surgery, by geographic region, bed size of hospital, age, and sex: United States, 1977

		Northeast			North Central			South			West		
Type of diagnosis or surgery, age, and sex	AII regions	6-99 beds	100- 499 beds	500 beds or more	6-99 beds	100- 499 beds	500 beds or more	6-99 beds	100- 499 beds	500 beds or more	6-99 beds	100- 499 beds	500 beds or more
First-listed diagnosis		Average length of stay in days											
All ages, both sexes ¹	7.3	7.6	8.5	9.8	6.1	7.4	8.4	6.0	6.8	8.1	4.9	6.3	6.9
Under 15 years 15-44 years 45-64 years 65 years and over	4.2 5.3 8.5 11.1	3.2 5.6 8.1 11.8	4.3 5.6 9.7 13.5	6.0 6.8 12.0 15.5	3.4 4.6 6.7 8.5	4.0 5.5 8.9 11.6	5.4 6.6 9.8 12.1	3.3 4.3 6.4 9.1	4.0 4.8 8.3 10.8	5.6 6.3 9.6 11.7	2.8 3.7 5.4 7.8	3.8 4.8 7.2 9.1	5.6 5.4 7.7 9.8
Single-listed diagnosis	5.4	5.9		6.7	4.5	5.4	6.5	4.5	5.0	6.2	3.7	4.6	5.3
All ages, both sexes ¹	5.4	5.9	6.0	6.7	4.5	5.4	0.5	4.5	<u> </u>				
Under 15 years	3.7 4.5 6.9 8.7	2.8 5.0 6.8 11.2	3.9 4.8 7.9 10.8	5.1 5.5 8.6 11.1	3.1 3.8 5.4 6.8	3.6 4.7 7.2 8.6	4.6 5.4 8.1 9.9	3.1 3.8 5.4 7.5	3.6 4.1 6.7 8.5	5.1 5.2 7.7 9.2	2.6 3.2 4.7 5.4	3.4 3.9 5.8 7.0	4.2 4.2 6.4 8.0
Single-listed surgery All ages, both sexes 1	6.6	5.9	7.6	8.3	5.5	6.4	7.5	5.5	6.2	7.6	4.3	5.8	6.5
Under 15 years	3.7 4.7 8.0 11.7	2.4 4.0	3.4 5.0 9.0 13.8	5.3 5.7 9.9 14.7	2.6 4.0 6.5 10.6	3.4 4.6 7.9 11.4	4.7 5.7 8.9 11.8	2.9 4.1 7.0 11.2	3.4 4.5 7.7 11.9	5.5 5.6 9.2 12.1	2.1 3.6 5.4 7.2	3.4 4.3 6.8 9.3	5.2 5.3 6.9 9.6

¹Includes discharge data for which sex was not stated.

hospitals with less than 100 beds had stays that did not significantly differ among the Northeast, North Central, and South Regions (table 18).

Average stays for single-listed surgery discharges in the Northeast Region were not always higher than comparable stays in the West Region (table 19). This statement is especially true in the age group 15-44 years where males discharged from small hospitals showed similar stays for the Northeast and West Regions (4.1 and 4.0 days, respectively). Female patients in the age group 15-44 years (excluding deliveries) in the West Region had stays similar to those of females discharged in the Northeast Region.

Table 11 shows the regional length of stay by type of hospital ownership, age, sex, and marital status for patients discharged. In government-owned hospitals, average stays for male patients (7.4 days) were 1.1 days longer than those for female patients (6.3 days). In contrast, in proprietary hospitals the stays for males were no longer than those for females. In




Figure 3. Percent of patients discharged from short-stay hospitals, by type of ownership of hospital and geographic region: United States, 1977

Table C. Percent distribution of patients discharged from short-stay hospitals with first-listed diagnoses, single-listed diagnoses, and single-listed surgery, by geographic region, bed size of hospital, age, and sex: United States, 1977

		Northeast			North Central			South			West		
Type of diagnosis or surgery, age, and sex	All regions	6-99 beds	100- 499 beds	500 beds or more	6-99 beds	100- 499 beds	500 beds or more	6-99 beds	100- 499 beds	500 beds or more	6-99 beds	100- 499 beds	500 beds or more
First-listed diagnosis		Percent distribution											
All ages, both sexes ¹	100.0	10.8	71.3	18.0	12.8	60.4	26.9	30.3	48.3	21.4	24.8	64.6	10.6
Under 15 years 15-44 years 45-64 years 65 years and over	100.0 100.0 100.0 100.0	7.7 10.8 12.2 10.5	73.8 69.5 70.5 74.1	18.7 19.7 17.4 15.4	10.4 11.4 12.6 16.8	63.6 61.0 58.3 59.7	26.1 27.5 29.1 23.6	28.4 28.0 29.6 35.9	51.3 49.4 48.1 45.2	20.2 22.6 22.3 19.0	25.6 25.0 24.0 25.1	63.4 64.2 64.7 65.8	11.0 10.8 11.3 9.1
Single-listed diagnosis													
All ages, both sexes ¹	100.0	10.7	70.3	19.0	11.2	60.4	28.5	29.1	48.7	22.2	25.4	63.9	10.7
Under 15 years	100.0 100.0 100.0 100.0	8.1 10.7 13.0 9.7	73.3 69.7 68.8 72.1	18.4 19.6 18.2 18.5	10.5 10.5 11.1 15.1	63.5 61.7 57.1 56.3	26.1 27.9 31.9 28.6	29.9 28.0 28.8 34.3	51.2 49.6 47.6 43.8	18.9 22.5 23.7 21.8	28.7 25.6 23.6 24.6	62.3 64.1 64.9 62.9	9.0 10.3 11.5 12.4
Single-listed surgery All ages, both sexes ¹	100.0	5.9	73.6	20.5	8.4	59.4	32.2	19.0	53.9	27.1	20.6	68 .5	11.0
Under 15 years	100.0 100.0 100.0 100.0	6.3 6.6 5.3 4.8	72.7 72.3 74.3 76.3	21.4 21.1 20.6 18.6	9.0 8.7 7.3 8.5	62.9 59.6 56.0 60.9	28.1 31.7 36.7 30.4	23.1 19.6 16.3 17.8	53.0 55.1 53.3 52.4	24.2 25.4 30.4 30.0	27.1 22.2 18.1 15.9	63.3 67.2 70.0 72.9	10.0 10.6 12.1 11.2

¹Includes discharge data for which sex was not stated.

both the voluntary and government-owned hospitals in each of the four regions, unmarried discharged persons had longer stays than married persons, especially in the Northeast Region. Overall, the number of discharges from government hospitals in the Northeast Region comprised the smallest percent of total discharges (7.4 percent) for any of the four regions (figure 3). The distribution of discharges by ownership in the South and West Regions showed a smaller percent of their discharges from voluntary nonprofit hospitals, 54.2 and 57.6 percent, respectively, in contrast to the Northeast and North Central Regions (86.4 and 82.4 percent, respectively). The South, like the West, when compared with the Northeast and North Central Regions, had a relatively larger percent of its discharges from smaller hospitals than from the largest ones. Hospitals with less than 100 beds in the South and West Regions comprised 30.3 and 24.8 percent, respectively, of patient discharges, compared with the Northeast and North

Central Regions where they comprised 10.8 and 12.8 percent, respectively, of patient discharges (table C).

RATIO COMPARISONS BY SELECTED CHARACTERISTICS

Table D shows the ratio of the average length of stay in the Northeast Region to those in the North Central, South, and West Regions by selected variables. By comparing the Northeast Region with the West Region by age, the average length of stay for ages 15-44 years in the Northeast Region is shown to be approximately 1.26 days for each 1-day stay in the West Region, compared with 1.43 days for each 1-day stay for the total of all ages. Similarly, by comparing hospital size, the ratio for hospitals with 100-499 beds is only 1.34 days; by ownership (voluntary), 1.39 days; and by marital status

Table D. Ratios of average length of stay in the Northeast Region to stays in the North Central, South, and West Regions, by selected characteristics: United States, 1977

Characteristic	North Cent.al	South	West			
	Ratio					
Total	1.15	1.24	1.43			
Age: Under 15 years	1.05	1.10	1.22			
15-44 years	1.02	1.16	1.26			
45-64 years65 years and over	1.11 1.21	1.24 1.31	1.46 1.53			
Hospital size: 6-99 beds	1.24	1.27	1.55			
100-499 beds	1.15	1.27	1.55 1.34			
500 beds or more	1.17	1.21	1.42			
Hospital ownership: Voluntary	1.10	1.20	1.39			
Government	1.40	1.44	1.54			
Proprietary	0.86	1.38	1.67			
Sex: Male	1.16	1.26	1.41			
Female	1.14	1.24	1.43			
Marital status:						
Married Unmarried	1.11 1.18	1.22 1.28	1.41 1.49			
Color:						
White	1.13	1.26	1.41			
All other	1.12	1.32	1.36			
Voluntary nonprofit hospitals, 100-499 beds, unmarried,						
ages 15-44 years ¹	0.98	1.15	1.18			
discharges)	1.08	1.20	1.39			
Single-listed surgery (all discharges)	1.13	1.17	1.38			

¹SOURCE: National Center for Health Statistics, Division of Health Resources Statistics: Unpublished data from the 1977 Hospital Discharge Survey.

(married), 1.41 days. Because each of the preceding characteristics shows the Northeast-to-West stay ratio to be less than or similar to the total ratio (1.43 days), the combination of these characteristics would yield a smaller ratio than any single characteristic. This situation occurs when stays for married patients, ages 15-44, discharged from voluntary hospitals with 100-499 beds in the Northeast Region are compared with similar discharges in the West Region. The average length-of-stay ratio is only 1.18 days, com-

pared with the total regional ratio of 1.43 days.

For discharges with a single-listed diagnosis, the Northeast-to-West length-of-stay ratio was 1.39 days (table D). The average stay in the Northeast Region for patients under 15 years of age discharged from hospitals with less than 100 beds was 2.8 days, and in the West Region, the stay was 2.6 days, or a ratio of 1.08 days (table 18).

The comparison of average stays for single-listed surgery discharges by region shows a significant reduction in regional differences by selected characteristics (table 19). Female discharges, ages 15-44, from hospitals with 500 beds or more had stays of 5.1 and 4.5 days, respectively, in the Northeast and West Regions, or a ratio of only 1.13 days. The same comparison for females (excluding deliveries) showed stays of 4.7 and 4.8 days, respectively, or a ratio of 0.98 days. The stays for the Northeast and South Regions for females (including deliveries) ages 15-44 years discharged from hospitals with 500 beds or more were both 5.1 days.

CONCLUSION

Analysis of average length of stay indicates the following:

- Only two characteristics—age and hospital size—are relatively unaffected by interaction with other characteristics. In general, older patients in larger hospitals had the longest stays regardless of other characteristics.
- Color and hospital ownership are influenced by hospital size; marital status differences are influenced by sex, age, and number of diagnoses and surgeries; and sex differences vary with age and bed size.
- Regional differences generally are smallest in the age group under 15 years discharged from medium-sized hospitals, and greatest in the age group over 64 years discharged with a single-listed diagnosis from small hospitals.
- Regional differences in length of stay can to some extent be explained by dif-

ferences in hospital size and ownership. The Northeast Region is characterized by large voluntary hospitals, compared with the West Region where hospitals are characteristically smaller and composed of a higher percent of proprietary hospitals.

REFERENCES

¹National Center for Health Statistics: Development of the design of the NCHS Hospital Discharge Survey. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 2-No. 39. Public Health Service. Washington. U.S. Government Printing Office, Sept. 1970.

²National Center for Health Statistics: Development and maintenance of a national inventory of hospitals and institutions. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 1-No. 3. Public Health Service. Washington. U.S. Government Printing Office, Feb. 1965.

³National Center for Health Statistics: Eighth Revision International Classification of Diseases, Adapted for Use in the United States. 2 Vols. PHS Pub. No. 1693. Public Health Service. Washington. U.S. Government Printing Office, 1967 and 1968.

⁴National Center for Health Statistics: Utilization of short-stay hospitals, summary of nonmedical statistics, United States, 1965. *Vital and Health Statistics*. PHS Pub. No. 1000-Series 13-No. 2. Public Health Service. Washington. U.S. Government Printing Office, Aug. 1967.

⁵National Center for Health Statistics: Utilization of short-stay hospitals by characteristics of discharged patients, United States, 1965. Vital and Health Statistics. PHS Pub. No. 1000-Series 13-No. 3. Public Health Service. Washington. U.S. Government Printing Office, Dec. 1967.

LIST OF DETAILED TABLES

1.	Average length of stay and number of patients discharged from short-stay hospitals with a first-listed diagnosis, by sex, color, and age: United States, 1977
2.	Average length of stay and number of patients discharged from short-stay hospitals with a single-listed diagnosis, by sex, color, and age: United States, 1977
3.	Average length of stay and number of patients discharged from short-stay hospitals with a single-listed surgery, by sex, color, and age: United States, 1977
4.	Average length of stay and number of patients discharged from short-stay hospitals with a first-listed diagnosis, by sex, bed size of hospital, and age: United States, 1977
5.	Average length of stay and number of patients discharged from short-stay hospitals with a single-listed diagnosis, by sex, bed size of hospital, and age: United States, 1977
6.	Average length of stay and number of patients discharged from short-stay hospitals with a single-listed surgery, by sex, bed size of hospital, and age: United States, 1977
7.	Average length of stay and number of patients discharged from short-stay hospitals with a first-listed diagnosis, by sex, type of ownership of hospital, and age: United States, 1977
8.	Average length of stay and number of patients discharged from short-stay hospitals with a single-listed diagnosis, by sex, type of ownership of hospital, and age: United States, 1977
9.	Average length of stay and number of patients discharged from short-stay hospitals with a single-listed surgery, by sex, type of ownership of hospital, and age: United States, 1977
10.	Average length of stay for patients discharged from short-stay hospitals with a first-listed diagnosis, by color, marital status, age, geographic region, and bed size of hospital: United States, 1977
11.	Average length of stay and number of patients discharged with a first-listed diagnosis, by type of ownership of hospital, geographic region, age, sex, marital status, and sex and marital status: United States, 1977
12.	Average length of stay for patients discharged from short-stay hospitals with a single-listed diagnosis, by color, marital status, age, geographic region, and bed size of hospital: United States, 1977
13.	Average length of stay for patients discharged from short-stay hospitals with a single-listed surgery, by color, marital status, age, geographic region, and bed size of hospital: United States, 1977
14.	Average length of stay and number of patients discharged from short-stay hospitals with a first-listed diagnosis, by sex, geographic region, and age: United States, 1977
15.	Average length of stay and number of patients discharged from short-stay hospitals with a single-listed diagnosis, by sex, geographic region, and age: United States, 1977
16.	Average length of stay and number of patients discharged from short-stay hospitals with a single-listed surgery, by sex, geographic region, and age: United States, 1977
17.	Average length of stay for patients discharged from short-stay hospitals with a first-listed diagnosis, by geographic region, bed size of hospital, sex, and age: United States, 1977
18.	Average length of stay for patients discharged from short-stay hospitals with a single-listed diagnosis, by geographic region, bed size of hospital, sex, and age: United States, 1977
19.	Average length of stay for patients discharged from short-stay hospitals with a single-listed surgery, by geographic region, bed size of hospital sex and age: United States 1977

Table 1. Average length of stay and number of patients discharged from short-stay hospitals with a first-listed diagnosis, by sex, color, and age: United States, 1977

Color and age	Both sexes	Male	Female	Both sexes	Male	Female	
<u>Total</u>	Avera	Average length of stay Number of particles in days discharged in the					
All ages	7.3	7.8	7.0	35,902	14,385	21,518	
Under 15 years	4.2 5.3 8.5 11.1	4.3 6.3 8.6 10.8	4.1 4.9 8.4 11.3	3,775 15,180 8,604 8,344	2,137 4,553 4,042 3,653	1,638 10,627 4,562 4,690	
White							
All ages	7.3	7.7	7.1	27,342	11,115	16,226	
Under 15 years	4.1 5.3 8.4 11.0	4.2 6.2 8.4 10.7	4.1 4.9 8.4 11.2	2,802 11,055 6,797 6,688	1,575 3,429 3,199 2,912	1,227 7,626 3,598 3,776	
All other							
All ages	7.6	8.9	6.8	3,963	1,459	2,504	
Under 15 years	5.2 5.7 10.6 12.6	5.4 7.5 11.5 12.4	4.8 5.1 9.7 12.9	497 2,155 741 569	293 553 348 266	204 1,603 393 303	
Color not stated							
All ages	7.0	7.4	6.8	4,598	1,810	2,788	
Under 15 years	3.6 5.1 8.0 11.1	3.7 5.9 8.0 10.5	3.5 4.7 8.0 11.5	477 1,969 1,066 1,086	269 571 495 475	207 1,398 571 611	

Table 2. Average length of stay and number of patients discharged from short-stay hospitals with a single-listed diagnosis, by sex, color, and age: United States, 1977

Color and age	Both sexes	Male	Female	Both sexes	Male	Female	
Total				-	nber of patients arged in thousands		
All ages	5.4	5.9	5.1	18,627	7,025	11,602	
Under 15 years	3.7 4.5 6.9 8.7	3.9 5.4 6.9 8.2	3.6 4.1 7.0 9.1	2,474 10,208 3,652 2,293	1,404 2,822 1,765 1,034	1,070 7,386 1,886 1,260	
White All ages	5.4	5.9	5.1	13,960	5,379	8,581	
Under 15 years	3.7 4.5 6.9 8.6	3.8 5.4 6.8 8.1	3.6 4.1 7.0 9.1	1,838 7,415 2,886 1,821	1,039 2,135 1,395 810	800 5,280 1,491 1,011	
All other						:	
All ages	5.4	6.6	4.7	2,188	727	1,461	
Under 15 years	4.5 4.6 7.7 10.3	4.9 6.3 8.1 10.1	4.1 4.1 7.3 10.4	329 1,428 283 147	194 323 136 74	135 1,105 147 74	
Color not stated							
All ages	5.1	5.6	4.8	2,479	919	1,560	
Under 15 years	3.3 4.3 6.6 8.2	3.4 5.1 6.7 7.8	3.1 4.0 6.5 8.5	307 1,365 482 325	172 363 234 150	135 1,001 249 176	

Table 3. Average length of stay and number of patients discharged from short-stay hospitals with a single-listed surgery, by sex, color, and age: United States, 1977

Color and age	Both sexes	Male	Female	Both sexes	Male	Female		
<u>Total</u>	Avera				mber of patients arged in thousands			
All ages	6.6	7.2	6.2	10,227	4,093	6,133		
Under 15 years	3.7 4.7 8.0 11.7	3.8 5.6 8.4 11.1	3.4 4.4 7.7 12.2	1,202 4,847 2,350 1,827	682 1,441 1,115 855	520 3,406 1,235 972		
White								
All ages	6.7	7.3	6.3	7,739	3,153	4,587		
Under 15 years	3.6 4.8 7.9 11.7	3.8 5.6 8.1 11.3	3.3 4.4 7.7 12.0	899 3,540 1,859 1,442	502 1,096 882 673	396 2,444 977 769		
All other								
All ages	6.9	8.2	6.2	1,093	388	705		
Under 15 years	4.7 5.0 11.1 13.7	4.5 6.5 12.3 12.1	5.2 4.6 10.0 15.4	139 654 180 121	87 151 87 63	52 502 94 58		
Color not stated								
All ages	6.1	6.4	5.8	1,394	553	841		
Under 15 years	3.0 4.4 7.1 10.8	3.1 5.0 7.5 9.9	2.9 4.1 6.8 11.5	165 654 310 265	93 194 147 119	72 460 164 146		

Table 4. Average length of stay and number of patients discharged from short-stay hospitals with a first-listed diagnosis, by sex, bed size of hospital, and age: United States, 1977

Bed size of hospital and age	Both sexes	Male	Female	Both sexes	Male	Female	
All sizes	Avera	ge length in day:	of stay		Number of patients discharged in thousand		
All ages	7.3	7.8	7.0	35,902	14,385	21,518	
Under 15 years	4.2 5.3 8.5 11.1	4.3 6.3 8.6 10.8	4.1 4.9 8.4 11.3	3,775 15,180 8,604 8,344	2,137 4,553 4,042 3,653	1,638 10,627 4,562 4,690	
6-99 beds							
Ail ages	6.0	6.1	6.0	7,126	2,871	4,255	
Under 15 years	3.2 4.4 6.5 9.1	3.2 4.7 6.4 8.6	3.2 4.2 6.5 9.4	678 2,869 1,678 1,901	361 931 772 807	317 1,938 906 1,093	
100-199 beds All ages	6.8	7.2	6.5	6,097	2,375	3,722	
Under 15 years	3.9 4.7 8.0 10.6	3.9 5.8 8.1 10.0	3.8 4.3 7.9 11.1	690 2,591 1,370 1,447	388 730 634 624	302 1,861 736 823	
200-299 beds	7.0	7.0	7.0	E 556	2 224	2 222	
All ages	7.3	7.8	7.0	5,556	2,224	3,332	
Under 15 years	4.1 5.2 8.5 11.6	4.2 6.1 8.5 11.3	3.8 4.8 8.6 11.8	623 2,334 1,259 1,340	355 688 594 587	268 1,646 664 754	
300-499 beds							
All ages	7.7	8.2	7.4	9,761	3,931	5,830	
Under 15 years	4.2 5.6 9.1 12.0	4.2 6.6 9.1 11.8	4.1 5.1 9.0 12.2	1,012 4,127 2,459 2,163	594 1,223 1,161 953	418 2,905 1,298 1,209	
500 beds or more							
All ages	8.4	9.3	7.8	7,362	2,984	4,378	
Under 15 years	5.6 6.4 10.0 12.4	5.8 8.0 10.3 12.3	5.3 5.8 9.7 12.5	772 3,258 1,838 1,493	440 981 881 682	333 2,277 957 811	

Table 5. Average length of stay and number of patients discharged from short-stay hospitals with a single-listed diagnosis, by sex, bed size of hospital, and age: United States, 1977

Bed size of hospital and age	Both sexes	Male	Female	Both sexes	Male	Female	
All sizes	Avera	ge lengt in day	h of stay s		Number of patients discharged in thousands		
All ages	5.4	5.9	5.1	18,627	7,025	11,602	
Under 15 years	3.7 4.5 6.9 8.7	3.9 5.4 6.9 8.2	3.6 4.1 7.0 9.1	2,474 10,208 3,652 2,293	1,404 2,822 1,765 1,034	1,070 7,386 1,886 1,260	
6-99 beds						_	
All ages	4.5	4.8	4.3	3,555	1,396	2,159	
Under 15 years	3.0 3.8 5.5 7.3	3.0 4.2 5.5 7.2	3.0 3.6 5.4 7.3	465 1,906 689 496	256 591 326 223	209 1,314 363 272	
100-199 beds All ages	5.0	5.5	4.7	3,116	1,107	2,008	
Under 15 years	3.5 4.0 6.9 8.4	3.6 5.1 6.6 7.8	3.4 3.7 7.1 8.9	448 1,750 559 359	250 440 263 155	199 1,310 296 204	
200-299 beds All ages	5.2	5.7	4.9	2,853	1,074	1,779	
Under 15 years	3.6 4.3 6.8 8.6	3.9 5.3 6.7 8.1	3.3 3.9 6.8 9.0	409 1,562 533 349	-233 426 259 156	176 1,136 274 193	
<u>300-499 beds</u> All ages	5.6	6.1	5.3	5,109	1,930	3,179	
Under 15 years	3.7 4.7 7.2 9.2	3.7 5.7 7.2 8.6	3.7 4.3 7.2 9.7	663 2,808 1,039 599	388 765 505 271	275 2,043 534 327	
500 beds or more							
All ages	6.3	7.1	5.8	3,995	1,518	2,477	
Under 15 years	4.8 5.3 7.9 9.7	5.2 6.7 7.9 9.0	4.4 4.7 7.9 10.4	488 2,183 831 492	278 599 412 229	211 1,583 419 263	

Table 6. Average length of stay and number of patients discharged from short-stay hospitals with a single-listed surgery, by sex, bed size of hospital, and age: United States, 1977

Bed size of hospital and age	Both sexes	Male	Female	Both sexes	Male	Female
All sizes	Average length of stay in days			Num dischar	tients ousands	
All ages	6.6	7.2	6.2	10,227	4,093	6,133
Under 15 years	3.7 4.7 8.0 11.7	3.8 5.6 8.4 11.1	3.4 4.4 7.7 12.2	1,202 4,847 2,350 1,827	682 1,441 1,115 855	520 3,406 1,235 972
<u>6-99 beds</u>				Ì		
All ages	5.2	5.3	5.1	1,338	505	832
Under 15 years	2.6 3.9 6.5 10.1	2.6 4.1 6.8 9.1	2.5 3.8 6.2 10.9	188 678 263 208	108 193 113 92	80 485 151 116
100-199 beds		;				
All ages	5.7	6.3	5.3	1,680	630	1,050
Under 15 years	3.2 4.0 7.1 10.5	3.2 4.9 7.5 9.9	3.1 3.7 6.8 11.1	199 843 337 300	108 223 158 141	92 620 179 160
All ages	6.6	6.9	6.4	1,662	683	979
Under 15 years	3.5 4.6 8.0 11.9	3.7 5.3 7.9 10.8	3.3 4.3 8.1 12.8	202 775 376 310	111 243 182 147	91 532 194 163
<u>300-499 beds</u>						
All ages	7.0	7.6	6.6	3,079	1,255	1,824
Under 15 years	3.5 5.0 8.2 12.3	3.5 6.0 8.5 11.9	3.4 4.5 8.0 12.7	343 1,410 752 575	196 437 359 263	146 973 393 312
500 beds or more						
All ages	7.6	8.5	6.9	2,468	1,020	1,448
Under 15 years	5.1 5.6 9.0 12.3	5.4 6.8 9.5 12.2	4.6 5.1 8.5 12.4	270 1,142 622 434	159 345 303 212	111 797 318 222

Table 7. Average length of stay and number of patients discharged from short-stay hospitals with a first-listed diagnosis, by sex, type of ownership of hospital, and age: United States, 1977

Type of ownership and age	Both sexes	Male	Female	Both sexes	Male	Female
All types	Avera	ge lengtl in day	n of stay s		per of disc nts in thou	•
All ages	7.3	7.8	7.0	35,902	14,385	21,518
Under 15 years	4.2 5.3 8.5 11.1	4.3 6.3 8.6 10.8	4.1 4.9 8.4 11.3	3,775 15,180 8,604 8,344	2,137 4,553 4,042 3,653	1,638 10,627 4,562 4,690
Voluntary nonprofit						
All ages	7.5	8.0	7.2	25,236	10,022	15,214
Under 15 years	4.2 5.4 8.8 11.6	4.3 6.3 8.9 11.2	4.1 5.0 8.7 11.9	2,627 10,520 6,136 5,953	1,501 3,045 2,870 2,607	1,126 7,475 3,267 3,346
Government						
All ages	6.7	7.4	6.3	7,469	3,041	4,429
Under 15 years	4.4 5.2 8.0 9.8	4.6 6.6 8.0 9.7	4.3 4.6 8.0 9.9	886 3,351 1,613 1,619	492 1,041 776 732	394 2,310 838 887
Proprietary						
All ages	6.8	6.9	6.8	3,197	1,322	1,875
Under 15 years	3.4 5.2 7.4 10.1	3.3 5.7 7.5 9.8	3.5 5.0 7.3 10.2	263 1,308 854 771	145 467 396 314	118 842 458 458

Table 8. Average length of stay and number of patients discharged from short-stay hospitals with a single-listed diagnosis, by sex, type of ownership of hospital, and age: United States, 1977

Type of ownership and age	Both sexes	Male	Female	Both sexes	Male	Female
All types	Avera	ge length in day:	-		er of disc ts in thou	-
All ages	5.4	5.9	5.1	18,627	7,025	11,602
Under 15 years	3.7 4.5 6 .9 8.7	3.9 5.4 6.9 8.2	3.6 4.1 7.0 9.1	2,474 10,208 3,652 2,293	1,404 2,822 1,765 1,034	1,070 7,386 1,886 1,260
Voluntary nonprofit				,	:	
All ages	5.5	6.0	5.2	13,139	4,895	8,244
Under 15 years	3.8 4.5 7.1 8.9	3.9 5.4 7.1 8.3	3.7 4.1 7.1 9.3	1,723 7,123 2,635 1,658	980 1,891 1,273 751	743 5,232 1,362 907
Government						
All ages	5.1	5.8	4.7	3,971	1,500	2,470
Under 15 years	3.8 4.4 6.5 8.3	4.1 5.8 6.2 7.9	3.5 3.9 6.7 8.6	573 2,288 673 437	324 649 327 200	249 1,639 346 237
Proprietary						
All ages	5.2	5.5	5.1	1,517	629	888
Under 15 years	3.0 4.5 6.6 8.0	3.1 5.0 6.7 7.5	3.0 4.2 6.4 8.3	179 797 343 198	101 281 165 82	78 516 178 115

Table 9. Average length of stay and number of patients discharged from short-stay hospitals with a single-listed surgery, by sex, type of ownership of hospital, and age: United States, 1977

Type of ownership and age	Both sexes	Male	Female	Both sexes	Male	Female
All types	Avera	ige lengt in day	h of stay s		er of disc nts in tho	
All ages	6.6	7.2	6.2	10,227	4,093	6,133
Under 15 years	3.7 4.7 8.0 11.7	3.8 5.6 8.4 11.1	3.4 4.4 7.7 12.2	1,202 4,847 2,350 1,827	682 1,441 1,115 855	520 3,406 1,235 972
Voluntary nonprofit						
All ages	6.7	7.3	6.4	7,590	3,034	4,556
Under 15 years	3.6 4.7 8.1 11.8	3.8 5.5 8.5 11.2	3.4 4.4 7.8 12.4	851 3,523 1,796 1,419	477 1,038 855 664	374 2,485 941 755
Government						
All ages	6.5	7.4	5.9	1,843	728	1,114
Under 15 years	4.0 5.0 8.5 11.4	4.2 6.7 8.6 11.1	3.8 4.3 8.3 11.6	258 943 361 280	151 274 170 134	107 669 191 147
Proprietary						
All ages	5.6	5.9	5.4	794	330	463
Under 15 years	2.6 4.2 6.5 10.7	2.5 4.6 6.8 10.4	2.8 3.9 6.2 10.9	93 381 193 128	54 128 91 58	39 252 102 70

Table 10. Average length of stay for patients discharged from short-stay hospitals with a first-listed diagnosis, by color, marital status, age, geographic region, and bed size of hospital: United States, 1977

			Color				Mai	ried			Unma	arried ²	
Region and bed size of hospital	Total	White	All other	Not stated	Total ¹	15 years and over	15-44 years	45-64 years	65 years and over	15 years and over	15-44 years	45-64 years	65 years and over
All regions					A	verage lei	ngth of st	ay in day:	s				
All sizes	7.3	7.3	7.6	7.0	7.7	7.1	5.1	8.2	10.3	8.7	5.8	9.6	11.9
6-99 beds 100-499 beds 500 beds or more	6.0 7.4 8.4	6.0 7.4 8.4	6.0 7.5 8.5	5.9 6.8 8.7	6.3 7.8 8.8	5.7 7.1 8.3	4.3 5.0 6.1	6.2 8.3 9.6	8.2 10.6 11.8	7.3 8.8 9.5	4.6 5.7 7.0	7.3 9.7 11.1	9.9 12.4 12.9
Northeast All sizes	8.6	8.6	9.1	8.7	9.1	8.2	5.5	9.3	12.4	10.4	6.3	11.5	14.6
6-99 beds 100-499 beds 500 beds or more	7.6 8.5 9.8	7.4 8.5 9.7	11.3 8.4 9.9	8.2 8.2 10.5	7.9 8.9 10.2	7.0 8.1 9.6	5.1 5.4 6.3	7.6 9.2 11.1	10.4 12.1 14.8	9.4 10.4 10.9	6.3 6.0 7.4	9.3 11.2 14.3	12.9 14.6 15.8
North Central	:												
All sizes	7.5	7.6	8.1	7.1	7.9	7.4	5.4	8.6	10.4	8.8	6.2	10.0	11.8
6-99 beds 100-499 beds 500 beds or more	6.1 7.4 8.4	6.2 7.6 8.4	7.5 8.1 8.1	5.2 7.0 8.4	6.4 7.9 8.7	5,9 7.3 8.3	4.4 5.3 6.3	6.5 8.5 9.7	8.0 10.8 11.3	7.3 8.9 9.5	5.0 6.0 7.1	7.5 10.3 10.5	9.1 12.3 12.8
South													
All sizes	6.9	6.8	6.9	7.4	7.2	6.7	4.8	7.8	9.9	8.1	5.4	8.9	10.8
6-99 beds 100-499 beds 500 beds or more	6.0 6.8 8.1	6.0 6.7 8.2	5.8 7.0 8.0	6.7 7.8 7.3	6.3 7.2 8.4	5.8 6.7 8.0	4.3 4.7 6.0	6.2 8.1 9.1	8.4 10.3 11.3	7.2 8.2 9.1	4.4 5.2 6.7	6.7 9.1 11.1	9.7 11.2 11.9
<u>West</u>													
All sizes	6.0	6.1	6.7	5.1	6.3	5.8	4.3	6.6	8.1	7.0	5.0	7.4	9.9
6-99 beds 100-499 beds 500 beds or more	4.9 6.3 6.9	5.1 6.5 6.8	4.4 7.0 7.8	3.3 5.3 6.6	5.2 6.5 7.0	4.7 6.2 6.7	3.7 4.4 5.0	5.0 7.1 7.6	6.6 8.5 9.6	6.1 7.2 7.3	3.6 5.3 6.1	6.7 7.6 7.0	9.6 10.0 9.9

Table 11. Average length of stay and number of patients discharged with a first-listed diagnosis, by type of ownership of hospital, geographic region, age, sex, marital status, and sex and marital status: United States, 1977

[Discussives from four	- Cuciai Siloi	-stay Mospitals.	Excitates nev		1	
Region, age, sex, marital status, and sex and marital status	Voluntary nonprofit	Government	Proprietary	Voluntary nonprofit	Government	Proprietary
ALL REGIONS	Averaç	ge length of stay	in days	Numb	er of discharged in thousands	patients
<u>Age</u>						
All ages	7.5	6.7	6.8	25,236	7,469	3,197
All ages, 15 years and over	7.9	7.0	7.1	22,609	6,584	2,934
15-24 years	4.7 5.8 8.8 11.6	4.5 5.7 8.0 9.8	4.4 5.6 7.4 10.1	3,908 6,612 6,136 5,953	1,417 1,934 1,613 1,619	414 8,941 8,540 7,715
Sex, all ages						
Male Female	8.0 7.2	7.4 6.3	6.9 6.8	10,022 15,214	3,041 4,429	1,322 1,875
Marital status, 15 years and over 1						
MarriedUnmarried	7.3 9.0	6.4 8.0	6.7 7.8	13,917 7,694	3,704 2,397	1,748 1,077
Marital status and sex, 15 years and over ¹						
Married males	8.6 6.4 8.8 9.1	7.5 5.6 8.6 7.6	7.4 6.2 7.4 7.9	5,638 8,278 2,504 5,190	1,513 2,190 840 1,556	782 966 352 725
NORTHEAST REGION						
Age						
All ages	8.5	9.4		6,749	579	•••
All ages, 15 years and over	9.0	9.9		6,091	519	
15-24 years	4.9 6.0 9.7 13.4	6.3 7.2 12.1 16.0	 	962 1,714 1,710 1,705	106 189 126 99	
Sex, all ages				1		
MaleFemale	9.1 8.1	10.6 8.4		2,714 4,034	258 322	
Marital status, 15 years and over ¹						
MarriedUnmarried	8.1 10.3	8.4 11.0		3,593 2,230	212 238	
Marital status and sex, 15 years and over ¹						
Married males	9.7 6.9 9.9 10.4	10.6 6.8 12.0 10.4		1,487 2,108 752 1,478	89 123 98 139	

¹Does not include marital status "Not stated."

Table 11. Average length of stay and number of patients discharged with a first-listed diagnosis, by type of ownership of hospital, geographic region, age, sex, marital status, and sex and marital status: United States, 1977—Con.

Region, age, sex, marital status, and sex and marital status	Voluntary nonprofit	Government	Proprietary	Voluntary nonprofit	Government	Proprietary
NORTH CENTRAL REGION	Averag	ge length of stay	in days	Numb	er of discharged in thousands	patients
Age						
All ages	7.7	6.7		8,931	1,850	
All ages, 15 years and over	8.1	7.0		7,905	1,627	
15-24 years	5.0 6.2 9.1 11.7	4.7 5.7 7.7 9.0		1,477 2,312 2,174 1,942	329 423 406 470	
Sex, all ages						
Male	8.1 7.4	7.1 6.4		3,611 5,320	754 1,096	
Marital status, 15 years and over ¹						
MarriedUnmarried	7.6 9.0	6.5 7.8		4,826 2,703	961 590	
Marital status and sex, 15 years and over 1						
Married males	8.8 6.8 8.9 9.0	7.4 5.8 7.9 7.7	 	1,981 2,846 886 1,816	399 562 207 383	
SOUTH REGION						
Age		:				
All ages	7.1	6.5		6,118	3,675	
All ages, 15 years and over	7.4	6.9		5,471	3,217	
15-24 years	4.3 5.4 8.4 10.8	4.2 5.5 7.8 10.0	•••	966 1,683 1,393 1,428	685 941 798 794	
MaleFemale	7.6 6.8	7.1 6.1		2,292 3,826	1,472 2,202	•••
Marital status, 15 years and over 1					·	
Married Unmarried	6.8 8.4	6.3 7.8		3,553 1,738	1,885 1,153	
Marital status and sex, 15 years and over ¹						
Married males	8.2 6.0 8.0 8.6	7.4 5.6 8.4 7.5	 	1,342 2,210 529 1,209	784 1,101 372 781	

¹Does not include marital status "Not stated."

Table 11. Average length of stay and number of patients discharged with a first-listed diagnosis, by type of ownership of hospital, geographic region, age, sex, marital status, and sex and marital status: United States, 1977—Con.

Region, age, sex, marital status, and sex and marital status	Voluntary nonprofit	Government	Proprietary	Voluntary Government P		Proprietary
WEST REGION	Averaç	ge length of stay	in days	Numb	er of discharged in thousands	patients
Age						
All ages	6.1	6.1		3,437	1,365	
All ages, 15 years and over	6.4	6.2		3,141	1,220	
15-24 years	3.7 4.8 6.8 9.1	4.5 5.3 7.3 8.4		503 903 859 877	298 381 284 257	
Sex, all ages						
Male Female	6.5 5.9	7.0 5.4		1,405 2,033	556 809	
Marital status, 15 years and over 1						
Married	5.9 7.2	5.7 7.0		1,943 1,024	646 415	
Marital status and sex, 15 years and over ¹						
Married males	6.8 5.2 7.3 7.2	6.8 5.0 8.0 6.3		827 1,115 337 686	241 405 162 253	

¹Does not include marital status "Not stated."

Table 12. Average length of stay for patients discharged from short-stay hospitals with a single-listed diagnosis, by color, marital status, age, geographic region, and bed size of hospital: United States, 1977

			Color				Mai	rried			Unma	rried ²	, , , , , , , , , , , , , , , , , , ,
Region and bed size of hospital	Total	White	All other	Not stated	Total ¹	15 years and over	15-44 years	45-64 years	65 years and over	15 years and over	15-44 years	45-64 years	65 years and over
All regions					A	verage ler	igth of st	ay in day:	5				
All sizes	5.4	5.4	5.4	5.1	5.6	5.3	4.2	6.7	7.8	6.3	4.9	7.8	9.7
6-99 beds	4.5 5.3 6.3	4.5 5.4 6.4	4.7 5.2 6.0	4.4 5.0 6.4	4.7 5.6 6.5	4.4 5.3 6.2	3.7 4.2 5.0	5.3 6.8 7.7	6.7 7.8 8.9	5.2 6.3 7.0	4.0 4.8 5.7	6.2 7.9 8.7	8.0 9.9 10.6
All sizes	6.1	6.1	6.4	6.1	6.4	6.0	4.6	7.6	9.5	7.2	5.4	9.1	12.1
6-99 beds 100-499 beds 500 beds or more	5.9 6.0 6.7	5.6 6.1 6.6	10.5 5.9 6.7	6.0 5.7 7.1	6.2 6.3 6.9	5.2 5.9 6.7	4.3 4.5 5.1	6.3 7.6 8.1	7.7 9.3 11.1	7.7 7.2 7.2	5.9 5.1 5.9	7.7 9.2 9.9	13.8 12.1 11.0
North Central													
All sizes	5.6	5.7	5.8	5.2	5.9	5.6	4.6	7.1	7.8	6.5	5.2	8.4	9.6
6-99 beds 100-499 beds 500 beds or more	4.5 5.4 6.5	4.6 5.5 6.6	6.2 5.8 5.9	4.1 5.1 6.4	4.7 5.8 6.7	4.5 5.4 6.4	3.7 4.5 5.1	5.3 6.9 8.1	6.4 7.6 8.8	5.2 6.4 7.4	3.9 5.1 6.0	6.2 8.8 8.6	7.2 9.4 11.5
South													
All sizes	5.1	5.1	4.9	5.6	5.3	5.0	4.1	6.4	7.8	5.9	4.6	7.4	8.8
6-99 beds 100-499 beds 500 beds or more	4.5 5.0 6.2	4.5 4.9 6.4	4.3 4.8 5.7	5.1 5.7 5.7	4.7 5.2 6.3	4.6 4.9 6.0	3.7 3.9 5.0	5.3 6.6 7.4	7.4 7.8 8.4	5.0 6.0 6.9	3.8 4.5 5.7	5.7 7.3 9.0	7.7 9.3 9.8
West All sizes	4.4	4.6	4.4	3.9	4.6	4.4	3.5	5.5	6.0	5.0	4.0	5.8	7.8
6-99 beds 100-499 beds 500 beds or more	3.7 4.6 5.3	3.8 4.8 5.3	3.4 4.5 4.8	2.7 4.0 5.4	3.8 4.8 5.4	3.7 4.6 5.1	3.1 3.7 4.0	4.5 5.8 6.4	5.2 6.2 6.8	3.9 5.2 5.7	3.2 4.2 4.4	5.5 5.9 5.9	5.9 8.0 9.5

¹Includes patients for whom marital status was not stated. ²Includes single, separated, divorced, and widowed persons.

Table 13. Average length of stay for patients discharged from short-stay hospitals with a single-listed surgery, by color, marital status, age, geographic region, and bed size of hospital: United States, 1977

			Color				Mar	ried			Unma	erried ²	
Region and bed size of hospital	Total	White	All other	Not stated	Total ¹	15 years and over	15-44 years	45-64 years	65 years and over	15 years and over	15-44 years	45-64 years	65 years and over
All regions					A	/erage lei	ngth of st	ay in day:	S				
All sizes	6.6	6.7	6.9	6.1	7.0	6.6	4.7	7.7	10.6	7.7	4.8	9.3	13.0
6-99 beds	5.2 6.5 7.6	5.2 6.7 7.6	5.6 6.5 8.0	4.7 6.0 6.9	5.6 7.0 7.9	5.2 6.5 7.6	4.0 4.6 5.5	6.0 7.6 8.6	8.4 10.5 11.8	6.3 7.7 8.3	3.7 4.6 5.7	7.8 9.1 10.1	12.2 13.2 12.9
Northeast All sizes	7.6	7.7	8.1	7.0	8.1	7.6	5.0	8.5	12.8	8.8	5.1	10.8	14.9
6-99 beds 100-499 beds 500 beds or more	5.9 7.6 8.3	5.9 7.6 8.4	7.9 7.7 8.7	6.0 7.0 7.3	6.3 8.0 8.6	5.3 7.5 8.5	3.9 5.0 5.6	5.9 8.6 9.0	10.7 12.3 15.6	8.1 8.9 8.7	4.1 4.9 5.8	11.7 10.2 12.5	13.5 15.3 13.6
North Central													
All sizes	6.7	6.7	6.8	6.4	7.1	6.8	5.0	7.9	10.4	7.5	4.7	9.2	12.6
6-99 beds 100-499 beds 500 beds or more	5.5 6.4 7.5	5.5 6.5 7.5	6.4 5.9 7.4	5.2 6.3 7.0	6.0 6.9 7.8	5.5 6.5 7.7	4.1 4.7 5.7	6.4 7.6 8.8	8.9 10.2 11.2	6.6 7.4 8.0	3.7 4.4 5.6	7.3 9.5 9.3	12.1 12.6 12.7
South													
All sizes	6.5	6.3	7.0	6.5	6.8	6.3	. 4.6	7.5	10.7	7.9	4.9	9.6	13.0
6-99 beds 100-499 beds 500 beds or more	5.5 6.2 7.6	5.5 6.0 7.6	5.6 6.7 8.3	5.0 7.1 6.1	6.0 6.6 7.9	5.6 6.0 7.5	4.1 4.4 5.5	6.8 7.2 8.6	9.4 10.9 11.1	6.7 7.8 8.7	4.0 4.7 5.8	7.0 9.3 11.0	13.0 12.9 13.2
West		5.0		4.0			4.0		-			- 4	40.5
All sizes	5.5	5.8	5.5	4.6	5.8	5.6	4.2	6.3	8.0	6.2	4.3	7.1	10.5
6-99 beds 100-499 beds 500 beds or more	4.3 5.8 6.5	4.4 6.2 6.4	4.9 5.3 7.3	2.6 4.7 7.6	4.6 6.0 6.7	4.5 5.8 6.1	3.8 4.2 4.6	4.9 6.6 6.8	6.2 8.4 8.7	4.7 6.5 7.2	3.1 4.4 6.0	7.4 7.1 6.4	9.7 10.6 10.6

¹Includes patients for whom marital status was not stated.
²Includes single, separated, divorced, and widowed persons.

Table 14. Average length of stay and number of patients discharged from short-stay hospitals with a first-listed diagnosis, by sex, geographic region, and age: United States, 1977

Region and age	Both sexes	Male	Female	Both sexes	Male	Female
All regions	Avera	ige lengtl in day			nber of pa rged in the	
All ages	7.3	7.8	7.0	35,902	14,385	21,518
Under 15 years	4.2 5.3 8.5 11.1	4.3 6.3 8.6 10.8	4.1 4.9 8.4 11.3	3,775 15,180 8,604 8,344	2,137 4,553 4,042 3,653	1,638 10,627 4,562 4,690
Northeast						
All ages	8.6	9.2	8.2	7,811	3,191	4,620
Under 15 years	4.5 5.8 9.9 13.6	4.6 7.2 10.1 13.2	4.4 5.2 9.7 14.0	739 3,182 1,982 1,908	432 973 954 832	307 2,209 1,028 1,076
North Central						
All ages	7.5	7.9	7.2	10,835	4,392	6,443
Under 15 years	4.3 5.7 8.9 11.2	4.4 6.7 8.9 10.8	4.1 5.3 8.8 11.5	1,251 4,567 2,594 2,424	714 1,395 1,220 1,063	537 3,172 1,374 1,360
South						
All ages	6.9	7.3	6.6	11,290	4,366	6,924
Under 15 years	4.1 5.0 8.0 10.4	4.2 5.8 8.2 10.2	4.0 4.7 7.9 10.5	1,214 4,843 2,573 2,660	667 1,389 1,182 1,128	547 3,454 1,391 1,532
West						
All ages	6.0	6.5	5.7	5,966	2,435	3,531
Under 15 years 15-44 years 45-64 years 65 years and over	3.7 4.6 6.8 8.9	3.8 5.5 6.7 8.8	3.7 4.2 6.9 9.0	571 2,588 1,455 1,352	324 796 686 629	247 1,792 769 723

Table 15. Average length of stay and number of patients discharged from short-stay hospitals with a single-listed diagnosis, by sex, geographic region, and age: United States, 1977

Region and age	Both sexes	Male	Female	Both sexes	Male	Female
All regions	Average length of stay in days					tients ousands
All ages	5.4	5.9	5.1	18,627	7,025	11,602
Under 15 years	3.7 4.5 6.9 8.7	3.9 5.4 6.9 8.2	3.6 4.1 7.0 9.1	2,474 10,208 3,652 2,293	1,404 2,822 1,765 1,034	1,070 7,386 1,886 1,260
Northeast						
All ages	6.1	6.8	5.7	4,004	1,531	2,473
Under 15 years	4.0 4.9 7.9 10.9	4.0 6.2 7.9 10.4	3.9 4.4 7.9 11.2	506 2,186 837 476	297 611 411 211	209 1,574 425 264
North Central						
All ages	5.6	6.1	5.3	5 ,6 57	2,170	3,486
Under 15 years	3.8 4.8 7.3 8.7	4.0 5.9 7.2 8.0	3.6 4.4 7.5 9.3	819 3,031 1,110 696	459 848 548 316	360 2,183 563 380
South		i				
All ages	5.1	5.6	4.8	5,802	2,119	3,683
Under 15 years	3.7 4.3 6.6 8.3	3.9 5.0 6.6 8.0	3.6 4.0 6.6 8.6	783 3,245 1,069 705	439 870 498 312	344 2,376 571 393
West						
All ages	4.4	4.9	4.2	3,165	1,204	1,960
Under 15 years	3.2 3.7 5.6 6.7	3.4 4.5 5.7 6.2	3.0 3.4 5.5 7.2	366 1,746 636 418	209 493 308 195	157 1,253 328 223

Table 16. Average length of stay and number of patients discharged from short-stay hospitals with a single-listed surgery, by sex, geographic region, and age: United States, 1977

Region and age	Both sexes	Male	Female	Both sexes	Male	Female
All regions	Avera	ge lengti in day	n of stay s		ber of patients ged in thousands	
All ages	6.6	7.2	6.2	10,227	4,093	6,133
Under 15 years	3.7 4.7 8.0 11.7	3.8 5.6 8.4 11.1	3.4 4.4 7.7 12.2	1,202 4,847 2,350 1,827	682 1,441 1,115 855	520 3,406 1,235 972
Northeast						
All ages	7.6	8.4	7.1	2,329	938	1,392
Under 15 years	3.8 5.1 9.1 13.9	3.8 6.2 9.4 13.2	3.8 4.6 8.8 14.5	238 1,071 564 456	139 311 271 216	99 760 294 240
North Central						
All ages	6.7	7.2	6.3	3,128	1,268	1,860
Under 15 years	3.7 4.9 8.2 11.5	4.1 5.6 8.6 10.8	3.3 4.5 7.8 12.0	399 1,441 736 552	220 448 351 250	179 994 386 302
South						
All ages	6.5	7.2	6.0	2,853	1,088	1,765
Under 15 years	3.8 4.7 8.0 11.8	3.9 5.5 8.6 11.6	3.6 4.4 7.5 12.0	355 1,429 595 473	205 384 281 219	151 1,045 314 255
West						
All ages	5.5	5.9	5.3	1,916	800	1,116
Under 15 years	3.2 4.3 6.5 9.0	3.2 5.2 6.4 8.5	3.2 3.8 6.7 9.5	210 905 454 347	118 297 213 171	92 607 241 176

Table 17. Average length of stay for patients discharged from short-stay hospitals with a first-listed diagnosis, by geographic region, bed size of hospital, sex, and age: United States, 1977

Sex and age		Northeast			North Central			South			West		
	All regions	6-99 beds	100- 499 beds	500 beds or more	6-99 beds	100- 499 beds	500 beds or more	6-99 beds	100- 499 beds	500 beds or more	6-99 beds	100- 499 beds	500 beds or more
Both sexes	Average length of stay in days												
All ages	7.3	7.6	8.5	9.8	6.1	7.4	8.4	6.0	6.8	8.1	4.9	6.3	6.9
Under 15 years 15-44 years 45-64 years 65 years and over	4.2 5.3 8.5 11.1	3.2 5.6 8.1 11.8	4.3 5.6 9.7 13.5	6.0 6.8 12.0 15.5	3.4 4.6 6.7 8.5	4.0 5.5 8.9 11.6	5.4 6.6 9.8 12.1	3.3 4.3 6.4 9.1	4.0 4.8 8.3 10.8	5.6 6.3 9.6 11.7	2.8 3.7 5.4 7.8	3.8 4.8 7.2 9.1	5.6 5.4 7.7 9.8
Male													
All ages	_/ 7.8	7.7	9.0	11.3	6.3	7.8	9.0	6.0	7.4	9.1	5.0	6.8	7.5
Under 15 years	4.3 6.3 8.6 10.8	2.9 6.3 7.9 11.8	4.2 6.8 9.9 12.7	6.5 9.0 12.5 15.8	3.4 5.2 6.9 8.0	4.2 6.5 8.7 11.3	5.5 8.0 10.2 11.2	3.2 ·4.4 6.3 8.6	4.1 5.7 8.5 10.6	5.9 7.7 9.9 12.0	3.0 3.9 4.9 7.7	3.9 5.9 7.2 9.0	5.2 6.8 7.5 9.8
Female including deliveries													
All ages	7.0	7.5	8.2	8.8	6.0	7.2	8.0	6.1	6.5	7.5	4.9	5.9	6.4
Under 15 years	4.1 4.9 8.4 11.3	3.5 5.1 8.3 11.8	4.3 5.1 9.4 14.1	5.2 5.9 11.6 15.2	3.4 4.3 6.5 9.0	3.8 5.1 9.0 11.7	5.2 6.0 9.5 12.7	3.4 4.3 6.4 9.5	3.9 4.5 8.2 10.9	5.2 5.7 9.4 11.4	2.5 3.6 5.8 7.9	3.7 4.3 7.3 9.3	6.1 4.8 7.8 9.7
Female excluding deliveries													
All ages	7.6	8.0	8.8	9.5	6.3	7.7	8.7	6.5	7.1	8.3	5.3	6.5	7.2
Under 15 years	4.1 5.4 8.4 11.3	3.5 5.5 8.3 11.8	4.3 5.4 9.4 14.1	5.2 6.2 11.6 15.2	3.4 4.5 6.5 9.0	3.8 5.5 .9.1 11.7	5.2 6.8 9.5 12.7	3.4 4.8 6.4 9.5	3.9 5.0 8.2 10.9	5.2 6.6 9.4 11.4	2.5 4.0 5.8 7.9	3.7 4.9 7.3 9.3	6.1 5.8 7.8 9.7

Table 18. Average length of stay for patients discharged from short-stay hospitals with a single-listed diagnosis, by geographic region, bed size of hospital, sex, and age: United States, 1977

[Discharges from non-Federal short-stay hospitals. Excludes newborn infants]

			Northea	st	No	orth Cen	itral		South			West	
Sex and age	All regions	6-99 beds	100- 499 beds	500 beds or more	6-99 beds	100- 499 beds	500 beds or more	6-99 beds	100- 499 beds	500 beds or more	6-99 beds	100- 499 beds	500 beds or more
Both sexes		Average length of stay in days											
All ages	5.4	5.9	6.0	6.7	4.5	5.4	6.5	4.5	5.0	6.2	3.7	4.6	5.3
Under 15 years	3.7 4.5 6.9 8.7	2.8 5.0 6.8 11.2	3.9 4.8 7.9 10.8	5.1 5.5 8.6 11.1	3.1 3.8 5.4 6.8	3.6 4.7 7.2 8.6	4.6 5.4 8.1 9.9	3.1 3.8 5.4 7.5	3.6 4.1 6.7 8.5	5.1 5.2 7.7 9.2	2.6 3.2 4.7 5.4	3.4 3.9 5.8 7.0	4.2 4.2 6.4 8.0
Male													
All ages	5.9	6.2	6.7	7.9	4.9	5.8	7.2	4.7	5.6	6.9	3.9	5.1	5.7
Under 15 years	3.9 5.4 6.9 8.2	2.6 5.7 6.4 12.8	3.8 6.0 8.1 10.1	5.8 7.5 8.5 10.7	3.0 4.2 5.8 7.0	3.8 5.6 6.8 8.0	4.9 7.0 8.2 8.6	3.0 4.1 5.4 7.2	3.8 5.0 6.7 8.2	5.4 6.3 7.8 9.1	2.9 3.5 4.6 5.2	3.5 4.8 5.9 6.3	4.1 5.2 6.2 7.4
Female including deliveries													
All ages	5.1	5.6	5.6	6.0	4.3	5.2	6.0	4.4	4.7	5.8	3.5	4.3	5.0
Under 15 years	3.6 4.1 7.0 9.1	3.2 4.5 7.4 10.0	3.9 4.3 7.8 11.3	4.1 4.7 8.6 11.5	3.2 3.6 5.2 6.7	3.4 4.3 7.6 9.1	4.3 4.8 8.1 11.1	3.2 3.6 5.4 7.8	3.4 3.8 6.8 8.8	4.7 4.8 7.7 9.2	2.3 3.1 4.7 5.7	3.2 3.5 5.6 7.5	4.5 3.8 6.7 8.7
Female excluding deliveries													
All ages	5.5	6.0	6.1	6.4	4.4	5.6	6.7	4.8	5.1	6.6	3.8	4.8	5,8
Under 15 years	3.6 4.4 7.0 9.1	3.2 4.8 7.4 10.0	3.9 4.4 7.8 11.3	4.1 4.9 8.7 11.5	3.2 3.6 5.2 6.7	3.4 4.5 7.6 9.1	4.3 5.4 8.1 11.1	3.2 4.1 5.4 7.8	3.4 4.1 6.8 8.8	4.7 5.7 7.7 9.2	2.3 3.3 4.7 5.7	3.2 3.9 5.6 7.5	4.5 4.5 6.7 8.7

Table 19. Average length of stay for patients discharged from short-stay hospitals with a single-listed surgery, by geographic region, bed size of hospital, sex, and age: United States, 1977

[Discharges from non-Federal short-stay hospitals. Excludes newborn infants]

			Northea	st	No	rth Cen	tral		South			West	
Sex and age	AII regions	6-99 beds	100- 499 beds	500 beds or more	6-99 beds	100- 499 beds	500 beds or more	6-99 beds	100- 499 beds	500 beds or more	6-99 beds	100- 499 beds	500 beds or more
Both sexes	·	Average length of stay in days											
All ages	. 6.6	5.9	7.6	8.3	5.5	6.4	7.5	5.5	6.2	7.6	4.3	5.8	6.5
Under 15 years	3.7 4.7 8.0 11.7	2.4 4.0 7.5 12.1	3.4 5.0 9.0 13.8	5.3 5.7 9.9 14.7	2.6 4.0 6.5 10.6	3.4 4.6 7.9 11.4	4.7 5.7 8.9 11.8	2.9 4.1 7.0 11.2	3.4 4.5 7.7 11.9	5.5 5.6 9.2 12.1	2.1 3.6 5.4 7.2	3.4 4.3 6.8 9.3	5.2 5.3 6.9 9.6
Male													
All ages	7.2	6.0	8.2	9.6	5.6	6.8	8.2	5.7	7.0	8.7	4.3	6.2	7.4
Under 15 years	3.8 5.6 8.4 11.1	2.5 ,4.1 8.5 10.0	3.3 6.0 9.3 12.9	5.7 7.4 10.1 15.2	2.8 3.9 6.8 10.1	3.8 5.4 8.2 10.7	5.0 6.6 9.6 11.2	2.7 4.1 7.8 10.2	3.6 5.4 8.2 11.5	5.8 6.8 9.8 12.5	2.4 4.0 5.0 6.4	3.2 5.3 6.6 8.8	5.0 7.0 7.4 9.3
Female including deliveries													
All ages	6.2	5.9	7.1	7.5	5.4	6.1	6.9	5.4	5.7	6.9	4.2	5.5	5.9
Under 15 years45-64 years	3.4 4.4 7.7 12.2	2.3 3.9 6.8 13.7	3.6 4.5 8.7 14.6	4.8 5.1 9.8 14.3	2.5 4.0 6.4 11.0	3.0 4.3 7.7 11.9	4.2 5.2 8.2 12.5	3.0 4.0 6.3 11.9	3.3 4.1 7.3 12.2	4.9 5.1 8.6 11.8	1.9 3.4 5.8 8.0	3.5 3.8 6.9 9.8	5.3 4.5 6.5 9.9
Female excluding deliveries													
All ages,	6.4	6.1	7.3	7.7	5.6	6.2	7.2	5.6	6.0	7.3	4.3	5.7	6.3
Under 15 years	3.4 4.2 7.7 12.2	2.3 3.5 6.8 13.7	3.5 4.1 8.7 14.6	4.8 4.7 9.8 14.3	2.5 3.9 6.4 11.0	3.0 4.0 7.7 11.9	4.1 5.2 8.2 12.5	3.0 4.0 6.3 11.9	3.3 4.1 7.3 12.2	4.9 5.1 8.6 11.8	1.8 3.3 5.8 8.0	3.5 3.8 6.9 9. 8	5.3 4.8 6.5 9.9

APPENDIXES

CONTENTS

I.	Technical Notes on Methods	35 35 37 39
II.	Definitions of Certain Terms Used in This Report Hospital and Hospital Characteristics Terms Relating to Hospitalization Terms Relating to Diagnoses Terms Relating to Surgery Demographic Terms	42 42 42 42 43 43
	LIST OF APPENDIX FIGURES	
I.	Medical Abstract for the Hospital Discharge Survey	38
II.	Approximate relative standard errors of estimated numbers of patients discharged, or of first-listed diagnoses, by selected patient and hospital characteristics	40
	LIST OF APPENDIX TABLES	
I.	Distribution of short-stay hospitals in the universe (Master Facility Inventory) and in the National Hospital Discharge Survey sample and the number of hospitals that participated in the survey, by geographic region and bed size of hospital: United States, 1977	36
II.	Civilian noninstitutionalized population by sex, age, and geographic region: United States, July 1, 1977	39
III.	Approximate standard errors of percents shown in this report for patient discharges: patient characteristics classified by geographic region and for all hospitals	41
IV.	Approximate standard errors of average lengths of stay shown in this report for patient discharges: patient characteristics cross-classified by geographic region, bed size of hospital, geographic region by bed size of hospital and for all hospitals	41

APPENDIX I

TECHNICAL NOTES ON METHODS

Statistical Design of the National Hospital Discharge Survey

Scope of the survey.—The scope of the National Hospital Discharge Survey (NHDS) encompasses patients discharged from noninstitutional hospitals, exclusive of military and Veterans Administration hospitals, located in the 50 States and the District of Columbia. Only hospitals having six beds or more for patient use and those in which the average length of stay for all patients is less than 30 days are included in the survey. Although all discharges of patients from these hospitals are within the scope of the survey, discharges of newborn infants from all hospitals are excluded from this report as well as discharges of all patients from Federal hospitals.

Sampling frame and size of sample.—The sampling frame (universe) for hospitals in the NHDS is the Master Facility Inventory of Hospitals and Institutions (MFI). A detailed description of how the MFI was developed, its contents, plans for maintaining it, and procedures for assessing the completeness of its coverage has been published.²

The universe for the survey consisted of 6,965 short-stay hospitals contained in the MFI in 1963, 442 hospitals that were added to the MFI in 1969, another 223 hospitals that were added in 1972, and 273 hospitals that were added in 1976. The distribution of the hospitals in the MFI and in the NHDS sample is shown by bed size and geographic region in table I.

The sample for 1977 consisted of 535 hospitals. Of these hospitals, 68 refused to participate

NOTE: A list of references follows the text.

and 44 were out of scope either because the hospital had gone out of business or because it failed to meet the definition of a short-stay hospital. Thus 423 hospitals participated in the survey during 1977 and provided approximately 224,000 abstracts of medical records.

Sample design.—All hospitals with 1,000 beds or more in the universe of short-stay hospitals were selected with certainty in the sample. All hospitals with fewer than 1,000 beds were stratified, the primary strata were the 24 size-by-region classes shown in table I. Within each of these 24 primary strata, the allocation of the hospitals was made through a controlled selection technique so that hospitals in the sample would be properly distributed with regard to type of ownership and geographic region. Sample hospitals were drawn with probabilities ranging from certainty for the largest hospitals to 1 in 40 for the smallest hospitals.

The within-hospital sampling ratio for selecting sample discharges varied inversely with the probability of selection of the hospital. The smallest sampling fraction of discharged patients was taken in the largest hospitals, and the largest fraction was taken in the smallest hospitals. This sampling was done to compensate for the fact that hospitals were selected with probabilities proportionate to their size class and to ensure that the overall probability of selecting a discharge would be approximately the same in each size class.

In nearly all hospitals, the daily listing sheet of discharges was the frame from which the subsamples of discharges were selected within the sample hospitals. The sample discharges were selected by a random technique, usually on the basis of the terminal digit(s) of the patient's

Table I. Distribution of short-stay hospitals in the universe (Master Facility Inventory) and in the National Hospital Discharge Survey sample and the number of hospitals that participated in the survey, by geographic region and bed size of hospital: United States, 1977

Bed size of hospital	AII regions	North- east	North Central	South	West	
All sizes	Number of hospitals					
Universe	7,903	1,177	2,214	3,130	1,472	
	535	132	152	170	81	
	423	113	120	125	65	
Universe	3,486	221	889	1,658	718	
	69	8	18	30	13	
	44	6	15	16	7	
Universe	1,871	299	482	721	369	
	79	14	20	31	14	
	58	10	14	23	11	
100-199 beds				!	:	
Universe Total sample Number participating	1,371	298	404	448	221	
	117	26	33	39	19	
	98	25	28	27	18	
200-299 beds						
Universe Total sample Number participating	614	193	164	160	97	
	97	31	27	24	15	
	76	26	21	17	12	
300-499 beds						
Universe Total sample Number participating	408	112	134	109	53	
	97	25	32	28	12	
	81	21	26	25	9	
500-999 beds						
Universe Total sample Number participating	135	45	48	29	13	
	58	19	19	13	7	
	49	17	13	12	7	
1,000 beds or more						
Universe Total sample Number participating	18	9	3	5	1	
	18	9	3	5	1	
	17	8	3	5	1	

medical record number—a number assigned when the patient was admitted to the hospital. If the hospital's daily discharge listing did not show the medical record numbers, the sample was selected by starting with a randomly selected discharge and taking every kth discharge thereafter.

Data Collection and Processing

Data collection.—Depending on the study procedure agreed on with the hospital administrator, the sample selection and the transcription of information from the hospital records to abstract forms were performed either by the hospital staff or by representatives of the National Center for Health Statistics (NCHS) or by both. In about two-thirds of the hospitals that participated in the NHDS during the year, this work was performed by the medical records department of the hospital. In the remaining hospitals, the work was performed by personnel of the U.S. Bureau of the Census acting for NCHS.

Survey hospitals used an abstract form to transcribe data from the hospital records. The abstract form provides for recording demographic data, admission and discharge dates, zip code of the patient's residence, expected sources of payment, disposition of patient at discharge, and information on discharge diagnoses and surgical operations or procedures (figure I). All discharge diagnoses and operations were listed on the abstract form in the order of the principal one, or the first-listed one if the principal one was not identified, followed by the order in which all other diagnoses or operations were entered on the face sheet of the medical record.

Shipments of completed abstract forms for each sample hospital were transmitted, along with sample selection control sheets, to a Census Regional Office. Every shipment of abstracts was reviewed and each abstract form was checked for completeness. Abstracts were then sent to NCHS for processing.

Medical coding and edit.—The medical information recorded on the sample patient abstracts was coded centrally by NCHS staff. A maximum of five diagnostic codes was assigned for each sample abstract; in addition, if the medical information included surgery, a maximum of three codes for surgical operations and procedures was

assigned. Following the conversion of the data on the Medical Abstract to computer tape, a final medical edit was accomplished by computer inspection runs and a review of rejected abstracts. If sex or age of patient was incompatible with the recorded medical information, priority was given to the medical information in the editing decision.

The basic system used for coding the diagnoses on the NHDS sample patient abstracts is the Eighth Revision International Classification of Diseases, Adapted for Use in the United States³ (ICDA).

Patient characteristics not stated.—If the age or sex of the patient was not stated on the hospital records of sample hospitals (the face sheet of patient's medical record), it was imputed by assigning the patient an age or sex consistent with the age or sex of other patients with the same diagnostic code. Color was identified as "not stated." If the dates of admission or discharge were not given and if they could not be obtained from the monthly sample listing sheet transmitted by the sample hospital, a length of stay was imputed by assigning the patient a stay characteristic of the stays of other patients of the same age. The age and sex of the patient were not stated for less than one-fourth of 1 percent of the discharges. However, color was not stated for 12.8 percent of all discharges, and, therefore, caution should be used in drawing conclusions from the data by color which are shown.

Rounded numbers.—Estimates of the numbers of inpatient discharges, discharges with surgery, and all-listed operations have been rounded to the nearest thousand for tabular presentation. For this reason, detailed figures within the tables do not always add to totals. Rates and percents were calculated on the basis of unrounded figures and will not necessarily agree with computations made from the rounded data.

Population estimates.—The population estimates used in computing rates are unpublished estimates for the U.S. civilian noninstitutionalized population on July 1 of the data year provided by the U.S. Bureau of the Census. The estimates by age and sex and by geographic region

NOTE: A list of references follows the text.

CONFIDENTIAL — All information which confidential, will be used only by persons e other persons or used for any other purpose.	engaged in and for the pu	ation of an individual or of an est rposes of the survey, and will not be	tablishment will be held disclosed or released to			
FORM HD5-1 (12-13-76) DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE PUBLIC HEALTH SERVICE HEALTH RESOURCES ADMINISTRATION NATIONAL CENTER FOR HEALTH STATISTICS						
MEDICAL A	ABSTRACT - HOSPIT	TAL DISCHARGE SURVEY				
A. PATIENT IDENTIFICATION 1. Hospital number 2. HDS number		4. Date of admission				
3. Medical Record number	6.	Residence ZIP Code				
B, PATIENT CHARACTERISTICS			<i>(.</i> –			
7. Date of birth	Year 8.	• Age (Complete only if Date of Birth not given)	Units 1 Years 2 Months 3 Days			
9. Sex (Mark one)	Male 2 Femal	le 3 Not stated				
10. Race or Color (Mark one) 1 White 2 Black 3 Other 4 Not stated						
11. Marital Status (Mark one)	Married 2 Single 3	☐ Widowed 4 ☐ Divorced 5 ☐ S	eparated 6 Not stated			
8 No charge	payments ommercial insurance	13. Disposition of Patient (Mo	charged home divice to another			
		□ Se	ee reverse side			
D. SURGICAL AND DIAGNOSTIC PROC Principal:Other/additional:			Date: Month Day Year			
Completed by		Date				

TUS, GOVERNMENT PRINTING OFFICE: 1976-758-250

Figure I. Medical Abstract for the Hospital Discharge Survey

are presented in table II and are consistent with the population estimates published in *Current Population Reports*, Series P-25. However, they are not official population estimates of the Bureau of the Census.

Table II. Civilian noninstitutionalized population by sex, age, and geographic region: United States, July 1, 1977

[Population estimates consistent with Series P-25, Current Population Reports, U.S. Bureau of the Census]

Age and region	Both sexes	Male	Female		
	Population in thousands				
All ages	212,182	102,399	109,782		
Northeast	48,529	23,292	25,237		
North Central	57,055	27,796	29,259		
South	68,278	32,694	35,584		
West	38,319	18,617	19,702		
0-14 years	51,481	26,260	25,220		
Under 1 year	3,161	1,619	1,542		
1-4 years	12,067	6,166	5,901		
5-14 years	36,253	18,475	17,778		
Northeast	11,081	5,662	5,419		
North Central	13,950	7,123	6,827		
South	17,063	8,687	8,375		
West	9,386	4,787	4,599		
15-44 years	95,060	46,231	48,828		
15-24 years	39,720	19,496	20,224		
25-34 years	32,247	15,642	16,604		
35-44 years	23,093	11,092	12,001		
Northeast	21,267	10,360	10,907		
North Central	25,722	12,684	13,038		
South	30,455	14,622	15,833		
West	17,616	8,566	9,051		
45-64 years	43,357	20,704	22,653		
45-54 years	23,174	11,174	11,999		
55-64 years	20,183	9,529	10,654		
Northeast	10,761	5,085	5,676		
North Central	11,387	5,490	5,896		
South	13,536	6,394	7,142		
West	7,674	3,735	3,938		
65 years and over	22,284	9,204	13,080		
65-74 years	14,278	6,205	8,072		
75 years and over	8,007	2,999	5,008		
Northeast	5,421	2,186	3,235		
North Central	5,996	2,499	3,497		
South	7,225	2,991	4,234		
West	3,643	1,529	2,114		

Reliability of Estimates

Estimation.—Statistics produced by NHDS are derived by a complex estimating procedure. The basic unit of estimation is the sample inpatient discharge abstract. The estimating procedure used to produce essentially unbiased national estimates in NHDS has three principal components: inflation by reciprocals of the probabilities of sample selection, adjustment for nonresponse, and ratio adjustment to fixed totals. These components of estimation are described in appendix I of two earlier publications.^{4,5}

Measurement errors.—As in any survey, results are subject to nonsampling or measurement errors, which include errors due to hospital non-response, missing abstracts, information incompletely or inaccurately recorded on abstract forms, and processing errors. Some of these errors were discussed under the previous section entitled "Patient characteristics not stated."

Sampling errors.—The standard error is primarily a measure of the variability that is attributed to using a value obtained from a sample as an estimate of a population value. In this report it also reflects part of the measurement error. The value that would have been obtained if a complete enumeration of the population had been made will be contained in an interval represented by the sample estimate plus or minus 1 standard error about 68 out of 100 times and plus or minus 2 standard errors about 95 out of 100 times.

The relative standard error is obtained by dividing the standard error by the estimate. The resulting value is multiplied by 100, which expresses the standard error as a percentage of the estimate.

The standard error of one statistic is generally different from that of another, even when the two come from the survey. To derive standard errors that would be applicable to a wide variety of statistics and that could be prepared at a moderate cost, a number of approximations are required. As a result, the figures shown in this appendix provide general relative standard errors for a wide variety of estimates rather than the specific error for a particular statistic.

NOTE: A list of references follows the text.

Approximate relative standard errors and standard errors have been prepared for measuring the variances applicable to (1) estimates of all patient discharges, or first-listed diagnoses, for patient characteristics (e.g., age, sex, and color) and hospital characteristics (e.g., region, bed size, and ownership); and (2) estimates of discharges for single-listed diagnosis and single-listed surgery (operations) for the patient and hospital characteristics.

The selection of the appropriate relative standard error is made as follows:

1. Discharges, or first-listed diagnoses, single-listed diagnosis, and single-listed

- surgery for patient characteristics: Relative standard errors of the estimated number of discharges are obtained from the curve in figure II.
- 2. The approximate standard error of estimated percents, when the characteristic(s) used to form the numerator of the percent is a subclass of the denominator, by patient and hospital characteristic, is presented in table III.
- 3. Approximate standard errors for estimates of discharges by average length of stay applicable to patient and hospital characteristics are presented in table IV.

Figure II. Approximate relative standard errors of estimated numbers of patients discharged, or of first-listed diagnoses, by selected patient and hospital characteristics




Illustration for use of figure II: As shown in table 14 in the West Region, an estimated 324,000 male patients who were under 15 years of age were discharged in 1977 from short-stay, non-Federal hospitals. The relative standard error of this estimate, as read from the curve "All variables" is approximately 7.5 percent. The standard error is approximately 24,000 (7.5 percent of 324,000).

Table III. Approximate standard errors of percents shown in this report for patient discharges: patient characteristics classified by geographic region and for all hospitals

[Standard errors for patient characteristics classified by size of hospital and geographic region by size of hospital are 1½ times and by type of ownership are 2½ times the standard errors shown in this table]

		Es	timate	d perce	ent	
Number of discharges (base of percent in thousands)	2 or 98	4 or 96	10 or 90	20 or 80	30 or 70	50
				r expre ge poin		
100	1.6	2.3	3.6	4.7	5.4	5.9
200	1.2	1.6		3.4	3.8	4.2
1,000	0.7	0.9	1.5	1.9	2.2	2.4
1,000	0.5	0.7	1.1	1.5	1.7	1.9
2,000	0.4	0.5	0.8	1.1	1.2	1.3
6,000	0.2	0.3	0.5	0.6	0.7	0.8
10,000	0.2	0.2	0.4	0.5	0.5	0.6
20,000	0.1	0.2	0.3	0.3	0.4	0.4
40,000	0.1	0.1	0.2	0.2	0.3	0.3

Illustration of use of table III: Figure 1 shows that 30.8 percent of the 11,115,000 white male patients (table 1) discharged during 1977 from all hospitals were 15-44 years of age. Linear interpolation between the values shown in table III yields an approximate standard error of 0.5 percent for an estimate of 30.8 percent with a base of 11,115,000.

Table IV. Approximate standard errors of average lengths of stay shown in this report for patient discharges: patient characteristics cross-classified by geographic region, bed size of hospital, geographic region by bed size of hospital and for all hospitals

[Standard errors for patient characteristics classified by type of ownership are 1½ times the standard errors shown in this table]

Number of discharges (base of average in thousands)	Average length of stay in days						
(base of average in thousands)		6	10	14			
	Stan	dard er	ror in	days			
100	0.3	0.6	8.0	1.1			
200	0.2	0.5	0.7	0.9			
600	0.2	0.4	0.5	0.7			
1,000	0.2	0.3	0.5	0.5			
2,000	0.1	0.2	0.3	0.5			
6,000	0.1	0.2	0.3	0.4			
10,000	0,1	0.2	0.3	0.4			
20,000	0.1	0.1	0.2	0.3			

Illustration of use of table IV: Table 15 shows that the average length of stay was 7.9 days for the estimated 837,000 patients age 45-64 years with a single-listed diagnosis discharged from hospitals in the Northeast Region. Linear interpolation between the values shown in table IV will yield an approximate standard error of 0.4 days for an estimated average length of stay of 7.9 days with a base of 837,000.

APPENDIX II

DEFINITIONS OF CERTAIN TERMS USED IN THIS REPORT

Hospitals and Hospital Characterisitcs

Hospitals.—Short-stay special and general hospitals having six beds or more for inpatient use and an average length of stay of less than 30 days. Federal hospitals and hospital units of institutions are not included.

Bed size of hospital.—Measured by the number of beds, cribs, and pediatric bassinets regularly maintained (set up and staffed for use) for patients; bassinets for newborn infants are not included. In this report the classification of hospitals by bed size is based on the number of beds at or near midyear reported by the hospitals.

Type of ownership of hospital.—The type of organization that controls and operates the hospital. Hospitals are grouped as follows:

Voluntary nonprofit.—Hospitals operated by a church or another nonprofit organization.

Government.—Hospitals operated by State or local governments.

Proprietary.—Hospitals operated by individuals, partnerships, or corporations for profit.

Terms Relating to Hospitalization

Patient.—A person who is formally admitted to the inpatient service of a short-stay hospital for observation, care, diagnosis, or treatment. In this report the number of patients refers to the number of discharges during the year including any multiple discharges of the same individual from one short-stay hospital or more. Infants admitted on the day of birth, directly or by transfer from another medical facility, with or without mention of a disease, disorder, or immaturity are included. All newborn infants, defined as those admitted by birth to the hospital,

are excluded. The terms "patient" and "inpatient" are used synonymously.

Days of care.—The total number of patient days accumulated at time of discharge by patients discharged from short-stay hospitals during a year. A stay of less than 1 day (patient admission and discharge on the same day) is counted as 1 day in the summation of total days of care. For patients admitted and discharged on different days, the number of days of care is computed by counting all days from (and including) the date of admission to (but not including) the date of discharge.

Average length of stay.—The total number of patient days accumulated at time of discharge by patients discharged during the year divided by the number of patients discharged.

Terms Relating to Diagnoses

Discharge diagnosis.—One or more diseases or injuries (or special conditions and examinations without sickness or tests with negative findings) that the attending physician assigns to the medical record of patients. In NHDS all discharge (or final) diagnoses listed on the face sheet (summary sheet) of the medical record for patients discharged from the inpatient service of short-stay hospitals are transcribed in the order listed. Each sample discharge is assigned a maximum of five 3- or 4-digit codes according to the Eighth Revision International Classification of Diseases, Adapted for Use in the United States³ (ICDA), and coding modifications for use in NHDS.

First-listed diagnosis.—The coded diagnosis identified as the principal diagnosis or else listed first on the face sheet of the medical record. The

NOTE: A list of references follows the text.

number of first-listed diagnoses is equivalent to the number of discharges and includes singlelisted diagnosis.

Terms Relating to Surgery

Discharges with surgery.—The estimated number of surgically treated patients discharged from non-Federal short-stay hospitals during the year.

Operation.—One or more surgical operations, procedures, or special treatments that are assigned by the physician to the medical record of patients discharged from the inpatient service of short-stay hospitals. In NHDS all terms listed on the face sheet (summary sheet) of the medical record under the captions "operation," "operative procedures," "operations and/or special treatments," and the like are transcribed in the order listed.

Demographic Terms

Age.—Patient's age refers to age at birthday prior to admission to the hospital inpatient service.

Color.—Patients are classified into two groups, "white" and "all other." The all other classification includes all categories other than white. Mexican and Puerto Rican are included in the white category unless specifically identified as all other.

Geographic region.—Hospitals are classified by location in one of the four geographic regions of the United States which correspond to those used by the U.S. Bureau of the Census.

Region	States included
Northeast	Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, and Pennsylvania
North Central	Michigan, Ohio, Illinois, Indiana, Wisconsin, Minnesota, Iowa, Missouri, North Dakota, South Dakota, Nebraska, and Kansas
South	Delaware, Maryland, District of Columbia, Virginia, North Carolina, South Carolina, Georgia, Florida, Kentucky, Tennessee, Alabama, Missis- sippi, Arkansas, Louisiana, Oklahoma, and Texas
West	Montana, Idaho, Wyoming, Colorado, New Mexico, Ari- zona, Utah, Nevada, Washing- ton, Oregon, California, Hawaii, and Alaska

VITAL AND HEALTH STATISTICS Series

- Series 1. Programs and Collection Procedures.—Reports which describe the general programs of the National Center for Health Statistics and its offices and divisions and data collection methods used and include definitions and other material necessary for understanding the data.
- Series 2. Data Evaluation and Methods Research.—Studies of new statistical methodology including experimental tests of new survey methods, studies of vital statistics collection methods, new analytical techniques, objective evaluations of reliability of collected data, and contributions to statistical theory.
- Series 3. Analytical Studies.—Reports presenting analytical or interpretive studies based on vital and health statistics, carrying the analysis further than the expository types of reports in the other series.
- Series 4. Documents and Committee Reports.—Final reports of major committees concerned with vital and health statistics and documents such as recommended model vital registration laws and revised birth and death certificates.
- Series 10. Data From the Health Interview Survey.—Statistics on illness, accidental injuries, disability, use of hospital, medical, dental, and other services, and other health-related topics, all based on data collected in a continuing national household interview survey.
- Series 11. Data From the Health Examination Survey and the Health and Nutrition Examination Survey.—Data from direct examination, testing, and measurement of national samples of the civilian noninstitutionalized population provide the basis for two types of reports: (1) estimates of the medically defined prevalence of specific diseases in the United States and the distributions of the population with respect to physical, physiological, and psychological characteristics and (2) analysis of relationships among the various measurements without reference to an explicit finite universe of persons.
- Series 12. Data From the Institutionalized Population Surveys.—Discontinued effective 1975. Future reports from these surveys will be in Series 13.
- Series 13. Data on Health Resources Utilization.—Statistics on the utilization of health manpower and facilities providing long-term care, ambulatory care, hospital care, and family planning services.
- Series 14. Data on Health Resources: Manpower and Facilities.—Statistics on the numbers, geographic distribution, and characteristics of health resources including physicians, dentists, nurses, other health occupations, hospitals, nursing homes, and outpatient facilities.
- Series 20. Data on Mortality.—Various statistics on mortality other than as included in regular annual or monthly reports. Special analyses by cause of death, age, and other demographic variables; geographic and time series analyses; and statistics on characteristics of deaths not available from the vital records based on sample surveys of those records.
- Series 21. Data on Natality, Marriage, and Divorce.—Various statistics on natality, marriage, and divorce other than as included in regular annual or monthly reports. Special analyses by demographic variables; geographic and time series analyses; studies of fertility; and statistics on characteristics of births not available from the vital records based on sample surveys of those records.
- Series 22. Data From the National Mortality and Natality Surveys.—Discontinued effective 1975. Future reports from these sample surveys based on vital records will be included in Series 20 and 21, respectively.
- Series 23. Data From the National Survey of Family Growth.—Statistics on fertility, family formation and dissolution, family planning, and related maternal and infant health topics derived from a biennial survey of a nationwide probability sample of ever-married women 15-44 years of age.
- For a list of titles of reports published in these series, write to:

Scientific and Technical Information Branch

National Center for Health Statistics

Public Health Service

Hyattsville, Md. 20782