

Occupational Health Guideline for EPN

INTRODUCTION

This guideline is intended as a source of information for employees, employers, physicians, industrial hygienists, and other occupational health professionals who may have a need for such information. It does not attempt to present all data; rather, it presents pertinent information and data in summary form.

SUBSTANCE IDENTIFICATION

- Formula: $(4\text{-NO}_2\text{C}_6\text{H}_4\text{O})(\text{C}_2\text{H}_5\text{O})\text{C}_6\text{H}_5\text{PSo}$
 $\text{C}_{14}\text{H}_{14}\text{NO}_4\text{PS}$
- Synonyms: O-ethyl O-p-nitrophenyl
thionobenzenephosphonate; O-ethyl O-p-nitrophenyl
benzenephosphonothioate
- Appearance: Yellow solid or brown liquid

PERMISSIBLE EXPOSURE LIMIT (PEL)

The current OSHA standard for EPN is 0.5 milligram of EPN per cubic meter of air (mg/m^3) averaged over an eight-hour work shift.

HEALTH HAZARD INFORMATION

- **Routes of exposure**
EPN can affect the body if it is inhaled, if it comes in contact with the eyes or skin, or if it is swallowed. It may enter the body through the skin.
- **Effects of overexposure**
 1. **Short-term Exposure:** After inhalation of EPN, breathing and eye effects are the first to appear. These include tightness of the chest, wheezing, a bluish discoloration of the skin, small pupils, aching in and behind the eyes, blurring of vision, tearing, runny nose, headache, and watering of the mouth. After swallowing EPN, loss of appetite, nausea, vomiting, abdominal cramps and diarrhea may appear within two hours. After skin absorption, sweating and twitching in the area of absorption may occur, usually within 15 minutes to four hours. With severe intoxication by all routes, in

addition to all the above symptoms, weakness, generalized twitching, and paralysis may occur and breathing may stop. In addition, dizziness, confusion, staggering, slurred speech, generalized sweating, slow heartbeat, convulsions and coma may occur.

2. **Long-term Exposure:** Repeated exposure to EPN may make a person more susceptible to the effects of this and related chemicals. Repeated exposure to concentrations which are too small to produce symptoms after a single exposure may result in the onset of symptoms.

3. **Reporting Signs and Symptoms:** A physician should be contacted if anyone develops any signs or symptoms and suspects that they are caused by exposure to EPN.

- **Recommended medical surveillance**
The following medical procedures should be made available to each employee who is exposed to EPN at potentially hazardous levels:

1. **Initial Medical Examination:**

—A complete history and physical examination: The purpose is to detect pre-existing conditions that might place the exposed employee at increased risk, and to establish a baseline for future health monitoring. Persons with a history of reduced pulmonary function, convulsive disorders, or recent exposure to anticholinesterase agents would be expected to be at increased risk from exposure. Examination of the respiratory system, nervous system, cardiovascular system, and attention to the cholinesterase levels in the blood should be stressed. The skin should be examined for evidence of chronic disorders.

—Cholinesterase determination: EPN causes depressed levels of activity of cholinesterase in the serum and erythrocytes. The cholinesterase activity in the serum and erythrocytes should be determined by using medically acceptable biochemical tests prior to any new period of exposure.

2. **Periodic Medical Examination:** The aforementioned medical examinations should be repeated on an annual basis, with the exception of the cholinesterase determination which should be performed quarterly or at any

These recommendations reflect good industrial hygiene and medical surveillance practices and their implementation will assist in achieving an effective occupational health program. However, they may not be sufficient to achieve compliance with all requirements of OSHA regulations.

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Public Health Service Center for Disease Control
National Institute for Occupational Safety and Health

U.S. DEPARTMENT OF LABOR
Occupational Safety and Health Administration

time overexposure is suspected or signs and symptoms of toxicity occur.

• **Summary of toxicology**

EPN is an anticholinesterase agent; absorption may occur from inhalation of the dust or mist, from skin absorption of solutions, or from ingestion. Signs and symptoms of overexposure are caused by the inactivation of the enzyme cholinesterase, which results in the accumulation of acetylcholine at synapses in the nervous system, skeletal and smooth muscle, and secretory glands. The sequence of the development of systemic effects varies with the route of entry. The onset of signs and symptoms may occur promptly or may be delayed for up to 12 hours. After inhalation, respiratory and ocular effects are the first to appear, often within a few minutes after exposure. Respiratory effects include tightness in the chest and wheezing due to bronchoconstriction and excessive bronchial secretion; laryngeal spasms and excessive salivation may add to the respiratory distress; cyanosis may also occur. Ocular effects include miosis, aching in and behind the eyes (attributed to ciliary spasm), blurring of distant vision, tearing, rhinorrhea, and frontal headache. After ingestion, gastrointestinal effects, such as anorexia, nausea, vomiting, abdominal cramps, and diarrhea appear within 15 minutes to 2 hours. After skin absorption, localized sweating and muscular fasciculations in the immediate area occur, usually within 15 minutes to 4 hours; skin absorption is somewhat greater at higher ambient temperatures and is increased by the presence of dermatitis. With severe intoxication by all routes, an excess of acetylcholine at the neuromuscular junctions of skeletal muscle causes weakness aggravated by exertion, involuntary twitchings, fasciculations, and eventually paralysis; the most serious consequence is paralysis of the respiratory muscles. Effects on the central nervous system include giddiness, confusion, ataxia, slurred speech, Cheyne-Stokes respiration, convulsions, coma, and loss of reflexes. The blood pressure may fall to low levels, and cardiac irregularities including complete heart block may occur; these effects may sometimes be reversed by establishing adequate pulmonary ventilation. Complete symptomatic recovery usually occurs within 1 week; increased susceptibility to the effects of anticholinesterase agents persists for weeks after exposure. Daily exposure to concentrations which are insufficient to produce symptoms following a single exposure may result in the onset of symptoms. Continued daily exposure may be followed by increasingly severe effects. In animals, EPN has produced a demyelination syndrome.

CHEMICAL AND PHYSICAL PROPERTIES

• **Physical data**

1. Molecular weight: 323.3
2. Boiling point (760 mm Hg): Data not available
3. Specific gravity (water = 1): 1.27
4. Vapor density (air = 1 at boiling point of EPN):

Not applicable

5. Melting point: 36 C (97 F)

6. Vapor pressure at 20 C (68 F): 0.0003 mm Hg

7. Solubility in water, g/100 g water at 20 C (68 F):

Insoluble

8. Evaporation rate (butyl acetate = 1): Not applicable

• **Reactivity**

1. Conditions contributing to instability: None.

2. Incompatibilities: Contact with strong oxidizers may cause fires and explosions.

3. Hazardous decomposition products: Toxic gases and vapors (such as oxides of sulfur and nitrogen, phosphoric acid mist, and carbon monoxide) may be released when EPN decomposes.

4. Special precautions: Liquid EPN will attack some forms of plastics, rubber, and coatings.

• **Flammability**

1. Not combustible

• **Warning properties**

1. Odor Threshold: No quantitative information is available concerning the odor threshold of EPN.

2. Eye Irritation Level: EPN is not known to be an eye irritant.

3. Evaluation of Warning Properties: Since no quantitative information is available relating warning properties to air concentrations of EPN, it is treated as a material with poor warning properties. The Pennsylvania *Hygienic Information Guide* for EPN states that the "lack of odor and warning signs, as well as the delayed onset of symptoms, make EPN a dangerous material to work with."

MONITORING AND MEASUREMENT PROCEDURES

• **General**

Measurements to determine employee exposure are best taken so that the average eight-hour exposure is based on a single eight-hour sample or on two four-hour samples. Several short-time interval samples (up to 30 minutes) may also be used to determine the average exposure level. Air samples should be taken in the employee's breathing zone (air that would most nearly represent that inhaled by the employee).

• **Method**

Sampling and analyses may be performed by collection of EPN using a filter with subsequent recovery with iso-octane and gas chromatographic analysis. A detailed analytical method for EPN may be obtained from the National Technical Information Service, U.S. Department of Commerce, Springfield, Virginia 22161, under the title "NIOSH Analytical Methods for Set T" (order number PB 262 404).

RESPIRATORS

• Good industrial hygiene practices recommend that engineering controls be used to reduce environmental

concentrations to the permissible exposure level. However, there are some exceptions where respirators may be used to control exposure. Respirators may be used when engineering and work practice controls are not technically feasible, when such controls are in the process of being installed, or when they fail and need to be supplemented. Respirators may also be used for operations which require entry into tanks or closed vessels, and in emergency situations. If the use of respirators is necessary, the only respirators permitted are those that have been approved by the Mine Safety and Health Administration (formerly Mining Enforcement and Safety Administration) or by the National Institute for Occupational Safety and Health.

- In addition to respirator selection, a complete respiratory protection program should be instituted which includes regular training, maintenance, inspection, cleaning, and evaluation.

PERSONAL PROTECTIVE EQUIPMENT

- Employees should be provided with and required to use impervious clothing, gloves, face shields (eight-inch minimum), and other appropriate protective clothing necessary to prevent any possibility of skin contact with EPN or liquids containing EPN.

- If employees' clothing has had any possibility of being contaminated with EPN or liquids containing EPN, employees should change into uncontaminated clothing before leaving the work premises.

- Clothing which has had any possibility of being contaminated with EPN should be placed in closed containers for storage until it can be discarded or until provision is made for the removal of EPN from the clothing. If the clothing is to be laundered or otherwise cleaned to remove the EPN, the person performing the operation should be informed of EPN's hazardous properties.

- Where there is any possibility of exposure of an employee's body to EPN or liquids containing EPN, facilities for quick drenching of the body should be provided within the immediate work area for emergency use.

- Non-impervious clothing which becomes contaminated with EPN should be removed immediately and not reworn until the EPN is removed from the clothing.

- Employees should be provided with and required to use dust- and splash-proof safety goggles where there is any possibility of EPN or liquids containing EPN contacting the eyes.

- Where there is any possibility that employees' eyes may be exposed to EPN or liquids containing EPN, an eye-wash fountain should be provided within the immediate work area for emergency use.

SANITATION

- Skin that becomes contaminated with EPN should be immediately washed or showered with soap or mild detergent and water to remove any EPN.

- Workers subject to skin contact with EPN or liquids containing EPN should wash with soap or mild detergent and water any areas of the body which may have contacted EPN at the end of each work day.

- Eating and smoking should not be permitted in areas where EPN or liquids containing EPN are handled, processed, or stored.

- Employees who handle EPN or liquids containing EPN should wash their hands thoroughly with soap or mild detergent and water before eating, smoking, or using toilet facilities.

COMMON OPERATIONS AND CONTROLS

The following list includes some common operations in which exposure to EPN may occur and control methods which may be effective in each case:

Operation	Controls
Formulation for use as an insecticide and acaricide	Process enclosure; dust collection; personal protective equipment; general dilution ventilation
Application as an insecticide and acaricide on agricultural crops, vegetables, fruit trees, and ornamentals	Material substitution; personal protective equipment
Manufacture of EPN	Process enclosure; general dilution ventilation; personal protective equipment; local exhaust ventilation

EMERGENCY FIRST AID PROCEDURES

In the event of an emergency, institute first aid procedures and send for first aid or medical assistance.

• Eye Exposure

If EPN or liquids or mists containing EPN get into the eyes, wash eyes immediately with large amounts of water, lifting the lower and upper lids occasionally. Get medical attention immediately. Contact lenses should not be worn when working with this chemical.

• Skin Exposure

If EPN or liquids or mists containing EPN get on the skin, immediately wash the contaminated skin using soap or mild detergent and water. If EPN or liquids or mists containing EPN penetrate through the clothing, remove the clothing immediately and wash the skin using soap or mild detergent and water. Get medical attention immediately.

- **Breathing**

If a person breathes in large amounts of EPN, move the exposed person to fresh air at once. If breathing has stopped, perform artificial respiration. Keep the affected person warm and at rest. Get medical attention as soon as possible.

- **Swallowing**

When EPN or liquids containing EPN have been swallowed and the person is conscious, give the person large quantities of water immediately. After the water has been swallowed, try to get the person to vomit by having him touch the back of his throat with his finger. Do not make an unconscious person vomit. Get medical attention immediately.

- **Rescue**

Move the affected person from the hazardous exposure. If the exposed person has been overcome, notify someone else and put into effect the established emergency rescue procedures. Do not become a casualty. Understand the facility's emergency rescue procedures and know the locations of rescue equipment before the need arises.

SPILL AND DISPOSAL PROCEDURES

- Persons not wearing protective equipment and clothing should be restricted from areas of spills until cleanup has been completed.

- If EPN is spilled, the following steps should be taken:

1. Ventilate area of spill.

2. Cover with soda ash, mix and spray with water. Place in a container of water and allow to stand for two days, then neutralize with 6 molar HCl.

- Waste disposal method:

EPN may be disposed of in sealed containers in a secured sanitary landfill.

REFERENCES

- American Conference of Governmental Industrial Hygienists: "EPN (O-Ethyl-O-p-Nitrophenyl Thionobenzene PhosphoDocumentation of the Threshold Limit

Values for Substances in Workroom Air (3rd ed., 2nd printing), Cincinnati, 1974.

- Christensen, H. E., and Luginbyhl, T. L. (eds.): *NIOSH Toxic Substances List*, 1974 Edition, HEW Publication No. 74-134, 1974.

- Deichmann, W. B., and Gerarde, H. W.: *Toxicology of Drugs and Chemicals*, Academic Press, New York, 1969.

- Goodman, L. S., and Gilman, A.: *The Pharmacological Basis of Therapeutics* (5th ed.), Macmillan, New York, 1975.

- Hamilton, A., and Hardy, H.: *Industrial Toxicology* (3rd ed.), Publishing Sciences Group, Acton, Massachusetts, 1974.

- Hayes, W. J., Jr.: *Clinical Handbook on Economic Poisons, Emergency Information for Treating Poisoning*, U.S. Public Health Service Publication No. 476, U.S. Government Printing Office, Washington, D.C., 1963.

- *Hygienic Information Guide No. 68 - EPN*, Commonwealth of Pennsylvania, Department of Environmental Resources, Bureau of Occupational Health, 1970.

- International Labour Office: *Encyclopedia of Occupational Health and Safety*, McGraw-Hill, New York, 1971.

- Koelle, G. B. (ed.): "Cholinesterases and Anticholinesterase Agents," Vol. 15 of *Handbuch der Experimentellen Pharmakologie*, Springer-Verlag, Berlin, 1963.

- Patty, F. A. (ed.): *Toxicology*, Vol. II of *Industrial Hygiene and Toxicology* (2nd ed. rev.), Interscience, New York, 1963.

- Spector, W. S. (Vols. I, II), Negherbon, W. O. (Vol. III), Grebe, R. M. (Vol. IV), and Dittmer, D. S. (Vol. V) (eds.): *Handbook of Toxicology*, Saunders, Philadelphia, 1956-1959.

- Spencer, E. Y.: *Guide to the Chemicals Used in Crop Protection* (6th ed.), Publication 1093, Research Branch Agriculture, Canada, 1973.

- Thienes, C. H., and Haley, T. J.: *Clinical Toxicology* (5th ed.), Lea and Febiger, Philadelphia, 1972.

RESPIRATORY PROTECTION FOR EPN

Condition	Minimum Respiratory Protection* Required Above 0.5 mg/m ³
Particulate or Vapor Concentration	
5 mg/m ³ or less	Any supplied-air respirator. Any self-contained breathing apparatus.
25 mg/m ³ or less	Any supplied-air respirator with a full facepiece, helmet, or hood. Any self-contained breathing apparatus with a full facepiece.
50 mg/m ³ or less	A Type C supplied-air respirator operated in pressure-demand or other positive pressure or continuous-flow mode.
Greater than 50 mg/m ³ ** or entry and escape from unknown concentrations	Self-contained breathing apparatus with a full facepiece operated in pressure-demand or other positive pressure mode. A combination respirator which includes a Type C supplied-air respirator with a full facepiece operated in pressure-demand or other positive pressure or continuous-flow mode and an auxiliary self-contained breathing apparatus operated in pressure-demand or other positive pressure mode.
Fire Fighting	Self-contained breathing apparatus with a full facepiece operated in pressure-demand or other positive pressure mode.
Escape	Any gas mask providing protection against organic vapors and particulates, including pesticide respirators which meet the requirements of this class. Any escape self-contained breathing apparatus.

*Only NIOSH-approved or MSHA-approved equipment should be used.

**Use of supplied-air suits may be necessary to prevent skin contact while providing respiratory protection from airborne concentrations of EPN; however, this equipment should be selected, used, and maintained under the immediate supervision of trained personnel. Where supplied-air suits are used above a concentration of 50 mg/m³, an auxiliary self-contained breathing apparatus operated in positive pressure mode should also be worn.