Table 20. Reported cases of HIV infection (not AIDS) for male adults and adolescents, by transmission category and race/ethnicity, 2006 and cumulative—45 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting

	200	6	Cumulat	C umulative ^a		
Transmission category	No.	%	No.	%		
		Whit	e, not Hispanic			
Male-to-male sexual contact	11,963	73	55,725	69		
Injection drug use	858	5	5,512	7		
Male-to-male sexual contact and injection drug use	1,092	7	5,578	7		
Hemophilia/coagulation disorder	29	0	355	0		
High-risk heterosexual contact ^b	522	3	2,658	3		
Sex with injection drug user	65	0	544	1		
Sex with person with hemophilia	0	0	8	0		
Sex with HIV-infected transfusion recipient	3	0	33	0		
Sex with HIV-infected person, risk factor not specified	454	3	2,073	3		
Receipt of blood transfusion, blood components, or tissue	14	0	197	0		
Other/risk factor not reported or identified	1,956	12	11,147	14		
Total	16,434	100	81,172	100		
		Black, not Hispanic				
Male-to-male sexual contact	6,203	44	30,148	37		
Injection drug use	1,594	11	11,997	15		
Male-to-male sexual contact and injection drug use	483	3	3,489	4		
Hemophilia/coagulation disorder	10	0	109	0		
High-risk heterosexual contact ^b	1,714	12	10,905	13		
Sex with injection drug user	169	1	1,554	2		
Sex with person with hemophilia	4	0	17	0		
Sex with HIV-infected transfusion recipient	6	0	77	0		
Sex with HIV-infected person, risk factor not specified	1,535	11	9,257	11		
Receipt of blood transfusion, blood components, or tissue	32	0	223	0		
Other/risk factor not reported or identified	4,151	29	24,632	30		
Total	14,187	100	81,503	100		
			Hispanic			
Male-to-male sexual contact	4,094	55	15,623	47		
Injection drug use	993	13	6,407	19		
Male-to-male sexual contact and injection drug use	316	4	1,402	4		
Hemophilia/coagulation disorder	3	0	26	0		
High-risk heterosexual contact ^b	582	8	2,963	9		
Sex with injection drug user	65	1	455	1		
Sex with person with hemophilia	0	0	6	0		
Sex with HIV-infected transfusion recipient	4	0	18	0		
Sex with HIV-infected person, risk factor not specified	513	7	2,484	7		
Receipt of blood transfusion, blood components, or tissue	11	0	55	0		
Other/risk factor not reported or identified	1,475	20	6,977	21		
2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2						

Table 20. Reported cases of HIV infection (not AIDS) for male adults and adolescents, by transmission category and race/ethnicity, 2006 and cumulative—45 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting (cont)

	2006		Cumulati	ve ^a	
Transmission category	No.	%	No.	%	
		Asian	Pacific Islander		
Male-to-male sexual contact	402	70	1,006	63	
Injection drug use	17	3	55	3	
Male-to-male sexual contact and injection drug use	15	3	38	2	
Hemophilia/coagulation disorder	0	0	1	0	
High-risk heterosexual contact ^b	25	4	96	6	
Sex with injection drug user	3	1	10	1	
Sex with person with hemophilia	0	0	0	0	
Sex with HIV-infected transfusion recipient	0	0	3	0	
Sex with HIV-infected person, risk factor not specified	22	4	83	5	
Receipt of blood transfusion, blood components, or tissue	4	1	7	0	
Other/risk factor not reported or identified	109	19	397	25	
Total	572	100	1,600	100	
	Am	erican l	Indian/Alaska Native		
Male-to-male sexual contact	104	61	516	57	
Injection drug use	13	8	101	11	
Male-to-male sexual contact and injection drug use	19	11	110	12	
Hemophilia/coagulation disorder	0	0	0	0	
High-risk heterosexual contact ^b	10	6	53	6	
Sex with injection drug user	2	1	14	2	
Sex with person with hemophilia	0	0	0	0	
Sex with HIV-infected transfusion recipient	0	0	0	0	
Sex with HIV-infected person, risk factor not specified	8	5	39	4	
Receipt of blood transfusion, blood components, or tissue	0	0	2	0	
Other/risk factor not reported or identified	24	14	125	14	
Total	170	100	907	100	
Male-to-male sexual contact	22,912	59	103,679	52	
Injection drug use	3,503	9	24,255	12	
Male-to-male sexual contact and injection drug use	1,946	5	10,704	5	
Hemophilia/coagulation disorder	43	0	495	0	
High-risk heterosexual contact ^b	2,874	7	16,830	8	
Sex with injection drug user	308	1	2,601	1	
Sex with person with hemophilia	4	0	31	0	
Sex with HIV-infected transfusion recipient	13	0	131	0	
Sex with HIV-infected person, risk factor not specified	2,549	7	14,067	7	
Receipt of blood transfusion, blood components, or tissue	61	0	489	0	
Other/risk factor not reported or identified	7,748	20	44,021	22	
Total	39,087 ^C	100	200,473 ^d	100	

Note. See Table 16 for list of 45 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting as of December 2006.

^a From the beginning of the epidemic through December 2006.

b Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

^c Includes 250 males of unknown race or multiple races.

 $^{^{\}rm d}$ Includes 1,838 males of unknown race or multiple races.