

**Table 8. Estimated numbers of persons living with HIV/AIDS, by year and selected characteristics, 2003–2006—33 states and 5 U.S. dependent areas with confidential name-based HIV infection reporting**

	2003	2004	2005	2006
<b>Data for 33 states</b>				
<b>Age at end of year (yrs)</b>				
<13	4,154	3,700	3,294	2,867
13–14	1,130	1,238	1,212	1,139
15–19	3,298	3,627	4,024	4,539
20–24	12,389	12,826	13,411	14,301
25–29	26,167	26,924	28,164	29,600
30–34	49,602	47,577	45,109	44,189
35–39	80,703	77,871	75,842	73,634
40–44	92,040	97,109	100,102	101,764
45–49	71,345	77,490	83,816	90,283
50–54	45,133	51,033	56,640	63,195
55–59	22,793	26,610	31,186	36,008
60–64	10,820	12,642	14,422	16,658
≥65	8,531	9,959	11,657	13,551
<b>Race/ethnicity</b>				
White, not Hispanic	145,081	151,694	158,375	166,000
Black, not Hispanic	202,951	212,425	221,439	231,957
Hispanic	72,612	76,503	80,491	84,720
Asian/Pacific Islander	2,234	2,529	2,858	3,187
American Indian/Alaska Native	1,788	1,889	2,008	2,119
<b>Transmission category</b>				
<b>Male adult or adolescent</b>				
Male-to-male sexual contact	181,771	193,245	205,423	218,676
Injection drug use	57,651	58,149	58,369	59,077
Male-to-male sexual contact and injection drug use	23,965	24,434	24,725	25,085
High-risk heterosexual contact <sup>a</sup>	40,084	42,559	44,915	47,562
Other <sup>b</sup>	3,218	3,278	3,325	3,424
Subtotal	306,689	321,665	336,756	353,825
<b>Female adult or adolescent</b>				
Injection drug use	32,346	32,735	32,973	33,470
High-risk heterosexual contact <sup>a</sup>	80,324	85,348	90,185	95,403
Other <sup>b</sup>	2,036	2,123	2,212	2,321
Subtotal	114,705	120,206	125,370	131,195
<b>Child (&lt;13 yrs at diagnosis)</b>				
Perinatal	6,137	6,154	6,178	6,143
Other <sup>c</sup>	570	575	571	560
Subtotal	6,707	6,730	6,748	6,703
<b>Subtotal for 33 states</b>	<b>428,107</b>	<b>448,605</b>	<b>468,879</b>	<b>491,727</b>
<b>Data for U.S. dependent areas</b>	<b>15,809</b>	<b>16,399</b>	<b>17,150</b>	<b>17,955</b>
<b>Total<sup>d</sup></b>	<b>443,916</b>	<b>465,005</b>	<b>486,028</b>	<b>509,681</b>

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor, but not for incomplete reporting.

Data include persons with a diagnosis of HIV infection (not AIDS), a diagnosis of HIV infection and a later diagnosis of AIDS, or concurrent diagnoses of HIV infection and AIDS.

See Technical Notes for the list of areas that have had laws or regulations requiring confidential name-based HIV infection reporting since at least 2003.

<sup>a</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>b</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

<sup>c</sup> Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

<sup>d</sup> Includes persons of unknown race or multiple races and persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.