

Table 4. Estimated numbers of AIDS cases in children <13 years of age, by year of diagnosis and selected characteristics, 2002–2006 and cumulative—50 states and the District of Columbia


| | Year of diagnosis | | | | | Cumulative ^a |
|---|-------------------|-----------|-----------|-----------|-----------|-------------------------|
| | 2002 | 2003 | 2004 | 2005 | 2006 | |
| Race/ethnicity | | | | | | |
| White, not Hispanic | 14 | 12 | 7 | 4 | 4 | 1,599 |
| Black, not Hispanic | 70 | 46 | 33 | 38 | 30 | 5,654 |
| Hispanic | 18 | 10 | 9 | 8 | 3 | 1,748 |
| Asian/Pacific Islander | 1 | 0 | 1 | 1 | 1 | 54 |
| American Indian/Alaska Native | 1 | 0 | 1 | 0 | 0 | 31 |
| Transmission category | | | | | | |
| Hemophilia/coagulation disorder | 0 | 0 | 0 | 0 | 0 | 226 |
| Mother with documented HIV infection or 1 of the following risk factors | 104 | 70 | 53 | 52 | 37 | 8,508 |
| Injection drug use | 12 | 8 | 5 | 3 | 4 | 3,220 |
| Sex with injection drug user | 4 | 6 | 3 | 2 | 1 | 1,397 |
| Sex with bisexual male | 2 | 0 | 3 | 1 | 1 | 209 |
| Sex with person with hemophilia | 0 | 0 | 0 | 0 | 0 | 35 |
| Sex with HIV-infected transfusion recipient | 0 | 0 | 0 | 0 | 0 | 22 |
| Sex with HIV-infected person, risk factor not specified | 36 | 20 | 20 | 25 | 13 | 1,530 |
| Receipt of blood transfusion, blood components, or tissue | 2 | 1 | 0 | 0 | 0 | 144 |
| Has HIV infection, risk factor not specified | 47 | 34 | 21 | 21 | 17 | 1,951 |
| Receipt of blood transfusion, blood components, or tissue | 2 | 0 | 0 | 0 | 0 | 374 |
| Other/risk factor not reported or identified | 0 | 0 | 0 | 0 | 0 | 36 |
| Total^b | 106 | 70 | 53 | 53 | 38 | 9,144 |

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor, but not for incomplete reporting.

^a From the beginning of the epidemic through 2006.

^b Includes children of unknown race or multiple races. Cumulative total includes 58 children of unknown race or multiple races. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.

Figure 1. Estimated numbers of AIDS cases in children <13 years of age, by year of diagnosis, 1992–2006—50 states and the District of Columbia



Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays, but not for incomplete reporting.