

**Table 3. Estimated numbers of AIDS cases, by year of diagnosis and selected characteristics, 2002–2006 and cumulative—United States and dependent areas**

	Year of diagnosis					Cumulative <sup>a</sup>
	2002	2003	2004	2005	2006	
<b>Data for 50 states and the District of Columbia</b>						
<b>Age at diagnosis (yrs)</b>						
<13	106	70	53	53	38	9,156
13–14	62	74	70	66	73	1,078
15–19	316	302	334	410	401	5,626
20–24	1,399	1,568	1,624	1,675	1,669	36,225
25–29	3,137	3,059	3,196	3,110	3,423	117,099
30–34	5,701	5,521	5,144	4,668	4,349	197,530
35–39	8,217	8,017	7,072	6,479	6,402	213,573
40–44	7,404	7,626	7,703	7,378	7,298	170,531
45–49	5,358	5,617	5,498	5,727	5,628	107,207
50–54	3,271	3,364	3,491	3,527	3,687	59,907
55–59	1,607	1,693	1,836	1,861	2,071	32,190
60–64	872	864	913	872	955	17,303
≥65	682	763	791	727	835	15,074
<b>Race/ethnicity</b>						
White, not Hispanic	11,233	10,948	11,066	10,676	10,929	394,024
Black, not Hispanic	19,246	19,512	18,909	18,081	17,960	409,982
Hispanic	6,712	7,102	6,771	6,823	6,907	161,505
Asian/Pacific Islander	425	450	444	450	519	7,951
American Indian/Alaska Native	177	176	182	170	155	3,345
<b>Transmission category</b>						
<b>Male adult or adolescent</b>						
Male-to-male sexual contact	15,709	16,078	16,054	15,711	16,001	465,965
Injection drug use	5,483	5,275	4,818	4,603	4,410	170,171
Male-to-male sexual contact and injection drug use	2,075	2,029	1,925	1,930	1,803	68,516
High-risk heterosexual contact <sup>b</sup>	4,547	4,471	4,506	4,306	4,558	65,241
Other <sup>c</sup>	253	226	228	237	217	13,893
Subtotal	28,067	28,079	27,532	26,787	26,989	783,786
<b>Female adult or adolescent</b>						
Injection drug use	2,897	2,836	2,810	2,586	2,385	74,718
High-risk heterosexual contact <sup>b</sup>	6,855	7,336	7,113	6,922	7,196	108,252
Other <sup>c</sup>	207	217	217	205	220	6,596
Subtotal	9,959	10,389	10,141	9,713	9,801	189,566
<b>Child (&lt;13 yrs at diagnosis)</b>						
Perinatal	104	70	53	52	37	8,508
Other <sup>d</sup>	2	0	0	0	0	636
Subtotal	106	70	53	53	38	9,144
<b>Region of residence</b>						
Northeast	10,092	10,342	9,412	9,396	9,486	306,241
Midwest	4,125	4,226	4,089	4,371	4,164	101,479
South	17,198	17,630	18,225	16,894	17,104	374,800
West	6,718	6,339	6,000	5,891	6,074	199,978
<b>Subtotal for 50 states and the District of Columbia</b>	<b>38,132</b>	<b>38,538</b>	<b>37,726</b>	<b>36,552</b>	<b>36,828</b>	<b>982,498</b>
<b>Data for U.S. dependent areas</b>	<b>1,058</b>	<b>1,060</b>	<b>921</b>	<b>955</b>	<b>833</b>	<b>31,217</b>
<b>Total<sup>e</sup></b>	<b>39,250</b>	<b>39,690</b>	<b>38,807</b>	<b>37,662</b>	<b>37,852</b>	<b>1,014,797</b>

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor, but not for incomplete reporting.

<sup>a</sup> From the beginning of the epidemic through 2006.

<sup>b</sup> Heterosexual contact with a person known to have, or to be at high risk for, HIV infection.

<sup>c</sup> Includes hemophilia, blood transfusion, perinatal exposure, and risk factor not reported or not identified.

<sup>d</sup> Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

<sup>e</sup> Includes persons of unknown race or multiple races and persons of unknown sex. Cumulative total includes 5,691 persons of unknown race or multiple races, 3 persons of unknown sex, 1,079 persons of unknown state of residence, and 3 persons who were residents of other areas. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.