

RHODE ISLAND – 2007 Profile

HIV/AIDS Epidemic

HIV/AIDS has claimed the lives of over 500,000 Americans. Today, about one million Americans are living with HIV, the virus that causes AIDS, and one guarter of those infected are unaware of their infection.





Figure 1: Source: CDC, HIV/AIDS Surveillance System

Rhode Island reported 2,503 AIDS cases to CDC, cumulatively from the beginning of the epidemic through December 2005. Rhode Island ranked 37th highest among the 50 states in number of reported AIDS cases in 2005. For more, visit: http://www.health.state.ri.us/hiv



Viral Hepatitis


Hepatitis A, B, and C Virus (HAV, HBV, HCV)

Safe and effective vaccines against HBV and HAV have led to dramatic declines in acute disease; for each, estimated incidence in 2005 was the lowest ever recorded. But there is no vaccine for HCV, and chronic HBV and HCV account for over 50% of new cases of chronic liver disease, a leading cause of death.

In Rhode Island, between 2000 and 2005:

- Reports of acute hepatitis A decreased by 41%.
- Reports of acute hepatitis B decreased by 89%.
- Chronic hepatitis C infection reporting to CDC was initiated to improve surveillance.

For more, visit: http://www.health.state.ri.us/hiv/hepatitis.php





Sexually Transmitted Diseases (STDs)




Figure 5: Source: CDC, STD Surveillance System

Figure 6: Source: CDC, STD Surveillance System

Syphilis - Primary and secondary (P&S) syphilis (the stages when syphilis is most infectious) remains a problem in the southern U.S. and some urban areas.

- Rhode Island ranked 17th among 50 states, with 2.2 cases of P&S syphilis per 100,000 persons.
- Between 1996 and 2005, Rhode Island reported 1 case of congenital syphilis, in 2004.

Chlamydia and Gonorrhea - Chlamydial and gonorrheal infections in women are usually asymptomatic and often go undiagnosed. Untreated, these infections can lead to pelvic inflammatory disease, which can cause tubal infertility, ectopic pregnancy, and chronic pelvic pain. In 2005, Rhode Island:

- Ranked 30th among 50 states in chlamydial infections (302.5 per 100,000 persons) and ranked 41st among 50 states in gonorrheal infections (40.5 per 100,000 persons).
- Reported rates of chlamydia among women (428.3 cases per 100,000) that were 2.6 times greater than those among men (166.5 cases per 100,000).

For more, visit: http://www.health.ri.gov/disease/communicable/std/index.php

Program Initiatives Supported by CDC

HIV/AIDS - CDC utilizes a comprehensive approach to HIV prevention that includes surveillance, research, interventions, capacity building, and evaluation. CDC supports 65 state, territorial and local health departments and over 100 community-based organizations to conduct HIV prevention programs. Programs are designed to meet the cultural needs, expectations, and values of the populations they serve, and CDC involves affected communities in the HIV prevention community planning process to ensure that funding goes to those who need it most.

STDs - CDC supports 65 state, territorial, and local health departments to conduct STD prevention programs through Comprehensive STD Prevention System grants. These grants support a community-wide, science-based, interdisciplinary approach to STD prevention that includes behavioral interventions, medical and laboratory services, disease surveillance, outbreak response, professional development, and STD awareness and education campaigns. CDC also assigns staff to state and local health departments.

CDC funding to Rhode Island, 2006	
HIV/AIDS	\$2,487,895
STDs	\$413,195
ТВ	\$485,224
Viral Hepatitis	\$190,376

TB - Through cooperative agreements with 68 state, territorial, and big-city

health departments, CDC supports TB prevention and control activities, including surveillance, case management, and directly observed therapy. These funds also support the identification and evaluation of persons exposed to TB, as well as laboratory services, medical consultation for complex TB cases, and training for state and local TB control staff. CDC also assigns staff to the field to provide direct assistance.

Viral Hepatitis - CDC supports Hepatitis C Coordinators to provide management, networking, and technical expertise for successful integration of hepatitis C prevention activities into existing public health programs. Coordinators facilitate activities in 52 health departments, including: 1) incorporating HCV counseling and testing into public health and clinical settings; 2) training of health care professionals; 3) identifying resources for hepatitis A and B vaccination; and 4) developing referral networks to address the needs of HCV-infected persons.

For More Information

- Rhode Island's programs, services, and statistics, visit: http://www.health.ri.gov
- CDC's programs, services, and statistics, visit: <u>http://www.cdc.gov/nchhstp</u>
- Funding opportunities, conferences, HIV testing resources, and other news, visit: <u>http://www.cdcnpin.org</u>