

*Unspecified and Other Pneumoconioses: Mortality***Unspecified and other pneumoconioses: Number of deaths, death rates (per million population), and years of potential life lost (YPLL) by state, U.S. residents age 15 and over, 1995–2004**

State	No. of Deaths		Death Rate				YPLL to Life Expectancy			
	Deaths	Rank	Crude	Rank	Age-Adjusted	Rank	Total	Rank	YPLL/death	Rank
Alabama	41	12	1.16	11	1.18	12	393	17	9.6	40
Alaska	1	45	0.21	35	0.37	23	15	46	14.6	6
Arizona	13	24	0.33	24	0.31	27	154	23	11.9	22
Arkansas	5	32	0.24	31	0.22	36	75	32	15.0	5
California	49	9	0.19	38	0.22	36	598	7	12.2	17
Colorado	43	11	1.27	9	1.69	9	408	15	9.5	43
Connecticut	7	30	0.26	28	0.23	33	80	31	11.4	24
Delaware	12	26	1.91	8	2.02	7	115	27	9.6	40
District of Columbia	1	45	0.21	35	0.23	33	38	42	38.4	1
Florida	56	7	0.43	20	0.33	25	568	8	10.1	36
Georgia	21	21	0.33	24	0.42	22	300	20	14.3	9
Hawaii	1	45	0.10	48	0.10	48	14	48	14.1	11
Idaho	2	42	0.20	37	0.21	39	30	43	15.2	4
Illinois	95	6	0.98	13	1.01	13	937	6	9.9	38
Indiana	37	14	0.78	15	0.79	15	394	16	10.6	32
Iowa	13	24	0.55	17	0.44	20	115	27	8.8	46
Kansas	5	32	0.24	31	0.23	33	66	33	13.1	14
Kentucky	240	3	7.37	4	7.73	3	2,670	3	11.1	29
Louisiana	8	29	0.23	33	0.24	31	114	29	14.3	9
Maine	2	42	0.19	38	0.18	41	23	45	11.3	25
Maryland	28	19	0.67	16	0.74	16	342	18	12.2	17
Massachusetts	19	22	0.37	23	0.34	24	194	22	10.2	35
Michigan	37	14	0.47	19	0.49	17	416	14	11.2	27
Minnesota	4	36	0.10	48	0.10	48	49	38	12.2	17
Mississippi	4	36	0.18	42	0.19	40	51	36	12.7	15
Missouri	11	27	0.25	30	0.24	31	116	26	10.5	33
Montana	2	42	0.28	27	0.27	29	29	44	14.6	6
Nebraska	-	-	-	-	-	-	-	-	-	-
Nevada	3	41	0.19	38	0.22	36	66	33	21.8	2
New Hampshire	4	36	0.41	22	0.43	21	42	40	10.4	34
New Jersey	34	18	0.51	18	0.49	17	478	12	14.1	11
New Mexico	16	23	1.14	12	1.28	10	144	24	9.0	45
New York	50	8	0.33	24	0.33	25	564	9	11.3	25
North Carolina	27	20	0.42	21	0.45	19	338	19	12.5	16
North Dakota	1	45	0.19	38	0.17	43	9	49	8.5	47
Ohio	178	5	1.98	7	1.91	8	1,919	5	10.8	31
Oklahoma	5	32	0.18	42	0.18	41	39	41	7.7	49
Oregon	4	36	0.15	45	0.14	45	48	39	12.1	20
Pennsylvania	789	1	7.98	3	6.50	4	7,238	1	9.2	44
Rhode Island	-	-	-	-	-	-	-	-	-	-
South Carolina	4	36	0.12	46	0.13	46	84	30	21.1	3
South Dakota	1	45	0.17	44	0.16	44	15	46	14.6	6
Tennessee	41	12	0.90	14	0.94	14	498	11	12.1	20
Texas	37	14	0.23	33	0.28	28	523	10	14.1	11
Utah	47	10	2.84	6	4.01	6	465	13	9.9	38
Vermont	6	31	1.22	10	1.23	11	58	35	9.6	40
Virginia	238	4	4.22	5	4.87	5	2,658	4	11.2	27
Washington	5	32	0.11	47	0.12	47	50	37	10.1	36
West Virginia	607	2	40.99	1	35.80	1	6,740	2	11.1	29
Wisconsin	11	27	0.26	28	0.25	30	131	25	11.9	22
Wyoming	35	17	8.91	2	10.04	2	293	21	8.4	48

- indicates no deaths listed.

NOTE: See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Census Bureau.