

**NORA mining sector and unspecified and other pneumoconioses: Proportionate mortality ratio (PMR) adjusted for age, sex, and race by industry and occupation, U.S. residents age 15 and over, selected states and years, 1990–1999**

Bureau of Census Industry/Occupation		Number of Deaths*	PMR	95% Confidence Interval	
				LCL	UCL
<b>CIC</b>	<b>Industry</b>				
041	Coal mining	833	41.6	38.9	44.6
040	Metal mining	18	4.9	2.9	7.8
050	Nonmetallic mining and quarrying, except fuel	7	2.5	1.0	5.2
	All other industries within this sector	2			
	<b>TOTAL</b>	<b>860</b>			
<b>COC</b>	<b>Occupation</b>				
616	Mining machine operators	756	40.3	37.5	43.3
575	Electricians	12	30.5	15.7	53.2
613	Supervisors, extractive occupations	17	17.7	10.3	28.2
804	Truck drivers	9	14.0	6.4	26.6
889	Laborers, except construction	10	8.7	4.2	15.9
019	Managers and administrators, n.e.c.	5	4.3	1.4	10.1
	All other occupations within this sector	51			
	Occupation not reported within this sector	0			
	<b>TOTAL</b>	<b>860</b>			

NORA - National Occupational Research Agenda    CIC - Census Industry Code    COC - Census Occupation Code

n.e.c. - not elsewhere classified    LCL - lower confidence limit    UCL - upper confidence limit

\* The total number of unspecified and other pneumoconioses deaths was 1,440 in these same selected states and years, and the comparable number of unspecified and other pneumoconioses deaths in the entire United States for this same time period was 3,384.

NOTE: See appendices for source description, methods, ICD codes, industry and occupation codes, industrial sector groupings, and list of selected states and years.

SOURCE: National Center for Health Statistics multiple cause-of-death data.