



Published October 21, 1994, for  
1993 / Vol. 42 / No. 53

# MMWR

MORBIDITY AND MORTALITY WEEKLY REPORT

---

- 1 Summaries of Notifiable Diseases in the United States, 1993
- 13 Graphs and Maps for Selected Notifiable Diseases in the United States
- 65 Historical Summary Tables Covering the Period 1944-1993

## Summary of Notifiable Diseases, United States 1993

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES  
Public Health Service  
Centers for Disease Control  
and Prevention (CDC)  
Atlanta, Georgia 30333



The statistical summary of notifiable diseases in the United States is published to accompany each volume of the *Morbidity and Mortality Weekly Report* by the Centers for Disease Control and Prevention (CDC), Public Health Service, U.S. Department of Health and Human Services, Atlanta, GA 30333.

SUGGESTED CITATION

Centers for Disease Control and Prevention. Summary of notifiable diseases, United States, 1993. MMWR 1993;42(53): [inclusive page numbers].

Centers for Disease Control and Prevention..... David Satcher, M.D., Ph.D.  
*Director*

The material in this report was collected and forwarded to CDC by the 56 state and territorial epidemiologists and was received and prepared by:

Epidemiology Program Office..... Stephen B. Thacker, M.D., M.Sc.  
*Director*

Richard A. Goodman, M.D., M.P.H.  
*Editor, MMWR Series*

Division of Surveillance  
and Epidemiology..... Donna F. Stroup, Ph.D., M.Sc.  
*Director*

Scientific Information and Communications Program  
Public Health Publications Branch ..... Suzanne M. Hewitt, M.P.A.  
*Chief*

Rachel J. Wilson  
*Project Editor*

Information Resources Management Branch ..... Elizabeth E. Rubery  
*Chief*

Peter M. Jenkins  
*Visual Information Specialist*

Use of trade names is for identification only and does not imply endorsement by the Public Health Service or the U.S. Department of Health and Human Services.

Copies can be purchased from Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402-9325. Telephone: (202) 783-3238.

The following CDC staff members prepared this report:

Denise T. Koo, M.D., M.P.H.  
Andrew G. Dean, M.D., M.P.H.

Ruth W. Slade

Carol M. Knowles

Deborah A. Adams

Wanda K. Fortune

Patsy A. Hall

Robert F. Fagan

Barbara Panter-Connah

Harry R. Holden

Gerald F. Jones

Clarence Lee Maddox

*Division of Surveillance and Epidemiology*

*Epidemiology Program Office*

**Consultant**

Willie J. Anderson

*Office of the Vice President for Health Affairs*

*Emory University*

## Table of Contents

<b>Foreword</b> .....	ii
<b>Background</b> .....	iii
<b>Data Sources</b> .....	iv
<b>1993 Highlights for Selected Diseases</b> .....	vi
<b>Selected Bibliography</b> .....	xi

### Part 1

Summaries of Notifiable Diseases in the United States, 1993 .....	1
Reported Cases, by Month, 1993.....	3
Reported Cases, by Geographic Division and Area, 1993 .....	4
Reported Cases, by Age Group, 1993 .....	10
Reported Cases, by Race, 1993.....	11
Reported Cases, by Ethnicity, 1993 .....	12

### Part 2

Graphs and Maps for Selected Notifiable Diseases in the United States.....	13
--	----

### Part 3

Historical Summary Tables Covering the Period 1944–1993 .....	65
Notifiable Diseases —	
Summary of Reported Cases, United States, 1984–1993 .....	67
Summary of Reported Cases, per 100,000 Population, United States, 1984–1993 .....	68
Summary of Reported Cases, United States, 1974–1983.....	69
Summary of Reported Cases, United States, 1964–1973.....	70
Summary of Reported Cases, United States, 1954–1963.....	71
Summary of Reported Cases, United States, 1944–1953.....	72
Deaths from Specified Notifiable Diseases, United States, 1982–1991 .....	73
<b>State and Territorial Epidemiologists and Laboratory Directors</b> .....	inside back cover

## Foreword

### *MMWR Summary of Notifiable Diseases, United States, 1993*

This publication contains summary tables of the official statistics for the occurrence of nationally notifiable diseases in the United States for calendar year 1993. This information is collected and compiled from reports to the National Notifiable Diseases Surveillance System (NNDSS).

Part 1 contains information on morbidity for each of the 49 currently notifiable conditions. In all tables, leprosy is listed as Hansen disease, typhus fever (flea-borne) as murine typhus fever, and typhus fever (tick-borne) as Rocky Mountain spotted fever (RMSF). The tables show the number of cases of notifiable diseases reported to CDC for 1993, as well as the distribution of cases by month and geographic location, and by patient's age, race, and ethnicity.

Part 2 contains graphs and maps depicting summary data for many of the notifiable conditions described in tabular form in Part I.

Part 3 includes tables showing the number of cases of notifiable diseases reported to CDC and to the National Office of Vital Statistics since 1944. It also includes a table on deaths associated with specified notifiable diseases reported to the National Center for Health Statistics, CDC, for the period 1982–1991.

## Background

As of January 1, 1994, 49 infectious diseases were designated as notifiable at the national level. *A notifiable disease is one for which regular, frequent, and timely information on individual cases is considered necessary for the prevention and control of the disease.* This section briefly summarizes the history of national notifiable disease reporting in the United States.

In 1878, Congress authorized the U.S. Public Health Service (PHS) to collect morbidity reports on cholera, smallpox, plague, and yellow fever from U.S. consuls overseas; this information was to be used for instituting quarantine measures to prevent the introduction and spread of these diseases into the United States. In 1879, a specific Congressional appropriation was made for the collection and publication of reports of these notifiable diseases. The authority for weekly reporting and publication was expanded by Congress in 1893 to include data from states and municipal authorities. To increase the uniformity of the data, Congress enacted a law in 1902 directing the Surgeon General to provide forms for the collection and compilation of data and for the publication of reports at the national level. In 1912, state and territorial health authorities—in conjunction with PHS—recommended weekly telegraphic reporting of five infectious diseases and monthly reporting by letter of 10 additional diseases. The first annual summary of *The Notifiable Diseases* in 1912 included reports of 10 diseases from 19 states, the District of Columbia, and Hawaii. By 1928, all states, the District of Columbia, Hawaii, and Puerto Rico were participating in national reporting of nearly 30 specified conditions. At their meeting in 1950, the State and Territorial Health Officers authorized a conference of state epidemiologists whose purpose was to determine which diseases should be reported to PHS. CDC assumed responsibility for the collection and publication of data on nationally notifiable diseases in 1961.

Public health officials at state health departments and CDC continue to collaborate in determining which diseases should be nationally notifiable; the Council of State and Territorial Epidemiologists (CSTE) makes recommendations annually for additions and deletions to the national notifiable disease list on the basis of CDC suggestions. However, reporting of nationally notifiable diseases to CDC by the states is voluntary. Reporting is currently mandated only at the state level. The list of diseases that are considered notifiable, therefore, varies slightly by state. All states generally report the internationally quarantinable diseases (i.e., cholera, plague, and yellow fever) in compliance with the World Health Organization's International Health Regulations.

The list of nationally notifiable diseases is revised periodically. Diseases are added to the list as new pathogens emerge; diseases are deleted as their incidence declines. During 1993, 49 infectious diseases were considered notifiable at the national level and were reported to CDC; 41 were reported on a weekly basis, and eight were reported monthly. CSTE will review the principles of notifiable disease reporting in late fall 1994.

## Data Sources

Provisional data on the reported occurrence of notifiable diseases are published routinely in *MMWR*; these data are compiled in final form in this summary. Notifiable disease reports published in the annual *MMWR Summary of Notifiable Diseases* are the authoritative and archival counts of cases. Data from surveillance records for selected diseases, which are useful for detailed epidemiologic analyses, are published on a periodic basis. Data appearing in those surveillance reports may not agree exactly with reports published in *MMWR* due to differences in the timing of reports or the source of the data or due to refinements in case definitions.

Data in this summary are primarily derived from electronic reports transmitted to the Epidemiology Program Office, CDC, by state and territorial health departments through the National Electronic Telecommunications System for Surveillance (NETSS). Final data for other diseases are from surveillance program records of the CDC programs listed below. Requests for further information regarding these data should be directed to the appropriate source.

### **National Center for Health Statistics**

Office of Vital and Health Statistics Systems (deaths from selected notifiable diseases)

### **National Center for Infectious Diseases**

Division of Bacterial and Mycotic Diseases (cholera and toxic-shock syndrome)  
Division of HIV/AIDS (human immunodeficiency virus/acquired immunodeficiency syndrome)

Division of Vector-Borne Infectious Diseases (arboviral infections of the central nervous system and plague)

Division of Viral and Rickettsial Diseases (animal rabies)

### **National Center for Prevention Services**

Division of Sexually Transmitted Diseases and HIV Prevention (gonorrhea, syphilis, chancroid, granuloma inguinale, and lymphogranuloma venereum)

Division of Tuberculosis Elimination (tuberculosis)

### **National Immunization Program**

Epidemiology and Surveillance Division (poliomyelitis)

Disease totals for the United States, unless otherwise stated, do not include data for American Samoa, Guam, Puerto Rico, the Virgin Islands, and the Commonwealth of the Northern Mariana Islands (CNMI).

Data on notifiable diseases before 1960 are obtained from publications of the National Office of Vital Statistics.

Population estimates for states are based on the July 1, 1993, post-censal estimates made by the U.S. Department of Commerce, Bureau of the Census, Population Division, Population Estimates Branch, Press Release CB93-219.

Population estimates for territories are from the 1990 census, U.S. Department of Commerce, Bureau of the Census, Press Releases CB91-142, 242, 243, 263, and 276.

Rates in the *1993 Summary of Notifiable Diseases* were based on data for the U.S. total resident population. However, population data from states in which diseases were not notifiable or not available were excluded from rate calculations.

### Interpreting Data

Although the data reported in this summary are important for analyzing disease trends and determining relative disease burdens, these data must be interpreted with caution. Some diseases, such as plague and rabies, that cause severe clinical illness and are associated with serious consequences probably are diagnosed and reported accurately. However, diseases that are clinically mild and infrequently associated with serious consequences (e.g., salmonellosis and mumps) are less likely to be reported. Additionally, subclinical cases are seldom detected except in the course of special studies. The degree of completeness of reporting also is influenced by the diagnostic facilities available, the control measures in effect, and the interests and priorities of state and local officials responsible for disease control and surveillance. Finally, factors such as the introduction of new diagnostic tests and the discovery of new disease entities may cause changes in disease reporting that are independent of the true incidence of disease.

Public health surveillance data are published for selected racial and ethnic population groups because these variables may be risk markers for certain notifiable diseases. Risk markers can identify potential risk factors for investigation in future studies. Data on race and ethnicity also can be useful for identifying groups to target for prevention efforts. Year 2000 Objectives for racial and ethnic groups have been established for several of the notifiable diseases.

EXPLANATION OF SYMBOLS USED IN TABLES, GRAPHS, AND MAPS	
Data not available .....	NA
No reported cases.....	-
Report of disease not required by state health department (not notifiable).....	NN



## 1993 HIGHLIGHTS FOR SELECTED DISEASES

This section highlights important developments in the reported occurrences of certain notifiable and non-notifiable infectious diseases. Those diseases that currently are not nationally notifiable are highlighted under the subheading "Emerging Infections."

### AIDS

In 1993, the number of AIDS cases reported to CDC increased 127% from the number reported in 1992. The large increase in the number of cases was largely due to changes in the 1993 AIDS surveillance case definition. As of January 1, 1993, HIV-infected persons with additional clinical conditions, as well as those with markers of severe immunosuppression, were defined as having AIDS.

### Botulism

In 1993, two restaurant-associated outbreaks of botulism occurred in which unusual vehicles of transmission were implicated. In one of the outbreaks, which occurred in Georgia, the toxin was transmitted by a canned cheese sauce; in the other outbreak in Texas, toxin was transmitted by a dip containing baked potatoes.

### Cholera

Epidemics of cholera caused by a new pathogen, *Vibrio cholerae* O139, occurred in late 1992 and early 1993 in southern Asia; in the United States, the first recognized case of infection caused by *Vibrio cholerae* O139 occurred during 1993 in a California resident who had visited India in January 1993.

### Diphtheria

Outbreaks of diphtheria were reported from the Newly Independent States of the former Soviet Union (especially Russia and Ukraine) and Ecuador. No cases were reported last year among U.S. travelers to these countries.

### *Haemophilus influenzae*

The rate of *Haemophilus influenzae* disease as reported through the National Notifiable Diseases Surveillance System (NNDSS) has continued to decline. Rates decreased 95% between 1987 and 1993. Data collected by active surveillance from selected sites indicate that the decline is primarily in *Haemophilus influenzae* type b (Hib) disease among children <5 years of age. This decline is associated with the use of the newly licensed Hib conjugate vaccines.

### Hepatitis B

The reported incidence of hepatitis B decreased 59% from 1985 through 1993. This decline was caused by decreases in the number of cases reported among homosexual men between 1985 and 1989 (61%) and in the number reported among injecting-drug users from 1989 through 1992 (51%). These decreases are thought to result from an increase in AIDS awareness, which has resulted in behavioral changes (e.g., safer sex and needle-using practices).

### **Hepatitis, Non-A, Non-B**

The steady increase in the number of cases of non-A, non-B hepatitis reported to NNDSS since 1990 is not thought to be caused by an increase in incidence, but rather caused by the screening of blood for the presence of antibody to the hepatitis C virus (anti-HCV, available beginning in May, 1990). Persons testing positive for anti-HCV, regardless of the presence of illness, have been reported as having non-A, non-B hepatitis. Surveillance at sentinel sites has actually shown a steady decrease in the incidence of non-A, non-B hepatitis, with a decline of >50% from 1988 through 1993. This decline is primarily attributable to a 58% decrease in the number of reported cases among injecting-drug users.

### **Legionnaires' Disease**

Although only 1,280 cases of Legionnaires' disease were reported through NNDSS in 1993, data from recent prospective pneumonia studies suggest that between 10,000 and 15,000 cases occur each year.

### **Lyme Disease**

In 1993, the number of cases of Lyme disease reported to CDC nationwide decreased by 15% from 1992. Hunterdon, New Jersey meanwhile experienced an increase in reported cases of Lyme disease from 75 cases in 1992 to 206 cases in 1993; upon investigation, the increase in reported cases was found to be due to both increased reporting by physicians and an increase in disease incidence.

### **Malaria**

In August 1993, three persons without recent bloodborne exposure and who had not recently traveled were diagnosed with *Plasmodium falciparum* malaria infection in New York City. The results of the epidemiologic investigation suggest local acquisition through mosquito-borne transmission.

### **Measles**

The total number of measles cases reported to CDC in 1993—312—was the lowest number ever recorded in the United States. This reduction in reported measles cases may reflect cyclical changes in measles incidence as well as increases in measles vaccination coverage among preschool-aged children, increased use of a second dose of measles vaccine among school- and college-aged persons, and increased efforts to control measles throughout the Western Hemisphere.

### **Meningococcal disease**

Several clusters of group C disease were recognized in 1993, resulting in mass vaccinations with meningococcal vaccine. Because approximately one half of severe meningococcal disease in this country is caused by serogroup B and because the vaccine is only effective for disease caused by serogroups A, C, Y, and W, the clusters illustrate the importance of serogroup determination and reporting for all cases of invasive *Neisseria meningitidis*.

### **Mumps**

The total number of mumps cases reported in 1993—1,692—was the lowest number ever reported in the United States.

### **Pertussis**

The 6,503 pertussis cases reported in the United States in 1993 represent the highest annual total since 1967. Of 1,347 cases of pertussis among children 7 months to 4 years of age with known vaccination status, 630 (46.8%) had received fewer than three doses of DTP vaccine—the minimum number of doses necessary for clinical protection.

### **Rubella**

The total number of rubella cases reported to the CDC remained low in 1993 (N=192), although the number increased by 19% over the record low number of cases reported in 1992.

### **Sexually Transmitted Diseases**

Nationally, the rates for gonorrhea and primary and secondary syphilis have declined since 1990, reaching low points in 1993 that were below or approaching the Year 2000 Objectives (1) ( $\leq 225$  cases gonorrhea and  $\leq 10$  cases primary and secondary syphilis per 100,000 persons). However, rates for both diseases remained higher than the Year 2000 Objectives for certain population subgroups: adolescents and young adults, minorities (especially blacks), and persons living in the southern United States.

### ***Shigella***

The number of cases of *Shigella* reported through NNDSS increased 35% from 1992 to 1993, with increases of 27%–405% reported from nine states. This increase was also reported through the national *Shigella* surveillance system, which receives reports of laboratory isolates of *Shigella* through the Public Health Laboratory Information System (PHLIS); 91% of *Shigella* isolates reported through PHLIS in 1993 were *Shigella sonnei*.

### **Tuberculosis**

Between 1992 and 1993, the number of reported tuberculosis (TB) cases in the United States decreased from 26,673 to 25,313. This decrease may be associated with the effectiveness of prevention and control measures implemented during the period 1989–1993. However, two other factors also may be responsible for the decrease, including a) delayed reporting caused by use of the new TB surveillance reporting form and the change from paper records to a computerized system and b) underreporting because of modification of the AIDS surveillance case definition in January 1993.

## **EMERGING INFECTIONS**

NOTE: The term “emerging infectious diseases” refers to those “infectious diseases whose incidence in humans has increased within the past two decades or threatens to increase in the near future” (2). These can be new or previously unrecognized infectious diseases, reemerging diseases, or infectious diseases that have developed resistance to previously effective antimicrobial drugs.

### **Coccidioidomycosis**

The outbreak of coccidioidomycosis in California that began in 1991 continued in 1993. From an annual average number of 428 cases reported per year in California during the period 1981–1990, 1,200 cases were reported in 1991, 4,516 in 1992, and

4,137 in 1993, 70% of which were reported from Kern County in central California. Key factors that may be associated with the ongoing outbreak include weather conditions that are conducive to the growth and spread of *Coccidioides immitis* (e.g., protracted drought followed by heavy rains), activities that disturb the soil and facilitate airborne spread of the organism, and an increasing population of persons who are susceptible to the organism because of migration from areas where coccidioidomycosis is not endemic.

### **Cryptosporidiosis**

In spring 1993, a municipal water supply in Milwaukee, Wisconsin, contaminated with *Cryptosporidium* caused the largest recognized outbreak of waterborne illness in the history of the United States. More than 400,000 persons became ill, 4,400 of whom required hospitalization.

### ***Escherichia coli* O157:H7**

In 1993, an outbreak of *E. coli* O157:H7 affected more than 500 people in four western states, resulting in 56 cases of hemolytic uremic syndrome and four deaths. Because of this outbreak, many clinical laboratories began screening stool samples for *E. coli* O157:H7, which resulted in the identification of many more cases and outbreaks. In May 1993, the Council of State and Territorial Epidemiologists (CSTE) passed a resolution recommending that *E. coli* O157:H7 infection be made reportable by all states and territories.

### **Group A Streptococcal Disease**

During 1993, CDC surveillance for invasive group A streptococcal infections consisted of a passive nationwide surveillance system. This system operated through the collection of isolates from normally sterile sites and the collection of case reports. Although current data on incidence and trends for invasive disease, streptococcal toxic shock syndrome, and necrotizing fasciitis are not available, population-based active surveillance for these infections has begun in several geographic areas and will be expanded in 1994 as part of surveillance for emerging infectious diseases.

### **Hantavirus Pulmonary Syndrome**

Hantavirus Pulmonary Syndrome (HPS), a newly recognized illness characterized by an influenza-like prodrome followed by the acute onset of respiratory failure, was first identified in the southwestern United States in June 1993 during the investigation of a cluster of unexplained deaths. A new hantavirus (Sin Nombre virus) and a rodent reservoir for the virus (the deer mouse [*Peromyscus maniculatus*]) were identified. As of August 31, 1994, national surveillance for HPS, initiated by CDC in coordination with CSTE, has identified 91 confirmed cases of HPS (with 48 deaths) in 20 states (case fatality rate: 53%).

### **Drug-Resistant Pneumococcus**

The increasing incidence of drug-resistant *Streptococcus pneumoniae* (DRSP) strains in the United States has created an emerging public health challenge. CDC surveillance data from 1992 indicated that the prevalence of pneumococcal strains that are highly resistant to penicillin increased 60-fold (from 0.02% to 1.3%) when compared with the prevalence of isolates collected from 1979 through 1987. CDC, CSTE, and the infectious diseases and microbiology communities are developing

recommendations for the surveillance of DRSP infections. This surveillance data will be used to determine optimal empiric treatment regimens for pneumococcal infections.

### **Vancomycin-Resistant Enterococci (VRE)**

From 1989 through 1993, the percentage of nosocomial enterococci resistant to vancomycin reported from hospitals participating in the National Nosocomial Infections Surveillance System increased from 0.3% to 7.9%. During this period, numerous VRE outbreaks (occurring primarily among immunocompromised patients) were reported. Because of the public health importance of the emergence of VRE, CDC has published draft *Guidelines for Preventing the Spread of Vancomycin Resistance* and is conducting studies to assess the effectiveness of these guidelines in preventing disease transmission.

### *References*

1. Public Health Service. Healthy people 2000: national health promotion and disease prevention objectives—full report, with commentary. Washington, DC: US Department of Health and Human Services, Public Health Service, 1991:152; DHHS publication no. (PHS)91-50212.
2. Institute of Medicine. Emerging infections: microbial threats to health in the United States. Washington, DC: National Academy Press, 1992.

## Bibliography

### General

- Benenson AS. Control of communicable diseases in man. 15th ed. Washington, DC: American Public Health Association, 1990.
- CDC. Case definitions for public health surveillance. MMWR 1990;39(No. RR-13).
- CDC. Mandatory reporting of infectious diseases by clinicians, and mandatory reporting of occupational diseases by clinicians. MMWR 1990;39(No. RR-9).
- CDC. National electronic telecommunications system for surveillance—United States, 1990–1991. MMWR 1991;40:502.
- CDC. Update: graphic method for presentation of notifiable disease data—United States, 1991. MMWR 1991;40:124–5.
- CDC. Use of race and ethnicity in public health surveillance. MMWR 1993;42(No. RR-10).
- CDC. Update: changes in notifiable disease surveillance data—United States, 1992–1993. MMWR 42(42):824–6.
- CDC. Sexually transmitted disease surveillance, 1993. Atlanta: US Department of Health and Human Services, Public Health Service, 1994.
- Stroup DF, Wharton M, Kafadar K, Dean AG. An evaluation of a method for detecting aberrations in public health surveillance data. Am J Epidemiol 1993;137(3):373–80.
- Teutsch SM, Churchill RE, eds. Principles and practice of public health surveillance. New York: Oxford University Press, 1994.
- Thacker SB, Choi K, Brachman PS. The surveillance of infectious diseases. JAMA 1983; 249:1181–5.

### AIDS

- CDC. 1993 revised classification for HIV infection and expanded surveillance case definition for AIDS among adolescents and adults. MMWR 1992;41(No. RR-17):1–19.
- CDC. Heterosexually acquired AIDS—United States, 1993. MMWR 1994;43(9):155–60.
- CDC. Update: impact of the expanded AIDS case definition for adolescents and adults on case reporting—United States, 1993. MMWR 1993;43(9):160–70.
- CDC. Recommendations of the U.S. Public Health Service Task Force on the use of zidovudine to reduce perinatal transmission of human immunodeficiency virus. MMWR 1994;43(No. RR-11):1–20.

### Amebiasis

- Katzenstein DA. Drug treatment of amebiasis. In: Peterson PK, Verhoef J, eds. Antimicrobial agents annual 2. New York: Elsevier Science Publishers, 1987:247–52.
- McAuley JB, Herwaldt BL, Stokes SL, et al. Diloxanide furoate for treating asymptomatic *Entamoeba histolytica* cyst passers: 14 years' experience in the United States. Clin Infect Dis 1992;15:464–8.
- Ravdin JI (ed.). Amebiasis: human infection by *Entamoeba histolytica*. New York: John Wiley and Sons, Inc., 1988.

### Anthrax

- Brachman PS. Anthrax. In: Hoepflich PD, Jordan MC, Roland AR, eds. Infectious diseases. 5th ed. Philadelphia: JB Lippincott Co., 1994;1003–8.

### Arboviral Infections

#### (St. Louis encephalitis, Eastern equine encephalitis, Western equine encephalitis, and California serogroup viruses)

- CDC. Arbovirus diseases—United States, 1993. MMWR 1994;43(21):385–7.
- CDC. Arboviral surveillance—United States, 1990. MMWR 1990;39(35):593–8.
- Tsai TF. Arboviral infections in the United States. Infect Dis Clin North Am 1991;5:73–102.

### Aseptic Meningitis

- CDC. Aseptic meningitis—New York State and United States, weeks 1–36, 1991. MMWR 1991;40(45):773–5.
- Moore M. Enteroviral disease in the United States, 1970–1979. J Infect Dis 1982;146:103–8.
- Morens DM, Zweighaft RM, Bryan JA. Nonpolio enterovirus disease in the United States, 1971–1975. Int J Epidemiol 1979;8:49–54.

Wilfert CM, Lehrman SN, Katz SL. Enteroviruses and meningitis. *Pediatr Infect Dis J* 1983; 2:333-41.

#### **Botulism**

St. Louis ME. Botulism. In: Evans AS, Brachman PS, eds. *Bacterial infections of humans: epidemiology and control*. 2nd ed. New York and London: Plenum Medical Book Co., 1991;115-31.

St. Louis ME, Peck SHS, Bowering D, et al. Botulism from chopped garlic: delayed recognition of a major outbreak. *Ann Intern Med* 1988;108:363-8.

Woodruff BA, Griffin PM, McCroskey LM, et al. Clinical and laboratory comparison of botulism from toxin types A, B, and E in the United States 1975-1988. *J Infect Dis* 1992;166:1281-6.

#### **Brucellosis**

Kaufmann AF, Fox MD, Boyce JM, et al. Airborne spread of brucellosis. *Ann NY Acad Sci* 1980;353:105-14.

Staskiewicz J, Lewis CM, Colville J, Zervos M, Band J. Outbreak of *Brucella melitensis* among microbiology laboratory workers in a community hospital. *J Clin Microbiol* 1991;29:287-90.

Young EJ. Human brucellosis. *Rev Infect Dis* 1983;5:821-42.

#### **Chancroid**

CDC. Chancroid in the United States, 1981-1990: evidence for underreporting of cases. *MMWR* 1992;41(No. SS-3):57-61.

Schmid GP. Chancroid. In: Evans AS, Brachman PS, eds. *Bacterial infections of humans: epidemiology and control*. 2nd ed. New York: Plenum Medical Book Company, 1991;173-8.

Schmid GP. Chancroid and granuloma inguinale. In: Kelley WB, ed. *Textbook of internal medicine*. 2nd ed. Philadelphia: JB Lippincott Co, 1991:1412-4.

#### **Cholera**

Blake PA. Epidemiology of cholera in the Americas. *Gastroenterol Clin North Am* 1993;22:639-60.

Swerdlow DL, Ries AA. *Vibrio cholerae* non-01—the eighth pandemic? *Lancet* 1993;342:382-3.

Wachsmuth IK, Blake PA, Olsvik O, eds. *Vibrio cholerae* and cholera: molecular to global perspectives. Washington, DC: American Society for Microbiology, 1994.

World Health Organization. Guidelines for cholera control. Geneva: World Health Organization, 1993.

#### **Congenital Syphilis**

CDC. Surveillance for geographic and secular trends in congenital syphilis—United States, 1983-1991. *MMWR* 1993;42(No. SS-6):59-71.

Zenker PN. Syphilis. In: Schdlower M, Shafer M-A, eds. *Adolescent medicine: state of the art reviews*. Philadelphia: Hanley and Belfus, Inc., 1990;1:511-29.

Zenker P. Congenital syphilis reporting (from the CDC). *Sex Transm Dis* 1991;18:44-54.

Zenker PN, Berman SM. Congenital syphilis: trends and recommendations for evaluation and management. *Pediatr Infect Dis J* 1991;10(7):516-22.

#### **Diphtheria**

CDC. Diphtheria outbreak—Russian Federation, 1990-1993. *MMWR* 1993;42:840-1,847.

Chen RT, Broome CV, Weinstein RA, Weaver R, Tsai TF. Diphtheria in the United States, 1971-81. *Am J Public Health* 1985;75:1393-7.

Farizo KM, Strebel PM, Chen RT, Kimbler A, Cleary TJ, Cochi SL. Fatal respiratory disease due to *Corynebacterium diphtheriae*: case report and review of guidelines for management, investigation, and control. *Clin Infect Dis* 1993;16:59-68.

#### **Gonorrhea**

CDC. Plasmid-mediated antimicrobial resistance in *Neisseria gonorrhoeae*—U.S. 1988 and 1989. *MMWR* 1990;39(17):284-7,293.

CDC. Special focus: surveillance for sexually transmitted diseases. *MMWR* 1993;42(No. SS-3): 1-11, 29-39.

Gershman KA, Rolfs RT. Diverging gonorrhea and syphilis trends in the 1980s: are they real? *Am J Public Health* 1991;81(10):1263-7.

### **Granuloma Inguinale**

Schmid GP. Chancroid and granuloma inguinale. In: Kelley WB, ed. Textbook of internal medicine. 2nd ed. Philadelphia: JB Lippincott Co, 1989:1555–6.

### ***Haemophilus influenzae***

Adams WG, Deaver KA, Cochi SL, et al. Decline of childhood *Haemophilus influenzae* type b (Hib) disease in the Hib vaccine era. JAMA 1993;269:221–6.

CDC. Recommendations for use of *Haemophilus b* conjugate vaccines and a combined diphtheria, tetanus, pertussis, and *Haemophilus b* vaccine: recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR 1993;42(No. RR-13).

CDC. Progress toward elimination of *Haemophilus influenzae* type b disease among infants and children—United States, 1987–1993. MMWR 1994;43(8):144–8.

### **Hansen Disease (Leprosy)**

Meyer WM. Leprosy. In: Strickland GT. Hunter's tropical medicine. WB Saunders, 1991:483–94.

Neill MA, Hightower AW, Broome CV. Leprosy in the United States 1971–1981. J Infect Dis 1985;152:1064–9.

Noordeen SK, Bravo LL, Sundaresan TK. Estimated number of leprosy cases in the world. Bull World Health Organ 1992;70(1):7–10.

### **Hepatitis**

Alter MJ, Mares A, Hadler SC, Maynard JE. The effect of underreporting on the apparent incidence and epidemiology of acute viral hepatitis. Am J Epidemiol 1987;125:133–9.

CDC. Hepatitis surveillance report no. 55. Atlanta: US Department of Health and Human Services, Public Health Service, 1994.

### **Hepatitis A**

Shapiro CN, Coleman PJ, McQuillan GM, et al. Epidemiology of Hepatitis A: seroepidemiology and risk groups in the U.S.A. Vaccine 1992;10(suppl 1):S59–62.

### **Hepatitis B**

Margolis HS, Alter MJ, Hadler SC. Hepatitis B: evolving epidemiology and implications for control. Semin Liver Dis 1991;11:84–92.

### **Hepatitis Non-A, Non-B**

Alter MJ, Hadler SC, Judson FN, et al. Risk factors for acute non-A, non-B hepatitis in the United States and association with hepatitis C virus infection. JAMA 1990;264:2231–5.

Alter MJ, Margolis HS, Krawczynski K, et al. The natural history of community-acquired hepatitis C in the United States. New Engl J Med 1992;327:1899–1905.

### **Legionellosis**

CDC. Legionnaires' disease associated with cooling towers—Massachusetts, Michigan, and Rhode Island, 1993. MMWR 1994;43(27):491–9.

Edelstein PH. Legionnaires' disease. Clin Infect Dis 1993;16:741–9.

Hoge CW, Breiman RF. Advances in the epidemiology and control of *Legionella* infections. Epidemiol Rev 1991;13:329–40.

World Health Organization. Epidemiology, prevention and control of legionellosis: memorandum from a WHO meeting. Bull World Health Organ 1990;68:155–64.

### **Leptospirosis**

Anderson DC, Folland DS, Fox MD, Patton CM, Kaufmann AF. Leptospirosis: a common-source outbreak due to leptospire of the Grippityphosa serogroup. Am J Epidemiol 1978;107:538–44.

Faine S. Guidelines for the control of leptospirosis. Geneva, Switzerland: World Health Organization, 1982.

Takafuji ET, Kirkpatrick JW, Miller RN, et al. An efficacy trial of doxycycline chemoprophylaxis against leptospirosis. N Engl J Med 1984;310:497–500.

### **Lyme Disease**

CDC. Lyme disease surveillance—United States, 1993. MMWR 1994;43(31):564–5,571–2.

Kalish R. Lyme disease. Rheum Dis Clin North Am 1993;19:399–426.

Steere AC. Lyme disease. N Engl J Med 1989;321:586–96.



### **Lymphogranuloma venereum**

Perine PL, Osoba AO. Lymphogranuloma venereum. In: Holmes KK, Mardh PA, Sparling PF, et al. eds. Sexually transmitted disease. New York: McGraw-Hill, 1990:195–204.

### **Malaria**

CDC. Health information for international travel 1994. Atlanta: CDC, 1994: HHS publication no. (CDC) 94-8280.

Lobel HO, Miani M, Eng T, Bernard KW, Hightower AW, Campbell CC. Long-term malaria prophylaxis with weekly mefloquine. *Lancet* 1993;341:848–51.

Zucker JR, Campbell CC. Malaria: principles of prevention and treatment. *Infect Dis Clin North Am* 1993;7:547–67.

### **Measles**

CDC. Measles prevention: recommendations of the Immunization Practices Advisory Committee. *MMWR* 1989;38(No. SS-9):1–18.

CDC. Measles—United States, first 26 weeks, 1993. *MMWR* 1993;42:813–6.

CDC. Absence of reported measles—United States, November 1993. *MMWR* 1993;42:925–6.

### **Meningococcal Disease**

CDC. Meningococcal vaccines. *MMWR* 1985;34(18):255–9.

CDC. Laboratory-based surveillance for meningococcal disease in selected areas—United States, 1989–1991. *MMWR* 1993;42(No. SS-2):21–30.

Schwartz B, Moore PS, Broome CV. Global epidemiology of meningococcal disease. *Clin Microbiol Rev* 1989;2(suppl):S118–24.

Wenger JD, Jackson LA, Raj P, Tonelli MJ. Issues in the control of outbreaks of group C meningococcal disease in the U.S. *Infect Dis Clin Prac* 1994;3:136–40.

### **Mumps**

Briss PA, Fehrs LJ, Parker RA, et al. Sustained transmission of mumps in a highly vaccinated population: assessment of primary vaccine failure and waning vaccine-induced immunity. *J Infect Dis* 1994;169:77–82.

CDC. Mumps prevention. *MMWR* 1989;38(22):388–92,397–400.

CDC. Mumps—United States, 1985–1988. *MMWR* 1989;38(7):101–5.

Hersch BS, Fine PEM, Kent WK, et al. Mumps outbreak in a highly vaccinated population. *J Pediatr* 1991;119:187–93.

### **Murine Typhus Fever**

Benenson AS. Control of communicable diseases in man. 15th ed. Washington, DC: American Public Health Association, 1990.

McDade JE, Fishbein DB. Rickettsiaceae: the rickettsiae. In: Laboratory diagnosis of infectious diseases: principles and practice. Vol II. Viral, rickettsial, and chlamydial diseases. New York: Springer-Verlag, 1988:864–89.

Woodward TE. Murine typhus fever: its clinical and biologic similarity of epidemic typhus. In: Biology of rickettsial disease. Vol II. Boca Raton, Florida: CRC Press, 1988:79–92.

### **Pertussis**

CDC. Pertussis surveillance—United States, 1989–1991. *MMWR* 1992;41(No. SS-8):11–9.

Cherry JD, Brunell PA, Golden GS, et al. Report of the Task Force on Pertussis and Pertussis Immunization, 1988. *Pediatrics* 1988;81(suppl):S939–84.

Strebel PM, Cochi SL, Farizo KM, et al. Pertussis in Missouri: evaluation of nasopharyngeal culture, direct fluorescent antibody testing and clinical case definitions in the diagnosis of pertussis. *Clin Infect Dis* 1993;16:276–85.

### **Plague**

Craven RB, Barnes AM. Plague and tularemia. *Infect Dis Clin North Am* 1991;5:165–75.

Poland JD. Plague. In: Hoeprich PD, ed. Infectious Diseases. 2nd ed. New York: Harper and Row, 1989:151;1296–1306.

### **Poliomyelitis**

CDC. Poliomyelitis—United States, 1975–1984. *MMWR* 1986;35(11):180–2.

- Prevots DR, Suter RW, Strebel PM, Weibel RE, Cochi SL. Completeness of reporting for paralytic poliomyelitis, United States, 1980 through 1991. *Archives of Pediatrics and Adolescent Medicine* 1994;148:479–85.
- Strebel PM, Sutter RW, Cochi SL, et al. Epidemiology of poliomyelitis in the United States: one decade after the last reported case of indigenous wild virus-associated disease. *Clin Infect Dis* 1992;14:568–79.

#### **Psittacosis**

- Anonymous. Compendium of chlamydiosis (psitticosis) control, 1994. National Association of State Public Health Veterinarians, Inc. *J Am Vet Med Assoc* 1993;203:1673–80.
- Barnes RC. Laboratory diagnosis of human chlamydial infections. *Clin Microbiol Rev* 1989; 2:119–36.
- Kuritsky JN, Schmid GP, Potter ME, Anderson DC, Kaufmann AF. Psittacosis: a diagnostic challenge. *J Occup Med* 1984;26:731–3.

#### **Rabies**

- Baer G. Natural history of rabies. 2nd ed. Boca Raton, Florida: CRC press, 1991.
- CDC. Compendium of animal rabies control, 1994. *MMWR* 1994;43(No. RR-10).
- CDC. Rabies prevention—United States, 1991: recommendations of the Immunization Practices Advisory Committee (ACIP). *MMWR* 1991;40(No. RR-3).
- Krebs JW, Strine TW, Childs JE. Rabies surveillance in the United States during 1992. *J Am Vet Med Assoc* 1993;203:1718–31.

#### **Rheumatic Fever**

- Dajani AS. Prevention of rheumatic fever: a statement for health professionals by the Committee on Rheumatic Fever, Endocarditis, and Kawasaki Disease of the Council on Cardiovascular Disease in the Young, the American Heart Association. *Pediatr Infect Dis J* 1989;8:263–6.
- Kaplan EL, Johnson DR, Cleary PP. Group A streptococcal serotypes isolated from patients and sibling contacts during the resurgence of rheumatic fever in the United States in the mid-1980's. *J Infect Dis* 1989;1259:101–3.
- Veasy GL, Wiedmeier SE, Orsmond GS, et al. Resurgence of acute rheumatic fever in the intermountain area in the United States. *N Engl J Med* 1987;316:421–8.

#### **Rocky Mountain Spotted Fever (RMSF)**

- Burgdorfer W, Anacker FL, eds. *Rickettsiae and Rickettsial Diseases*. New York: Academic Press, 1981.
- McDade JE, Fishbein DB. Rickettsiaceae: the rickettsiae. In: *Laboratory diagnosis of infectious diseases: principles and practice*. Vol II. Viral, rickettsial, and chlamydial diseases. New York: Springer-Verlag, 1988:864–89.

#### **Rubella**

- CDC. Increase in rubella and congenital rubella syndrome—United States, 1988–1990. *MMWR* 1991;40(6):93–9.
- CDC. Rubella and congenital rubella syndrome—United States, January 1, 1991–May 7, 1994. *MMWR* 1994;43(21):391,397–401.
- CDC. Outbreaks of rubella among the Amish—United States, 1991. *MMWR* 1991;40(16):264.
- CDC. Congenital rubella syndrome among the Amish—Pennsylvania, 1991–1992. *MMWR* 1992;41(26):468–9,475–6.
- CDC. Rubella prevention: recommendations of the Immunization Practices Advisory Committee (ACIP). *MMWR* 1990;39(No. RR-15).
- Lindegren ML, Fehrs LJ, Hadler SC, Hinman AR. Update: rubella and congenital rubella syndrome, 1980–1990. *Epidemiol Rev* 1991;13:341–8.

#### **Salmonellosis**

- Lee LA, Puhr ND, Maloney EK, Bean NH, Tauxe RV. Increase in antimicrobial-resistant *Salmonella* infections in the United States, 1989–1990. *J Infect Dis* 1994;170:128–34.
- Mishu B, Griffin PM, Tauxe RV, Cameron DN, Hutcheson RH, Schaffner W. *Salmonella enteritidis* gastroenteritis transmitted by intact chicken eggs. *Ann Intern Med* 1991;115:190–4.
- St. Louis ME, Morse DL, Potter ME, et al. The emergence of grade A eggs as a major source of *Salmonella enteritidis* infections: new implications for the control of salmonellosis. *JAMA* 1988;259:2103–7.

Tauxe RV. *Salmonella*: a postmodern pathogen. *Journal of Food Protection* 1991;54:563–8.

#### **Shigellosis**

Lee LA, Shapiro CN, Hargrett-Bean N, Tauxe RV. Hyperendemic shigellosis in the United States: a review of surveillance data for 1967–1988. *J Infect Dis* 1991;164:894–900.

Parsonnet J, Greene KD, Gerber AR, et al. *Shigella dysenteriae* type 1 infections in U.S. travellers to Mexico. *Lancet* 1989;543–5.

Ries AA, Wells JG, Olivola D, et al. Epidemic *Shigella dysenteriae* type 1 in Burundi: panresistance and implications for prevention. *J Infect Dis* 1994;169:1035–41.

#### **Syphilis**

Aral SO, Wasserheit JN, Green SB, Judson FN, Sparling FP. The NIAID/NIH working group on integrated behavioral research for prevention and control of STDs. Part III: Issues in evaluating behavioral interventions. *Sex Transm Dis* 1990;17:208–10.

CDC. Primary and secondary syphilis—United States, 1981–1990. *MMWR* 1991;40(19):314–5, 321–3.

CDC. Regional and temporal trends in the surveillance of syphilis, United States, 1986–1990. *MMWR* 1991;40(No. SS-3):29–33.

CDC. Special focus: surveillance for sexually transmitted diseases. *MMWR* 1993;42(No. SS-3):13–9.

#### **Tetanus**

CDC. Surveillance of tetanus—United States, 1989–1990. *MMWR* 1992;41(No. SS-8):1–9.

Sutter RW, Cochi SL, Brink EW, Sirotkin BI. Assessment of vital statistics and surveillance data for monitoring tetanus mortality, United States, 1979–1984. *Am J Epidemiol* 1990;131:132–42.

#### **Toxic Shock Syndrome**

CDC. Reduced incidence of menstrual toxic shock syndrome—United States, 1980–1990. *MMWR* 1990;39(25):421–3.

Gaventa S, Reingold AL, Hightower AW, et al. Active surveillance for toxic shock syndrome in the United States, 1986. *Rev Infect Dis* 1989;(suppl):S28–34.

Schuchat A, Broome CV. Toxic shock syndrome and tampons. *Epidemiol Rev* 1991;13:99–112.

#### **Trichinosis**

Bailey TM, Schantz PM. Trends in the incidence and transmission patterns of human trichinosis in the United States, 1982–1986. *Rev Infect Dis* 1990;12:5–11.

Campbell WC (ed.). *Trichinella* and trichinosis. New York: Plenum Press, 1983.

CDC. Trichinosis surveillance—United States, 1987–1990. *MMWR* 1991;40(No. SS-3):35–42.

McAuley JB, Michelson MK, Hightower AW, Engeran S, Wintermeyer LA, Schantz PM. A trichinosis outbreak among Southeast Asian refugees. *Am J Epidemiol* 1992;135:1404–10.

#### **Tuberculosis**

American Thoracic Society/CDC. Treatment of tuberculosis and tuberculosis infection in adults and children. *Am J Respir Crit Care Med* 1994;149:1359–74.

CDC. Recommendations for counting reported tuberculosis cases. Atlanta: US Department of Health and Human Services, Public Health Service, 1977.

CDC. Expanded tuberculosis surveillance and tuberculosis morbidity—United States, 1993. *MMWR* 1994;43(20):361–6.

#### **Tularemia**

Evans ME, Gregory DW, Schaffner W, McGee ZA. Tularemia: a 30 year experience with 88 cases. *Medicine* 1985;64:251–69.

Taylor JP, Istre GR, McChesny TC, Satalowich FT, Parker RE, McFarland LM. Epidemiologic characteristics of human tularemia in the southwest-central states, 1981–1987. *Am J Epidemiol* 1991;133:1032–8.

#### **Typhoid Fever**

CDC. Typhoid immunization: recommendations of the Immunization Practices Advisory Committee. *MMWR* 1990;39(No. RR-10).

Ryan CA, Hargrett-Bean NT, Blake PA. *Salmonella typhi* infections in the United States, 1975–1984: increasing role of foreign travel. *Rev Infect Dis* 1989;11:1–7.

Woodruff BA, Pavia AT, Blake PA. A new look at typhoid vaccination: information for the practicing physician. *JAMA* 1991;265(6):756–9.

### **Varicella**

- CDC. Varicella outbreak in a women's prison—Kentucky. *MMWR* 1989;38(37):635–6,641–2.
- Gershon AA, LaRussa P, Hardy I, Steinberg S, Silverstein S. Varicella vaccine: the American experience. *J Infect Dis* 1992;166(suppl 1):S63–8.
- Lieu TA, Cochi SL, Black SB, et al. Cost-effectiveness of a routine varicella vaccination program for U.S. children. *JAMA* 1994;271:375–81.

# **PART 1:**

## **Summaries of Notifiable Diseases in the United States**

**NOTIFIABLE DISEASES — summary of reported cases, by month, United States, 1993**

Disease	Total	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Unk.
AIDS*	103,691	7,153	7,222	21,244	6,725	10,081	8,254	7,597	8,524	9,176	5,072	5,797	6,846	-
Amebiasis	2,970	144	198	202	211	218	228	284	313	272	310	245	345	-
Anthrax	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Aseptic meningitis	12,848	519	532	502	478	698	739	1,569	1,891	1,797	1,909	1,005	1,209	-
Botulism, total	97	6	5	4	5	6	6	11	11	8	16	10	9	-
Brucellosis	120	4	6	6	7	8	6	12	8	3	11	7	42	-
Chancroid†	1,399	-	401	-	-	469	-	-	229	-	-	300	-	-
Cholera	18	5	3	1	3	2	-	2	-	1	1	-	-	-
Diphtheria	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Encephalitis, primary infections	919	43	49	45	33	53	45	84	93	158	152	72	92	-
Post-infectious	170	7	20	16	14	17	17	13	15	9	10	14	18	-
Gonorrhea†	439,673	-	103,178	-	-	102,890	-	-	120,498	-	-	113,107	-	-
Granuloma inguinale†	19	-	15	-	-	4	-	-	-	-	-	-	-	-
<i>Haemophilus influenzae</i>	1,419	99	104	149	97	141	100	105	68	78	126	155	197	-
Hansen disease (leprosy)	187	11	9	20	17	27	10	8	18	14	30	11	12	-
Hepatitis A	24,238	1,739	1,718	1,985	1,739	2,104	1,678	2,132	1,661	1,838	2,535	1,916	3,193	-
Hepatitis B	13,361	772	918	1,000	1,038	1,311	999	1,191	994	1,003	1,207	1,036	1,892	-
Hepatitis, non-A, non-B <sup>§</sup>	4,786	272	354	337	293	383	301	347	340	364	496	384	915	-
Hepatitis, unspecified	627	44	47	52	43	77	34	61	40	55	49	39	86	-
Legionellosis	1,280	110	81	88	77	112	86	104	107	104	155	88	168	-
Leptospirosis	51	1	8	1	1	3	2	4	7	5	3	8	8	-
Lyme disease	8,257	175	310	323	234	433	524	1,474	1,156	845	840	738	1,205	-
Lymphogranuloma venereum†	285	-	69	-	-	73	-	-	71	-	-	72	-	-
Malaria	1,411	51	69	74	118	77	98	144	154	124	200	110	192	-
Measles (rubeola)	312	17	41	27	11	28	72	28	24	17	26	9	12	-
Meningococcal infections	2,637	155	211	311	251	273	180	204	109	121	202	160	460	-
Mumps	1,692	101	132	169	144	186	191	120	80	97	138	126	208	-
Murine typhus fever	25	-	-	1	-	3	-	5	-	4	4	4	4	-
Pertussis (whooping cough)	6,586	214	236	202	239	298	329	777	876	988	1,061	469	897	-
Plague	10	-	-	1	-	2	-	1	4	-	2	-	-	-
Poliomyelitis, paralytic <sup>¶</sup>	3	-	-	-	-	-	-	1	1	-	-	-	1	-
Psittacosis	60	3	7	4	3	8	6	5	7	4	1	7	5	-
Rabies, animal	9,377	408	512	649	801	994	724	876	930	899	1,001	713	870	-
Rabies, human	3	-	-	-	-	-	-	-	1	-	-	-	2	-
Rheumatic fever, acute	112	5	16	12	4	10	28	13	7	1	3	4	9	-
Rocky Mountain spotted fever	456	9	3	3	3	16	38	90	95	69	74	24	32	-
Rubella (German measles)	192	10	8	25	15	30	20	30	19	7	5	6	17	-
Rubella, congenital syndrome	5	1	1	-	2	-	-	1	-	-	-	-	-	-
Salmonellosis	41,641	1,909	2,099	2,196	2,188	3,131	3,256	4,819	5,119	4,367	4,980	3,164	4,413	-
Shigellosis	32,198	1,224	1,554	1,507	1,572	2,511	2,799	3,494	3,665	3,060	4,005	2,725	4,082	-
Syphilis, total all stages†	101,259	-	25,621	-	-	26,942	-	-	24,692	-	-	24,004	-	-
Primary and secondary†	26,498	-	6,952	-	-	6,684	-	-	6,621	-	-	6,241	-	-
Congenital <1 year†	3,211	-	678	-	-	834	-	-	864	-	-	835	-	-
Tetanus	48	1	1	1	3	5	5	4	7	5	4	3	9	-
Toxic-shock syndrome	212	7	21	26	23	16	15	19	25	23	15	6	16	-
Trichinosis	16	1	3	3	-	-	1	-	2	-	2	1	3	-
Tuberculosis	25,313	778	1,322	1,881	2,105	1,979	2,371	2,003	2,009	1,938	2,028	1,769	5,130	-
Tularemia	132	3	5	4	3	13	20	24	21	12	12	2	13	-
Typhoid fever	440	26	24	26	25	40	26	46	38	46	62	30	51	-
Varicella (chickenpox)	134,722	12,815	15,322	18,553	17,373	23,933	13,866	5,129	3,093	2,534	4,038	6,536	11,530	-

\* AIDS total updated through December 31, 1993.

† Cases updated through February 28, 1994.

<sup>§</sup>The number of reported cases of non-A, non-B hepatitis is misleading because in some states, reported cases included persons positive for antibody to hepatitis C virus (anti-HCV) identified in routine screening programs but who did not have acute hepatitis.

<sup>¶</sup> Subject to change due to retrospective case evaluations or late reports.

SUMMARY TABLES — 1993

NOTIFIABLE DISEASES — reported cases, by geographic division and area,  
United States, 1993

Area	Total resident population (in thousands)	AIDS	Amebiasis	Anthrax	Aseptic meningitis	Botulism			Brucellosis
						Foodborne	Infant	Other	
<b>United States</b>	<b>257,908</b>	<b>103,691*</b>	<b>2,970</b>	-	<b>12,848</b>	<b>27</b>	<b>65</b>	<b>5<sup>†</sup></b>	<b>120</b>
<b>New England</b>	<b>13,230</b>	<b>5,156</b>	<b>112</b>	-	<b>425</b>	-	-	-	<b>1</b>
Maine	1,239	149	11	-	41	-	-	-	-
N.H.	1,125	124	5	-	55	-	-	-	-
Vt.	576	74	7	-	45	-	-	-	-
Mass.	6,012	2,703	87	-	177	-	-	-	<b>1</b>
R.I.	1,000	348	2	-	107	-	-	-	-
Conn.	3,277	1,758	NN	-	NN	-	-	-	-
<b>Mid. Atlantic</b>	<b>38,125</b>	<b>26,115</b>	<b>619</b>	-	<b>1,133</b>	-	<b>8</b>	<b>1</b>	<b>3</b>
N.Y. (excl. NYC)	10,921	3,565	116	-	575	-	-	-	<b>1</b>
N.Y.C.	7,276	13,902	457	-	257	-	2	1	-
N.J.	7,879	5,434	16	-	NN	-	3	-	<b>1</b>
Pa.	12,048	3,214	30	-	301	-	3	-	<b>1</b>
<b>E.N. Central</b>	<b>43,017</b>	<b>8,069</b>	<b>208</b>	-	<b>2,234</b>	-	<b>6</b>	<b>1</b>	<b>10</b>
Ohio	11,091	1,585	18	-	720	-	5	1	<b>1</b>
Ind.	5,713	954	25	-	244	-	1	-	-
Ill.	11,697	2,959	50	-	562	-	-	-	<b>6</b>
Mich.	9,478	1,840	48	-	643	-	-	-	<b>1</b>
Wis.	5,038	731	67	-	65	-	-	-	<b>2</b>
<b>W.N. Central</b>	<b>18,054</b>	<b>3,181</b>	<b>173</b>	-	<b>817</b>	-	-	-	<b>3</b>
Minn.	4,517	659	58	-	118	-	-	-	-
Iowa	2,814	202	31	-	159	-	-	-	<b>2</b>
Mo.	5,234	1,745	54	-	275	-	-	-	-
N. Dak.	635	11	3	-	20	-	-	-	-
S. Dak.	715	29	1	-	22	-	-	-	<b>1</b>
Nebr.	1,607	179	4	-	21	-	-	-	-
Kans.	2,531	356	22	-	202	-	-	-	-
<b>S. Atlantic</b>	<b>45,738</b>	<b>22,783</b>	<b>198</b>	-	<b>2,645</b>	<b>6</b>	<b>2</b>	-	<b>36</b>
Del.	700	375	2	-	70	-	1	-	-
Md.	4,965	2,528	9	-	220	-	-	-	-
D.C.	578	1,585	-	-	37	-	-	-	-
Va.	6,491	1,625	34	-	343	-	-	-	-
W. Va.	1,820	106	2	-	56	-	-	-	-
N.C.	6,945	1,368	11	-	273	-	-	-	<b>27</b>
S.C.	3,643	1,476	NN	-	35	-	-	-	-
Ga.	6,917	2,789	90	-	173	5	1	-	<b>5</b>
Fla.	13,679	10,931	50	-	1,438	1	-	-	<b>4</b>
<b>E.S. Central</b>	<b>15,717</b>	<b>2,720</b>	<b>7</b>	-	<b>774</b>	<b>2</b>	<b>4</b>	-	<b>1</b>
Ky.	3,789	323	1	-	335	1	1	-	-
Tenn.	5,099	1,203	NN	-	166	1	-	-	-
Ala.	4,187	733	1	-	192	-	-	-	<b>1</b>
Miss.	2,643	461	5	-	81	-	3	-	-
<b>W.S. Central</b>	<b>27,983</b>	<b>10,136</b>	<b>102</b>	-	<b>1,500</b>	-	<b>3</b>	-	<b>40</b>
Ark.	2,424	404	4	-	79	-	1	-	<b>3</b>
La.	4,295	1,464	5	-	91	-	-	-	<b>2</b>
Okla.	3,231	725	7	-	1	-	-	-	<b>1</b>
Tex.	18,031	7,543	86	-	1,329	-	2	-	<b>34</b>
<b>Mountain</b>	<b>14,776</b>	<b>3,913</b>	<b>177</b>	-	<b>732</b>	<b>3</b>	<b>4</b>	-	<b>6</b>
Mont.	839	32	-	-	1	-	-	-	-
Idaho	1,099	77	7	-	11	-	2	-	-
Wyo.	470	46	3	-	8	1	-	-	-
Colo.	3,566	1,324	66	-	240	1	-	-	<b>3</b>
N. Mex.	1,616	294	21	-	115	-	-	-	<b>1</b>
Ariz.	3,936	1,238	70	-	202	-	1	-	<b>2</b>
Utah	1,860	264	4	-	74	1	1	-	-
Nev.	1,389	638	6	-	81	-	-	-	-
<b>Pacific</b>	<b>41,269</b>	<b>21,460</b>	<b>1,374</b>	-	<b>2,588</b>	<b>16</b>	<b>38</b>	<b>3</b>	<b>20</b>
Wash.	5,255	1,564	49	-	NN	4	5	-	-
Oreg.	3,032	778	95	-	NN	-	2	-	-
Calif.	31,211	18,689	1,193	-	2,428	3	29	3	<b>19</b>
Alaska	599	70	6	-	23	9	-	-	-
Hawaii	1,172	359	31	-	137	-	2	-	<b>1</b>
Guam	133	2	-	-	7	-	-	-	-
P.R.	3,522	3,199	-	-	72	-	-	-	-
V.I.	102	57	-	-	-	-	-	-	-
C.N.M.I.	43	-	-	-	2	-	-	-	-
American Samoa	47	-	-	-	-	-	-	-	-

\*Total reported through December 31, 1993. Total includes 158 cases with unknown state of residence.

†Includes wound and unspecified botulism.

**NOTIFIABLE DISEASES — reported cases, by geographic division and area,  
United States, 1993 (continued)**

Area	Encephalitis					Gonor- rhea	Granuloma inguinale	<i>Haemophilus influenzae</i>	Hansen disease (leprosy)
	Chancroid	Cholera	Diphtheria	Primary infections	Post- infectious				
<b>UNITED STATES</b>	<b>1,399*</b>	<b>18</b>	<b>-</b>	<b>919</b>	<b>170</b>	<b>439,673*</b>	<b>19*</b>	<b>1,419</b>	<b>187</b>
<b>New England</b>	<b>5</b>	<b>2</b>	<b>-</b>	<b>26</b>	<b>8</b>	<b>8,391</b>	<b>-</b>	<b>59</b>	<b>7</b>
Maine	-	-	-	3	-	80	-	4	-
N.H.	3	-	-	-	2	83	-	6	-
Vt.	-	-	-	6	-	25	-	3	-
Mass.	2	-	-	12	4	3,118	-	35	7
R.I.	-	1	-	5	2	427	-	2	-
Conn.	-	1	-	-	-	4,658	-	9	-
<b>Mid. Atlantic</b>	<b>618</b>	<b>-</b>	<b>-</b>	<b>63</b>	<b>11</b>	<b>54,796</b>	<b>-</b>	<b>135</b>	<b>18</b>
N.Y. (excl. NYC)	5	-	-	45	6	10,887	-	43	1
N.Y.C.	613	-	-	3	-	19,240	-	34	14
N.J.	-	-	-	-	-	6,444	-	27	3
Pa.	-	-	-	15	5	18,225	-	31	-
<b>E.N. Central</b>	<b>115</b>	<b>3</b>	<b>-</b>	<b>226</b>	<b>22</b>	<b>80,638</b>	<b>1</b>	<b>221</b>	<b>4</b>
Ohio	21	1	-	72	4	22,286	1	125	1
Ind.	3	-	-	20	5	8,656	-	12	-
Ill.	91	2	-	62	3	28,412	-	57	2
Mich.	-	-	-	54	10	18,014	-	18	-
Wis.	-	-	-	18	-	3,270	-	9	1
<b>W.N. Central</b>	<b>3</b>	<b>-</b>	<b>-</b>	<b>56</b>	<b>17</b>	<b>22,415</b>	<b>1</b>	<b>180</b>	<b>1</b>
Minn.	1	-	-	18	-	2,543	-	21	-
Iowa	-	-	-	7	2	1,915	1	8	-
Mo.	1	-	-	11	15	13,148	-	135	-
N. Dak.	-	-	-	5	-	54	-	-	-
S. Dak.	-	-	-	7	-	270	-	2	-
Nebr.	-	-	-	1	-	714	-	9	1
Kans.	1	-	-	7	-	3,771	-	5	-
<b>S. Atlantic</b>	<b>86</b>	<b>2</b>	<b>-</b>	<b>231</b>	<b>78</b>	<b>124,475</b>	<b>13</b>	<b>244</b>	<b>13</b>
Del.	1	-	-	3	-	1,586	-	-	-
Md.	-	-	-	25	-	13,548	-	52	1
D.C.	1	-	-	-	1	6,162	-	1	-
Va.	3	1	-	44	7	12,022	-	28	-
W. Va.	1	-	-	117	-	635	-	10	-
N.C.	13	-	-	31	-	24,187	1	23	1
S.C.	-	-	-	-	-	10,953	-	40	-
Ga.	21	-	-	2	-	31,483	12	71	-
Fla.	46	1	-	9	70	23,899	-	19	11
<b>E.S. Central</b>	<b>35</b>	<b>-</b>	<b>-</b>	<b>37</b>	<b>7</b>	<b>45,173</b>	<b>-</b>	<b>56</b>	<b>2</b>
Ky.	4	-	-	21	6	4,627	-	16	-
Tenn.	8	-	-	10	-	14,285	-	12	-
Ala.	23	-	-	3	-	15,793	-	18	2
Miss.	-	-	-	3	1	10,468	-	10	-
<b>W.S. CENTRAL</b>	<b>517</b>	<b>2</b>	<b>-</b>	<b>80</b>	<b>2</b>	<b>55,795</b>	<b>1</b>	<b>109</b>	<b>33</b>
Ark.	170	-	-	3	-	7,590	-	8	-
La.	310	-	-	10	-	13,323	1	4	-
Okla.	-	-	-	8	-	4,759	-	46	2
Tex.	37	2	-	59	2	30,123	-	51	31
<b>Mountain</b>	<b>8</b>	<b>3</b>	<b>-</b>	<b>21</b>	<b>5</b>	<b>11,549</b>	<b>1</b>	<b>125</b>	<b>6</b>
Mont.	-	-	-	-	1	81	-	5	-
Idaho	-	-	-	-	-	171	-	3	1
Wyo.	1	-	-	-	-	85	-	5	-
Colo.	-	1	-	14	-	3,803	-	24	-
N. Mex.	-	-	-	4	2	1,014	-	9	-
Ariz.	3	-	-	-	-	4,176	1	31	3
Utah	4	2	-	1	1	350	-	12	1
Nev.	-	-	-	2	1	1,869	-	36	1
<b>PACIFIC</b>	<b>12</b>	<b>6</b>	<b>-</b>	<b>179</b>	<b>20</b>	<b>36,441</b>	<b>2</b>	<b>290</b>	<b>103</b>
Wash.	-	-	-	1	-	3,740	-	19	9
Oreg.	-	-	-	-	-	1,189	-	11	-
Calif.	12	6	-	167	20	29,970	2	240	75
Alaska	-	-	-	10	-	678	-	6	1
Hawaii	-	-	-	1	-	864	-	14	18
Guam	-	-	-	-	-	83	-	-	4
P.R.	25	-	-	-	-	527	1	3	-
V.I.	5	-	-	-	-	84	-	-	-
C.N.M.I.	-	-	-	1	-	-	-	24	2
American Samoa	-	-	-	-	-	-	-	-	-

\*Cases updated through February 28, 1994.



SUMMARY TABLES — 1993

NOTIFIABLE DISEASES — reported cases, by geographic division and area, United States, 1993 (continued)

Area	Hepatitis A	Hepatitis B	Hepatitis non-A, non-B	Hepatitis unsp.	Legionel- losis	Lepto- spirosis	Lyme disease	Lympho- granuloma venereum	Malaria
<b>United States</b>	<b>24,238</b>	<b>13,361</b>	<b>4,786*</b>	<b>627</b>	<b>1,280</b>	<b>51</b>	<b>8,257</b>	<b>285†</b>	<b>1,411</b>
<b>New England</b>	<b>448</b>	<b>356</b>	<b>106</b>	<b>13</b>	<b>66</b>	-	<b>1,815</b>	<b>8</b>	<b>97</b>
Maine	14	11	2	-	6	-	18	-	7
N.H.	18	27	8	-	2	-	15	1	4
Vt.	9	10	6	-	3	-	12	-	3
Mass.	213	214	77	13	32	-	148	7	46
R.I.	77	19	13	-	23	-	272	-	7
Conn.	117	75	-	-	NN	-	1,350	-	30
<b>Mid. Atlantic</b>	<b>1,665</b>	<b>1,574</b>	<b>398</b>	<b>7</b>	<b>253</b>	<b>3</b>	<b>4,689</b>	<b>184</b>	<b>325</b>
N.Y. (excl. NYC)	464	457	259	1	89	2	2,758	12	123
N.Y.C.	722	353	3	-	3	-	60	172	126
N.J.	295	407	98	-	33	1	786	-	51
Pa.	184	357	38	6	128	-	1,085	-	25
<b>E.N. Central</b>	<b>2,727</b>	<b>1,497</b>	<b>595</b>	<b>15</b>	<b>327</b>	<b>5</b>	<b>505</b>	<b>4</b>	<b>93</b>
Ohio	338	192	37	-	154	2	30	4	15
Ind.	646	249	15	1	49	-	32	-	2
Ill.	1,023	344	102	7	26	3	19	-	46
Mich.	213	393	400	7	62	-	23	-	19
Wis.	507	319	41	-	36	-	401	-	11
<b>W.N. Central</b>	<b>2,372</b>	<b>802</b>	<b>79</b>	<b>27</b>	<b>95</b>	<b>8</b>	<b>319</b>	<b>2</b>	<b>45</b>
Minn.	497	96	14	4	3	-	141	-	21
Iowa	60	36	9	4	19	3	8	-	5
Mo.	1,443	585	25	19	33	3	108	1	9
N. Dak.	80	1	3	-	2	-	2	-	1
S. Dak.	18	-	-	-	-	-	-	1	2
Nebr.	195	19	11	-	30	2	6	-	4
Kans.	79	65	17	-	8	-	54	-	3
<b>S. Atlantic</b>	<b>1,329</b>	<b>2,408</b>	<b>723</b>	<b>95</b>	<b>219</b>	<b>2</b>	<b>639</b>	<b>60</b>	<b>338</b>
Del.	12	18	1	-	4	1	143	-	2
Md.	159	265	42	3	56	-	180	2	48
D.C.	11	43	3	-	15	-	2	8	11
Va.	156	157	54	43	11	-	95	7	41
W. Va.	29	44	43	-	4	-	50	-	2
N.C.	94	315	80	-	27	1	86	16	145
S.C.	18	51	5	1	19	-	9	-	7
Ga.	145	670	451	1	36	-	44	22	20
Fla.	705	845	44	47	47	-	30	5	62
<b>E.S. Central</b>	<b>364</b>	<b>1,748</b>	<b>1,025</b>	<b>4</b>	<b>44</b>	<b>2</b>	<b>40</b>	<b>3</b>	<b>29</b>
Ky.	142	99	16	-	19	-	16	1	5
Tenn.	104	1,152	994	3	17	1	20	1	12
Ala.	58	107	5	1	2	1	4	1	7
Miss.	60	390	10	-	6	-	-	-	5
<b>W.S. Central</b>	<b>3,190</b>	<b>1,909</b>	<b>645</b>	<b>173</b>	<b>51</b>	<b>6</b>	<b>78</b>	<b>14</b>	<b>65</b>
Ark.	74	90	5	3	6	1	8	-	4
La.	105	269	178	4	10	5	3	12	7
Okla.	213	195	50	9	13	-	19	-	6
Tex.	2,798	1,355	412	157	22	-	48	2	48
<b>Mountain</b>	<b>4,164</b>	<b>686</b>	<b>366</b>	<b>79</b>	<b>81</b>	-	<b>20</b>	<b>1</b>	<b>40</b>
Mont.	77	28	3	-	7	-	-	-	2
Idaho	298	89	-	3	1	-	2	-	1
Wyo.	17	34	119	-	7	-	9	-	-
Colo.	877	81	62	44	9	-	-	-	25
N. Mex.	400	215	107	3	6	-	2	-	5
Ariz.	1,493	96	13	14	17	-	-	1	1
Utah	828	69	42	13	16	-	2	-	2
Nev.	174	74	20	2	18	-	5	-	4
<b>Pacific</b>	<b>7,979</b>	<b>2,381</b>	<b>849</b>	<b>214</b>	<b>144</b>	<b>25</b>	<b>152</b>	<b>9</b>	<b>379</b>
Wash.	926	247	219	11	12	-	9	4	41
Oreg.	532	221	55	2	1	1	8 <sup>§</sup>	1	14
Calif.	5,677	1,876	560	198	121	-	134	4	314
Alaska	775	15	12	-	-	-	-	-	3
Hawaii	69	22	3	3	10	24	1	-	7
Guam	3	6	-	11	-	-	-	-	3
P.R.	79	412	90	2	-	11	-	1	-
V.I.	1	6	-	-	-	-	-	-	-
C.N.M.I.	-	2	-	1	-	1	-	-	2
American Samoa	20	1	-	-	-	-	-	-	-

\*The number of reported cases of non-A, non-B hepatitis is misleading because in some states, reported cases included persons positive for antibody to hepatitis C virus (anti-HCV) identified in routine screening programs but who did not have hepatitis.

†Cases updated through February 28, 1994.

§Voluntarily reportable for this state.

**NOTIFIABLE DISEASES — reported cases, by geographic division and area,  
United States, 1993 (continued)**

Area	Measles		Meningo- coccal infections	Mumps	Murine typhus fever	Pertussis	Plague	Polio- myelitis, paralytic
	Indigenous	Imported						
<b>United States</b>	<b>237</b>	<b>75*</b>	<b>2,637</b>	<b>1,692</b>	<b>25</b>	<b>6,586</b>	<b>10</b>	<b>3†</b>
<b>New England</b>	<b>54</b>	<b>9</b>	<b>133</b>	<b>15</b>	-	<b>834</b>	-	-
Maine	1	-	15	-	-	20	-	-
N.H.	2	-	6	3	-	168	-	-
Vt.	30	1	7	1	-	122	-	-
Mass.	11	7	72	1	-	408	-	-
R.I.	1	1	-	2	-	14	-	-
Conn.	9	-	33	8	-	102	-	-
<b>Mid. Atlantic</b>	<b>22</b>	<b>19</b>	<b>310</b>	<b>135</b>	<b>2</b>	<b>991</b>	-	<b>1</b>
N.Y. (excl. NYC)	4	7	126	46	1	373	-	-
N.Y.C.	12	7	37	2	1	116	-	1
N.J.	6	5	50	17	-	85	-	-
Pa.	-	-	97	70	-	417	-	-
<b>E.N. Central</b>	<b>20</b>	<b>11</b>	<b>375</b>	<b>259</b>	-	<b>1,627</b>	-	-
Ohio	6	3	108	74	-	523	-	-
Ind.	1	-	51	8	-	178	-	-
Ill.	5	4	104	74	-	434	-	-
Mich.	5	1	65	80	-	116	-	-
Wis.	3	3	47	23	-	376	-	-
<b>W.N. Central</b>	<b>1</b>	<b>2</b>	<b>142</b>	<b>67</b>	-	<b>626</b>	-	-
Minn.	-	-	23	2	-	393	-	-
Iowa	-	-	28	11	-	38	-	-
Mo.	1	-	34	46	-	144	-	-
N. Dak.	-	-	3	6	-	5	-	-
S. Dak.	-	-	7	-	-	8	-	-
Nebr.	-	-	11	1	-	14	-	-
Kans.	-	2	36	1	-	24	-	-
<b>S. Atlantic</b>	<b>25</b>	<b>8</b>	<b>470</b>	<b>463</b>	<b>1</b>	<b>673</b>	-	<b>1</b>
Del.	-	-	15	1	-	11	-	-
Md.	-	4	60	82	-	133	-	1
D.C.	-	-	6	1	-	14	-	-
Va.	-	4	52	40	1	75	-	-
W. Va.	-	-	19	23	-	8	-	-
N.C.	1	-	67	231	-	199	-	-
S.C.	-	-	31	17	-	73	-	-
Ga.	-	-	94	20	-	56	-	-
Fla.	24	-	126	48	-	104	-	-
<b>E.S. Central</b>	<b>1</b>	-	<b>154</b>	<b>58</b>	<b>1</b>	<b>297</b>	-	-
Ky.	-	-	25	-	-	38	-	-
Tenn.	-	-	45	19	1	183	-	-
Ala.	1	-	53	22	-	65	-	-
Miss.	-	-	31	17	-	11	-	-
<b>W.S. Central</b>	<b>8</b>	<b>3</b>	<b>264</b>	<b>274</b>	<b>13</b>	<b>239</b>	<b>1</b>	-
Ark.	-	-	27	10	1	18	-	-
La.	1	-	46	20	-	14	-	-
Okla.	-	-	34	13	-	86	-	-
Tex.	7	3	157	231	12	121	1	-
<b>Mountain</b>	<b>4</b>	<b>3</b>	<b>211</b>	<b>81</b>	-	<b>464</b>	<b>10</b>	-
Mont.	-	-	13	-	-	11	-	-
Idaho	-	-	21	6	-	101	-	-
Wyo.	-	-	5	5	-	2	-	-
Colo.	3	-	41	18	-	187	2	-
N. Mex.	-	-	8	NN	-	43	6	-
Ariz.	-	3	86	19	-	70	-	-
Utah	-	-	24	6	-	45	1	-
Nev.	1	-	13	27	-	5	-	-
<b>Pacific</b>	<b>102</b>	<b>20</b>	<b>578</b>	<b>340</b>	<b>8</b>	<b>835</b>	-	<b>1</b>
Wash.	-	-	97	14	-	91	-	1
Oreg.	-	4	110	NN	-	106	-	-
Calif.	89	7	346	285	8	619	-	-
Alaska	2	-	15	11	-	5	-	-
Hawaii	11	9	10	30	-	14	-	-
Guam	22	3	1	13	-	-	-	-
P.R.	356	-	14	3	-	11	-	-
V.I.	-	-	-	5	-	-	-	-
C.N.M.I.	92	1	-	15	-	1	-	-
American Samoa	-	-	-	5	-	2	-	-

\*For measles only, imported includes both out-of-state and international importations.

†Ten suspected cases of paralytic poliomyelitis were reported in 1993. Three cases have been confirmed as of August 12, 1994, two of which were vaccine-associated; one was classified as imported. The number of reported cases is subject to change due to retrospective case evaluations or late reports.

SUMMARY TABLES — 1993

NOTIFIABLE DISEASES — reported cases, by geographic division and area, United States, 1993 (continued)

Area	Psittacosis	Rabies		Rheumatic fever, acute	RMSF*	Rubella		Salmonellosis	Shigellosis
		Animal	Human			Rubella	Cong. syndrome		
<b>United States</b>	<b>60</b>	<b>9,377</b>	<b>3</b>	<b>112</b>	<b>456</b>	<b>192</b>	<b>5</b>	<b>41,641</b>	<b>32,198</b>
<b>New England</b>	<b>2</b>	<b>1,695</b>	-	<b>2</b>	<b>4</b>	<b>10</b>	-	<b>3,882</b>	<b>605</b>
Maine	-	-	-	1	-	1	-	158	8
N.H.	-	148	-	NN	-	-	-	220	17
Vt.	1	45	-	-	-	-	-	421	8
Mass.	1	720	-	-	4	9	-	2,041	285
R.I.	-	2	-	-	-	-	-	231	42
Conn.	-	780	-	1	-	-	-	811	245
<b>Mid. Atlantic</b>	<b>16</b>	<b>3,585</b>	<b>1</b>	-	<b>28</b>	<b>59</b>	<b>2</b>	<b>7,005</b>	<b>2,218</b>
N.Y. (excl. NYC)	5	2,692	1	NN	7	17	1	2,390	790
N.Y.C.	-	53	-	NN	1	22	-	1,601	554
N.J.	2	458	-	-	10	15	-	1,215	347
Pa.	9	382	-	NN	10	5	1	1,799	527
<b>E.N. Central</b>	<b>8</b>	<b>110</b>	-	<b>33</b>	<b>18</b>	<b>8</b>	-	<b>5,810</b>	<b>5,111</b>
Ohio	4	6	-	5	8	1	-	1,214	1,411
Ind.	-	12	-	-	4	3	-	543	811
Ill.	1	23	-	2	4	1	-	2,232	1,722
Mich.	1	18	-	21	2	2	-	815	829
Wis.	2	51	-	5	-	1	-	1,006	338
<b>W.N. Central</b>	<b>3</b>	<b>351</b>	-	<b>26</b>	<b>34</b>	<b>1</b>	-	<b>1,924</b>	<b>1,509</b>
Minn.	-	45	-	7	1	-	-	551	240
Iowa	2	79	-	2	7	-	-	241	69
Mo.	1	35	-	-	20	1	-	529	674
N. Dak.	-	57	-	NN	-	-	-	60	7
S. Dak.	-	49	-	14	3	-	-	99	111
Nebr.	-	7	-	NN	2	-	-	145	200
Kans.	-	79	-	3	1	-	-	299	208
<b>S. Atlantic</b>	<b>8</b>	<b>2,165</b>	-	-	<b>215</b>	<b>7</b>	-	<b>8,502</b>	<b>7,049</b>
Del.	-	135	-	NN	-	-	-	185	178
Md.	2	605	-	NN	14	3	-	936	384
D.C.	-	18	-	NN	-	-	-	102	87
Va.	1	387	-	NN	14	-	-	1,055	776
W. Va.	-	90	-	-	6	-	-	109	38
N.C.	4	107	-	NN	129	-	-	1,121	2,305
S.C.	1	165	-	NN	11	-	-	738	414
Ga.	-	471	-	NN	37	-	-	1,316	474
Fla.	-	187	-	NN	4	4	-	2,940	2,393
<b>E.S. Central</b>	<b>1</b>	<b>260</b>	-	-	<b>63</b>	<b>1</b>	-	<b>1,879</b>	<b>1,849</b>
Ky.	-	20	-	NN	16	1	-	302	171
Tenn.	1	124	-	NN	32	-	-	558	938
Ala.	-	116	-	NN	4	-	-	554	375
Miss.	-	-	-	-	11	-	-	465	365
<b>W.S. Central</b>	<b>2</b>	<b>635</b>	<b>1</b>	-	<b>77</b>	<b>24</b>	-	<b>3,297</b>	<b>5,748</b>
Ark.	-	42	-	-	17	-	-	402	201
La.	-	17	-	NN	2	1	-	650	482
Okla.	1	65	-	NN	51	1	-	321	484
Tex.	1	511	1	NN	7	22	-	1,924	4,581
<b>Mountain</b>	<b>3</b>	<b>178</b>	-	<b>28</b>	<b>16</b>	<b>12</b>	-	<b>2,018</b>	<b>1,921</b>
Mont.	-	24	-	NN	2	-	-	106	56
Idaho	1	8	-	NN	-	2	-	173	46
Wyo.	-	25	-	1	10	-	-	59	22
Colo.	-	29	-	5	4	3	-	550	611
N. Mex.	1	9	-	3	-	-	-	326	411
Ariz.	-	60	-	NN	-	2	-	519	693
Utah	1	8	-	19	-	4	-	154	42
Nev.	-	15	-	NN	-	1	-	131	40
<b>Pacific</b>	<b>17</b>	<b>398</b>	<b>1</b>	<b>23</b>	<b>1</b>	<b>70</b>	<b>3</b>	<b>7,324</b>	<b>6,188</b>
Wash.	4	-	-	-	-	-	-	832	797
Oreg.	4	7	-	NN	1	-	-	349	169
Calif.	9	363	1	20	-	41	3	5,739	5,093
Alaska	-	28	-	3	-	1	-	59	30
Hawaii	-	-	-	NN	-	28	-	345	99
Guam	-	-	-	2	-	-	-	119	35
P.R.	-	43	-	-	-	-	-	734	46
V.I.	-	-	-	-	-	-	-	5	1
C.N.M.I.	-	-	-	10	-	-	-	54	73
American Samoa	-	-	-	1	-	-	-	55	17

\*Rocky Mountain spotted fever.

NOTIFIABLE DISEASES — reported cases, by geographic division and area,  
United States, 1993 (continued)

Area	Syphilis			Tetanus	Toxic-shock syndrome	Trichinosis	Tuberculosis	Tularemia	Typhoid fever	Varicella (chickenpox)
	Primary & secondary	Cong. (<1 yr.)	All stages							
<b>United States</b>	<b>26,498*</b>	<b>3,211*</b>	<b>101,259*</b>	<b>48</b>	<b>212</b>	<b>16</b>	<b>25,313</b>	<b>132</b>	<b>440</b>	<b>134,722</b>
<b>New England</b>	<b>330</b>	<b>17</b>	<b>1,713</b>	<b>2</b>	<b>13</b>	-	<b>649</b>	-	<b>31</b>	<b>12,655</b>
Maine	7	-	19	-	3	-	27	-	-	1,733
N.H.	26	-	50	-	3	-	26	-	2	1,635
Vt.	1	-	1	-	1	-	7	-	-	NN
Mass.	122	6	935	2	5	-	370	-	23	6,619
R.I.	16	1	146	-	1	-	64	-	-	2,668
Conn.	158	10	562	-	-	-	155	-	6	NN
<b>Mid. Atlantic</b>	<b>2,415</b>	<b>1,070</b>	<b>19,306</b>	<b>2</b>	<b>33</b>	<b>4</b>	<b>5,611</b>	<b>2</b>	<b>139</b>	<b>4,773</b>
N.Y. (excl. NYC)	258	94	1,980	-	17	4	718	2	20	NA
N.Y.C.	1,132	654	10,513	-	1	-	3,235	-	95	4,773
N.J.	328	161	2,556	-	-	-	912	-	18	NN
Pa.	697	161	4,257	2	15	-	746	-	6	NN
<b>E.N. Central</b>	<b>4,070</b>	<b>527</b>	<b>11,789</b>	<b>7</b>	<b>48</b>	<b>2</b>	<b>2,385</b>	<b>5</b>	<b>40</b>	<b>64,380</b>
Ohio	1,180	70	2,889	2	12	-	315	-	7	5,472
Ind.	362	1	1,019	1	1	-	248	1	2	NN
Ill.	1,489	368	4,881	1	11	2	1,242	3	23	26,447
Mich.	543	84	1,952	3	24	-	480	1	7	32,461
Wis.	496	4	1,048	-	-	-	100	-	1	NA
<b>W.N. Central</b>	<b>1,631</b>	<b>111</b>	<b>3,271</b>	<b>10</b>	<b>15</b>	-	<b>582</b>	<b>39</b>	<b>2</b>	<b>16,814</b>
Minn.	66	9	261	6	3	-	141	-	-	NN
Iowa	64	1	175	1	7	-	59	-	-	5,057
Mo.	1,354	97	2,500	1	2	-	257	17	2	9,609
N. Dak.	1	-	4	-	-	-	7	-	-	37
S. Dak.	2	-	3	-	-	-	16	17	-	420
Nebr.	14	1	35	-	-	-	22	2	-	4
Kans.	130	3	293	2	3	-	80	3	-	1,687
<b>S. Atlantic</b>	<b>6,513</b>	<b>586</b>	<b>24,237</b>	<b>8</b>	<b>24</b>	<b>6</b>	<b>4,626</b>	<b>4</b>	<b>63</b>	<b>8,332</b>
Del.	94	3	274	-	-	-	66	-	1	3
Md.	359	30	1,865	2	1	3	406	-	8	NN
D.C.	291	74	1,652	-	-	-	161	-	-	4
Va.	660	23	1,970	-	7	1	458	-	7	2,917
W. Va.	8	6	195	-	-	-	75	-	-	5,288
N.C.	1,937	54	4,448	-	4	-	594	2	3	NN
S.C.	921	82	2,339	1	-	-	401	-	-	120
Ga.	1,052	79	4,077	-	2	-	810	-	3	NN
Fla.	1,191	235	7,417	5	10	2	1,655	2	41	NN
<b>E.S. Central</b>	<b>4,117</b>	<b>158</b>	<b>10,494</b>	<b>3</b>	<b>11</b>	-	<b>1,727</b>	<b>4</b>	<b>7</b>	<b>3,981</b>
Ky.	331	9	651	-	3	-	405	1	2	1,429
Tenn.	1,156	52	3,241	2	4	-	556	2	2	2,552
Ala.	869	27	2,333	1	2	-	487	1	3	NN
Miss.	1,761	70	4,269	-	2	-	279	-	-	NN
<b>W.S. Central</b>	<b>5,969</b>	<b>409</b>	<b>19,079</b>	<b>9</b>	<b>2</b>	-	<b>3,181</b>	<b>58</b>	<b>19</b>	<b>14,292</b>
Ark.	559	8	1,600	1	-	-	209	36	2	NN
La.	2,598	144	6,854	-	-	-	367	-	1	NN
Okla.	282	11	721	1	2	-	209	17	1	-
Tex.	2,530	246	9,904	7	-	-	2,396	5	15	14,291
<b>Mountain</b>	<b>256</b>	<b>27</b>	<b>1,251</b>	<b>1</b>	<b>16</b>	<b>1</b>	<b>592</b>	<b>12</b>	<b>11</b>	<b>7,192</b>
Mont.	1	-	4	-	-	-	22	5	-	84
Idaho	2	-	15	-	2	-	12	-	-	NN
Wyo.	4	-	9	-	-	1	7	3	-	NN
Colo.	90	8	287	1	2	-	102	1	5	NN
N. Mex.	34	-	172	-	1	-	74	-	2	NN
Ariz.	95	16	557	-	2	-	231	-	3	6,811
Utah	10	-	68	-	7	-	46	2	1	297
Nev.	20	3	139	-	2	-	98	1	-	NN
<b>Pacific</b>	<b>1,197</b>	<b>306</b>	<b>10,119</b>	<b>6</b>	<b>50</b>	<b>3</b>	<b>5,960</b>	<b>8</b>	<b>128</b>	<b>2,303</b>
Wash.	67	4	360	1	7	-	286	2	8	NN
Oreg.	39	4	179	-	-	-	154	3	4	NN
Calif.	1,073	298	9,488	5	42	1	5,212	3	113	NN
Alaska	11	-	51	-	-	-	57	-	-	NN
Hawaii	7	-	41	-	1	2	251	-	3	2,303
Guam	-	-	5	-	-	-	NA	-	4	553
P.R.	470	18	2,482	2	-	-	253	-	-	8,228
V.I.	12	1	39	1	-	-	4	-	-	486
C.N.M.I.	-	-	-	-	-	-	NA	-	-	178
American Samoa	-	-	-	-	-	-	NA	-	1	176

\*Cases updated through February 28, 1994.

## NOTIFIABLE DISEASES — summary of reported cases, by age group, United States, 1993

Disease	Total	Under 1	1-4	5-9	10-14	15-19	20-24	25-29	30-39	40-49	50-59	60+	Age not stated
AIDS*	103,691	328	352	148	138	487	3,771	14,795	47,064	26,602	7,423	2,583	-
Amebiasis	2,970	13	119	147	70	82	123	200	394	234	121	98	1,369
Anthrax	-	-	-	-	-	-	-	-	-	-	-	-	-
Aseptic meningitis	12,848	2,668	917	1,297	913	803	1,091	1,276	2,068	702	267	328	518
Botulism, total	97	63	1	-	-	-	1	1	3	5	3	11	9
Brucellosis	120	1	1	12	1	7	11	11	27	18	13	12	6
Cholera	18	-	2	-	-	1	-	4	5	1	3	2	-
Diphtheria	-	-	-	-	-	-	-	-	-	-	-	-	-
Encephalitis, primary infections	919	65	78	115	99	58	58	51	115	79	50	136	15
Post-infectious	170	12	25	28	9	10	11	9	20	13	9	23	1
Gonorrhea <sup>†</sup>	440,149	-	-	-	8,249	116,974	119,739	62,217	64,110	23,563	-	-	43,660
<i>Haemophilus influenzae</i>	1,419	288	147	57	32	34	23	28	93	70	100	476	71
Hansen disease (leprosy)	187	-	1	-	3	5	18	16	30	25	18	50	21
Hepatitis A	24,238	141	1,442	3,075	2,120	1,845	2,520	2,772	4,378	1,912	904	1,237	1,892
Hepatitis B	13,361	63	78	100	201	915	1,768	2,108	3,670	1,868	800	796	994
Hepatitis, non-A, non-B <sup>§</sup>	4,786	34	16	23	28	74	245	522	1,881	1,106	290	439	128
Hepatitis, unspecified	627	4	28	49	33	26	54	74	147	85	49	51	27
Legionellosis	1,280	4	5	-	7	9	20	35	136	180	188	646	50
Leptospirosis	51	-	-	4	5	2	10	3	11	7	2	1	6
Lyme disease	8,257	13	334	621	490	325	312	399	1,161	1,169	845	1,391	1,197
Malaria	1,411	16	77	69	70	99	234	177	324	158	77	65	45
Measles (rubeola)	312	34	85	52	35	27	24	14	31	5	1	-	4
Meningococcal infections	2,637	444	480	227	199	297	173	82	127	104	93	270	141
Mumps	1,692	14	284	497	316	186	88	50	111	44	18	8	76
Murine typhus fever	25	-	-	-	1	2	-	1	5	7	2	7	-
Pertussis (whooping cough)	6,586	2,838	1,378	737	552	275	98	92	242	164	45	34	131
Plague	10	-	-	-	-	-	2	-	2	1	-	5	-
Poliomyelitis, paralytic <sup>¶</sup>	3	-	1	-	-	-	-	-	2	-	-	-	-
Psittacosis	60	-	-	-	2	4	1	2	19	16	7	6	3
Rabies, human	3	-	-	-	1	-	-	-	-	-	-	2	-
Rheumatic fever, acute	112	-	4	12	34	4	1	1	10	2	-	1	43
Rocky Mountain spotted fever	456	4	38	58	40	24	21	17	70	54	46	69	15
Rubella (German measles)	192	15	17	19	13	8	33	24	34	16	4	2	7
Salmonellosis	41,641	5,162	5,747	2,344	1,457	1,584	2,337	2,283	4,099	2,796	1,752	4,169	7,911
Shigellosis	32,198	744	8,849	5,859	1,500	863	1,501	1,690	2,457	930	455	652	6,698
Syphilis, primary and secondary <sup>†</sup>	26,546	-	-	-	174	2,940	5,469	5,176	8,040	4,686	-	-	35
Tetanus	48	-	-	-	-	1	3	3	3	4	8	26	-
Toxic-shock syndrome	212	-	6	2	15	32	27	28	55	25	11	8	3
Trichinosis	16	-	-	-	-	2	1	2	1	4	4	-	2
Tuberculosis	25,313	220	856	355	290	580	1,263	1,965	5,255	4,302	3,021	7,162	44
Tularemia	132	1	18	13	14	9	3	9	12	13	7	27	6
Typhoid fever	440	4	39	58	38	38	55	56	83	29	13	18	9
Varicella (chickenpox)	134,722	775	6,896	20,068	3,682	848	330	254	227	68	35	73	101,466

\* AIDS total updated through December 31, 1993.

<sup>†</sup> Age data are collected on aggregate forms different from those used for numbers of reported cases. Therefore, total cases reported on this table may differ slightly from other tables.

Cases among persons ≤9 years of age are not shown because some of these may not be due to sexual transmission; however, they are included in the totals.

<sup>§</sup> The number of reported cases of non-A, non-B hepatitis is misleading because in some states, reported cases included persons positive for antibody to hepatitis C virus (anti-HCV) who were identified in routine screening programs but who did not have acute hepatitis.

<sup>¶</sup> Subject to change due to retrospective case evaluations or late reports.

**NOTIFIABLE DISEASES — summary of reported cases, by race,\* United States, 1993**

Disease	Total	American Indian or Alaska Native	Asian or Pacific Islander	Black	White	Other	Race not stated <sup>†</sup>
AIDS <sup>§</sup>	103,691	339	765	38,514	48,222	-	15,851
Amebiasis	2,970	7	140	88	570	3	2,162
Anthrax	-	-	-	-	-	-	-
Aseptic meningitis	12,848	42	211	1,780	7,210	36	3,569
Botulism, total	97	9	2	8	54	-	24
Brucellosis	120	-	2	25	58	-	35
Cholera	18	-	1	1	6	-	10
Diphtheria	-	-	-	-	-	-	-
Encephalitis, primary infections	919	4	12	88	603	2	210
Post-infectious	170	-	1	21	110	2	36
Gonorrhea <sup>¶</sup>	440,149	1,708	1,340	270,829	41,346	-	124,926
<i>Haemophilus influenzae</i>	1,419	17	21	211	805	1	364
Hansen disease (leprosy)	187	-	61	7	52	2	65
Hepatitis A	24,238	1,217	461	2,336	12,528	84	7,612
Hepatitis B	13,361	76	708	2,953	5,435	48	4,141
Hepatitis, non-A, non-B**	4,786	46	40	646	2,382	1	1,671
Hepatitis, unspecified	627	4	10	64	339	9	201
Legionellosis	1,280	3	13	117	865	1	281
Leptospirosis	51	-	8	1	25	-	17
Lyme disease	8,257	13	39	121	4,855	2	3,227
Malaria	1,411	4	293	347	424	20	323
Measles (rubeola)	312	1	32	24	191	-	64
Meningococcal infections	2,637	15	37	323	1,626	19	617
Mumps	1,692	5	40	286	817	5	539
Murine typhus fever	25	-	-	3	21	-	1
Pertussis (whooping cough)	6,586	33	38	434	2,539	7	3,535
Plague	10	1	-	-	7	-	2
Poliomyelitis, paralytic <sup>††</sup>	3	-	-	1	2	-	-
Psittacosis	60	-	-	3	38	-	19
Rabies, human	3	-	-	-	3	-	-
Rheumatic fever, acute	112	13	2	-	30	1	66
Rocky Mountain spotted fever	456	24	1	18	341	-	72
Rubella (German measles)	192	1	31	4	115	-	41
Rubella, congenital syndrome	5	-	-	1	2	-	2
Salmonellosis	41,641	166	481	3,580	17,196	30	20,188
Shigellosis	32,198	341	147	5,566	12,847	16	13,281
Syphilis, primary and secondary <sup>¶</sup>	26,546	37	84	22,330	2,230	-	1,865
Tetanus	48	-	1	4	33	-	10
Toxic-shock syndrome	212	-	4	10	160	1	37
Trichinosis	16	-	1	2	9	-	4
Tuberculosis	25,313	278	3,739	9,175	11,978	-	143
Tularemia	132	18	-	2	93	-	19
Typhoid fever	440	-	117	40	78	14	191
Varicella (chickenpox)	134,722	2	94	1,615	477	-	132,534

\*Some information on race/ethnicity was reported using one variable with the following categories: American Indian or Alaska Native, Asian or Pacific Islander, Black non-Hispanic, White non-Hispanic, and Hispanic. Race not stated includes cases originally reported as Hispanic.

† Includes cases originally reported as Hispanic: 15,681 for AIDS; 16,150 for Gonorrhea; and 1,338 for Syphilis, primary and secondary.

§ AIDS total reported through December 31, 1993.

¶ Race data are collected on aggregate forms different from those used for numbers of reported cases. Therefore, total cases reported on this table may differ slightly from other tables.

\*\*The number of reported cases of non-A, non-B hepatitis is misleading because in some states, reported cases included persons positive for antibody to hepatitis C virus (anti-HCV) who were identified in routine screening programs but who did not have acute hepatitis.

†† Subject to change due to retrospective case evaluations or late reports.

**NOTIFIABLE DISEASES — summary of reported cases, by ethnicity,\* United States, 1993**

Disease	Total	Hispanic	Non-Hispanic	Ethnicity not stated
AIDS†	103,691	15,681	86,736	1,274
Amebiasis	2,970	260	518	2,192
Anthrax	-	-	-	-
Aseptic meningitis	12,848	1,545	6,096	5,207
Botulism, total	97	11	55	31
Brucellosis	120	58	15	47
Cholera	18	13	1	4
Diphtheria	-	-	-	-
Encephalitis, primary infections	919	52	409	458
Post-infectious	170	18	94	58
Gonorrhea§	440,149	16,150	312,175	111,824
<i>Haemophilus influenzae</i>	1,419	78	722	619
Hansen disease (leprosy)	187	67	90	30
Hepatitis A	24,238	4,251	11,912	8,075
Hepatitis B	13,361	928	6,525	5,908
Hepatitis, non-A, non-B¶	4,786	364	1,962	2,460
Hepatitis, unspecified	627	151	294	182
Legionellosis	1,280	21	652	607
Leptospirosis	51	-	15	36
Lyme disease	8,257	92	2,980	5,185
Malaria	1,411	105	897	409
Measles (rubeola)	312	43	201	68
Meningococcal infections	2,637	219	1,403	1,015
Mumps	1,692	271	716	705
Murine typhus fever	25	6	16	3
Pertussis (whooping cough)	6,586	422	1,740	4,424
Plague	10	2	5	3
Poliomyelitis, paralytic**	3	-	3	-
Psittacosis	60	3	29	28
Rabies, human	3	1	2	-
Rheumatic fever, acute	112	5	21	86
Rocky Mountain spotted fever	456	4	246	206
Rubella (German measles)	192	36	104	52
Rubella, congenital syndrome	5	-	3	2
Salmonellosis	41,641	2,248	13,620	25,773
Shigellosis	32,198	3,436	11,523	17,239
Syphilis, primary and secondary§	26,546	1,338	24,560	648
Tetanus	48	9	19	20
Toxic-shock syndrome	212	7	119	86
Trichinosis	16	1	6	9
Tuberculosis	25,313	5,197	19,876	240
Tularemia	132	2	47	83
Typhoid fever	440	141	191	108
Varicella (chickenpox)	134,722	1,297	2,178	131,247

\*Some information on race/ethnicity was reported using one variable with the following categories: American Indian or Alaska Native, Asian or Pacific Islander, Black non-Hispanic, White non-Hispanic, and Hispanic. Ethnicity not stated includes cases originally reported as American Indian or Alaska Native and Asian or Pacific Islander.

†AIDS total reported through December 31, 1993.

§Ethnicity data are collected on aggregate forms different from those used for numbers of reported cases. Therefore, total cases reported on this table may differ slightly from other tables.

¶The number of reported cases of non-A, non-B hepatitis is misleading because in some states, reported cases included persons positive for antibody to hepatitis C virus (anti-HCV) who were identified in routine screening programs but who did not have acute hepatitis.


\*\*Subject to change due to retrospective case evaluations or late reports.

# **PART 2:**

## **Graphs and Maps for Selected Notifiable Diseases in the United States**




**ACQUIRED IMMUNODEFICIENCY SYNDROME (AIDS) — reported cases and known deaths, by year, United States,\* 1981–1993†**




\* Includes Guam, Puerto Rico, the U.S. Pacific Islands, and the U.S. Virgin Islands (N=361,164).

† The case definition for AIDS was most recently revised as of January 1, 1993.

**ACQUIRED IMMUNODEFICIENCY SYNDROME (AIDS) — annual rates, by selected age group and sex for reported cases, United States, 1993**




ACQUIRED IMMUNODEFICIENCY SYNDROME (AIDS) — reported pediatric cases,\* United States and Puerto Rico, 1993




\*Children <13 years of age.

**ACQUIRED IMMUNODEFICIENCY SYNDROME (AIDS) — reported cases, per 100,000 population, United States and Puerto Rico, 1993\***



\*The denominator for Puerto Rico is based on extrapolations from U.S. Bureau of Census population data from 1980 and 1990 censuses.

ACQUIRED IMMUNODEFICIENCY SYNDROME (AIDS) — reported adult/adolescent cases among men, by exposure category,\* United States,† 1993




\* For surveillance purposes, AIDS cases are counted only once in a hierarchy of exposure categories. Persons with more than one reported mode of exposure to HIV are classified in the exposure category listed first in the hierarchy except for men with both a history of sexual contact with other men and injecting-drug use. These persons make up a separate exposure category.

† Includes Guam, Puerto Rico, the U.S. Pacific Islands, and the U.S. Virgin Islands.

§ Includes transfusion, tissue, and organ recipients from donors who were screened as negative for HIV antibody at the time of donation.

¶ "Other" refers to persons who developed AIDS after exposure to HIV-infected blood within the health-care setting, as documented by evidence of seroconversion or other laboratory studies. "Risk not reported or identified" refers to patients whose mode of exposure to HIV is unknown. This includes patients under investigation; patients who died, were lost to follow-up, or refused interview; and patients whose mode of exposure to HIV remains undetermined after investigation.

**ACQUIRED IMMUNODEFICIENCY SYNDROME (AIDS) — reported adult/adolescent cases among women, by exposure category,\* United States,† 1993**



\* For surveillance purposes, AIDS cases are counted only once in a hierarchy of exposure categories. Persons with more than one reported mode of exposure to HIV are classified in the exposure category listed first in the hierarchy.


† Includes Guam, Puerto Rico, the U.S. Pacific Islands, and the U.S. Virgin Islands.

§ Includes transfusion, tissue, and organ recipients from donors who were screened as negative for HIV antibody at the time of donation.


¶ "Other" refers to persons who developed AIDS after exposure to HIV-infected blood within the health-care setting, as documented by evidence of seroconversion or other laboratory studies. "Risk not reported or identified" refers to patients whose mode of exposure to HIV is unknown. This includes patients under investigation; patients who died, were lost to follow-up, or refused interview; and patients whose mode of exposure to HIV remains undetermined after investigation.

## GRAPHS AND MAPS


**ARBOVIRAL INFECTIONS (of the central nervous system) — cases due to St. Louis encephalitis virus, by month, United States, 1975–1993**




**ARBOVIRAL INFECTIONS (of the central nervous system) — cases due to California serogroup viruses, by month, United States, 1975–1993**



**ARBOVIRAL INFECTIONS (of the central nervous system) — cases due to Western equine encephalitis virus, by month, United States, 1975–1993**




**ARBOVIRAL INFECTIONS (of the central nervous system) — cases due to Eastern equine encephalitis virus, by month, United States, 1975–1993**




**GRAPHS AND MAPS**

**AMEBIASIS — by year, United States, 1955–1993**




**ASEPTIC MENINGITIS — by month, United States, 1986–1993**






**ASEPTIC MENINGITIS — reported cases, per 100,000 population, United States and territories, 1993**



## GRAPHS AND MAPS


### BOTULISM (foodborne) — by year, United States, 1975–1993




### BOTULISM (infant) — by year, United States, 1975–1993




**BRUCELLOSIS — by year, United States, 1945–1993**




CHOLERA — reported cases, United States and territories, 1993



**DIPHTHERIA — by year, United States, 1955–1993**




**ENCEPHALITIS (post-infectious) — by year, United States, 1984–1993**



## GRAPHS AND MAPS


### GONORRHEA — by sex, United States, 1960–1993



### GONORRHEA — by race and ethnicity, United States, 1981–1993




**GONORRHEA — reported cases, per 100,000 population,\* United States, 1993**




\*The total rate of gonorrhea for the United States was 172.40 per 100,000 population. The Year 2000 Objective is  $\leq 225$  per 100,000 population.

**HAEMOPHILUS INFLUENZAE — reported cases, per 100,000 population, United States and territories, 1993**







**HAEMOPHILUS INFLUENZAE — by age group, United States, 1993**



**HANSEN DISEASE (leprosy) — by year, United States, 1955-1993**




## HEPATITIS — by year, United States, 1965–1993




\*The first hepatitis B vaccine was licensed June 1982.

†Cases reported as acute non-A, non-B hepatitis may include many chronic hepatitis C virus (HCV) infections because the test for antibody to HCV (anti-HCV) does not distinguish acute infections from chronic infections.


HEPATITIS A — reported cases, per 100,000 population, United States and territories, 1993



HEPATITIS B — reported cases, per 100,000 population, United States and territories, 1993




LEGIONELLOSIS — reported cases, per 100,000 population, United States and territories, 1993




GRAPHS AND MAPS

LEPTOSPIROSIS — by year, United States, 1955-1993




LYME DISEASE — reported cases, per 100,000 population, United States and territories, 1993




## GRAPHS AND MAPS

### MALARIA — by year, United States, 1930–1993




Y-AXIS IS LOG SCALE

### MEASLES (rubeola) — by year, United States, 1950–1993







MEASLES (rubeola) — counties reporting cases, United States, 1993




MENINGOCOCCAL INFECTION — counties reporting cases, United States, 1993



**MENINGOCOCCAL INFECTION — by year, United States, 1935–1993**




**MUMPS — by year, United States, 1968–1993**




NOTE: Mumps vaccine licensed December 1967.

GRAPHS AND MAPS

MURINE TYPHUS FEVER — by year, United States, 1955-1993




**PERTUSSIS (whooping cough) — by year, United States, 1957–1993**




NOTE: DTP vaccine licensed 1948.

\* Data on mortality are not yet available for 1992 and 1993.


**PERTUSSIS (whooping cough) — by age group, United States, 1993**




PERTUSSIS (whooping cough) — reported cases, per 100,000 population, United States and territories, 1993



**PLAGUE — among humans, by year, United States, 1955–1993**




**POLIOMYELITIS (paralytic) — by year, United States, 1951–1993**




Y-AXIS IS LOG SCALE

## GRAPHS AND MAPS

### PSITTACOSIS — by year, United States, 1955–1993




### RABIES — wild and domestic animals, by year, United States and Puerto Rico, 1955–1993







RHEUMATIC FEVER — reported cases, United States and territories, 1993



ROCKY MOUNTAIN SPOTTED FEVER — counties reporting cases, United States, 1993




ROCKY MOUNTAIN SPOTTED FEVER — by year, United States, 1955–1993




## GRAPHS AND MAPS

### RUBELLA (German measles) — by year, United States, 1966–1993




### SALMONELLOSIS (excluding typhoid fever) — by year, United States, 1955–1993




**SHIGELLOSIS — by year, United States, 1955–1993**




**SYPHILIS — by year, United States, 1941–1993**






\*The total rate of primary and secondary syphilis for the United States was 10.4 per 100,000 population. The Year 2000 Objective is  $\leq 10.0$  per 100,000 population.

**SYPHILIS (primary and secondary) — counties with rates above and below the Year 2000 Objective, United States, 1993**



**GRAPHS AND MAPS**

**SYPHILIS (primary and secondary) — by race and ethnicity, United States, 1981–1993**




**SYPHILIS (primary and secondary) — by sex, United States, 1956–1993**






**CONGENITAL SYPHILIS — reported cases in infants <1 year of age and rate of primary and secondary syphilis among women — United States, 1970–1993**



\* The surveillance case definition for congenital syphilis changed in 1989.


**GRAPHS AND MAPS**

**TETANUS — by year, United States, 1955–1993**




NOTE: Tetanus toxoid was first available in 1933.

**TETANUS — by age group, United States, 1993**




**TOXIC SHOCK SYNDROME — by quarter, United States, 1979–1993\***




\* Includes only cases meeting the CDC case definition (N=3,245) for staphylococcal disease.

**TRICHINOSIS — by year, United States, 1950–1993**




## TUBERCULOSIS — rates by state, United States,\* 1993



\*Includes Puerto Rico and the U.S. Virgin Islands.


TUBERCULOSIS — percentage of cases, by race and ethnicity,\* United States, 1993




\* Excludes 300 cases (1.2%) with race and/or ethnicity unknown.

## GRAPHS AND MAPS


### TUBERCULOSIS – by year, United States, 1975–1993



### TUBERCULOSIS — frequency distribution of cases by age, race, and ethnicity, United States, 1993




**TUBERCULOSIS — cases, among foreign-born persons and persons born in the United States, 1993**




\* Includes persons born in Puerto Rico.

**TULAREMIA — by year, United States, 1955–1993**




TULAREMIA — counties reporting cases, United States, 1993






**TYPHOID FEVER — by year, United States, 1955–1993**



**VARICELLA (chickenpox) — by month, United States, 1986–1993**



# **PART 3:**

**Historical  
Summary  
Tables**



TABLE 2. NOTIFIABLE DISEASES — summary of reported cases, per 100,000 population, United States, 1984–1993

Disease	1993	1992	1991	1990	1989	1988	1987	1986	1985	1984
AIDS	40.20	17.83	17.32	16.72	13.58	12.61	8.66	5.36	3.46	1.88
Amebiasis	1.21	1.21	1.23	1.38	1.34	1.20	1.33	1.47	1.92	2.30
Anthrax	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Aseptic meningitis	5.39	5.18	6.26	4.77	4.14	2.94	4.72	4.72	4.50	3.57
Botulism, total (including wound and unsp.)	0.04	0.04	0.05	0.04	0.04	0.03	0.03	0.05	0.05	0.05
Foodborne	0.01	0.00	0.01	0.01	0.01	0.01	0.01	0.01	0.02	0.01
Infant	0.03	0.03	0.03	0.03	0.02	0.02	0.02	0.03	0.03	0.04
Brucellosis	0.05	0.04	0.04	0.03	0.04	0.04	0.05	0.04	0.06	0.06
Chancroid*	0.54	0.80	1.40	1.70	1.90	2.04	2.07	1.57	0.87	0.28
Cholera	0.00	0.04	0.01	0.00	0.00	0.00	0.00	0.01	0.00	0.00
Diphtheria	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Encephalitis, primary	0.36	0.30	0.40	0.54	0.40	0.36	0.58	0.54	0.58	0.53
Post-infectious	0.07	0.05	0.03	0.04	0.04	0.05	0.05	0.05	0.07	0.05
Gonorrhea*	172.40	201.60	249.48	276.60	297.36	298.74	323.14	376.37	384.51	374.75
Granuloma inguinale*	0.00	0.00	0.01	0.00	0.00	0.00	0.01	0.03	0.02	0.01
<i>Haemophilus influenzae</i>	0.55	0.55	1.10	.....	.....	.....	↑	.....	.....	.....
Hansen disease (leprosy)	0.07	0.07	0.06	0.08	0.07	0.07	0.10	0.11	0.15	0.12
Hepatitis A	9.40	9.06	9.67	12.64	14.43	11.60	10.39	10.02	10.03	9.33
Hepatitis B	5.18	6.32	7.14	8.48	9.43	9.43	10.65	11.17	11.50	11.06
Hepatitis, non-A, non-B <sup>§</sup>	1.86	2.36	1.42	1.03	1.02	1.07	1.23	1.55	1.81	1.64
Hepatitis, unspecified	0.24	0.35	0.50	0.67	0.93	1.00	1.27	1.69	2.38	2.34
Legionellosis	0.50	0.53	0.53	0.55	0.48	0.44	0.43	0.43	0.37	0.35
Leptospirosis	0.02	0.02	0.02	0.03	0.04	0.02	0.02	0.02	0.02	0.02
Lyme disease	3.20	0.12	3.80	.....	.....	.....	↑	.....	.....	.....
Lymphogranuloma venereum*	0.10	0.10	0.19	0.10	0.08	0.07	0.13	0.16	0.10	0.07
Malaria	0.55	0.43	0.51	0.52	0.51	0.45	0.39	0.47	0.44	0.43
Measles (rubeola)	0.12	0.88	3.82	11.17	7.33	1.38	1.50	2.61	1.18	1.10
Meningococcal infections	1.02	0.84	0.84	0.99	1.10	1.21	1.20	1.08	1.04	1.16
Mumps	0.66	1.03	1.72	2.17	2.34	2.05	5.43	3.37	1.30	1.32
Murine typhus fever	0.01	0.01	0.02	0.02	0.02	0.02	0.02	0.03	0.02	0.02
Pertussis (whooping cough)	2.55	1.60	1.08	1.84	1.67	1.40	1.16	1.74	1.50	0.96
Plague	0.00	0.01	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.01
Poliomyelitis, paralytic	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Psittacosis	0.02	0.04	0.04	0.05	0.05	0.05	0.04	0.09	0.05	0.07
Rabies, human	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Rheumatic fever, acute	0.08	0.06	0.12	0.09	0.13	0.14	0.13	0.12	0.07	0.08
Rocky Mountain spotted fever	0.18	0.20	0.25	0.26	0.25	0.25	0.25	0.32	0.30	0.36
Rubella (German measles)	0.07	0.06	0.56	0.45	0.16	0.09	0.13	0.23	0.26	0.32
Rubella, congenital syndrome <sup>¶</sup>	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Salmonellosis, excluding typhoid fever	16.15	16.04	19.10	19.54	19.26	19.91	20.92	20.73	27.37	17.30
Shigellosis	12.48	9.38	9.34	10.89	10.07	12.46	9.80	7.11	7.14	7.36
Smallpox	..... Last documented case occurred in 1949.....									
Syphilis, primary and secondary*	10.40	13.70	17.26	20.10	18.07	16.43	14.54	11.65	11.45	12.20
Total, all stages*	39.70	45.30	51.69	53.80	44.94	42.37	35.81	28.50	28.50	29.81
Tetanus	0.02	0.02	0.02	0.03	0.02	0.02	0.02	0.03	0.03	0.03
Toxic-shock syndrome	0.08	0.10	0.11	0.13	0.16	0.16	0.15	0.19	0.18	0.24
Trichinosis	0.01	0.02	0.02	0.05	0.01	0.02	0.02	0.02	0.03	0.03
Tuberculosis	9.82	10.46	10.42	10.33	9.46	9.13	9.25	9.44	9.30	9.42
Tularemia	0.05	0.06	0.08	0.06	0.06	0.08	0.09	0.07	0.07	0.12
Typhoid fever	0.17	0.16	0.20	0.22	0.19	0.18	0.16	0.15	0.17	0.17
Varicella (chickenpox)	118.54	176.54	135.82	120.06	121.77	122.43	136.68	122.42	123.23	138.44
Yellow fever	..... Last indigenous case reported 1911; last imported, 1924.....									

NOTE: Rates <0.01 after rounding are shown as 0.00.

\* Post-censal population data from 1992 were used to calculate 1993 rates.

† Not previously notifiable nationally.

§ The number of reported cases of non-A, non-B hepatitis is misleading because in some states, reported cases included persons positive for antibody to hepatitis C virus (anti-HCV) identified in routine screening programs but who did not have acute hepatitis.

¶ Per 1,000 live births.

**TABLE 3. NOTIFIABLE DISEASES — summary of reported cases, United States, 1974-1983**

Disease	1983	1982	1981	1980	1979	1978	1977	1976	1975	1974
U.S. total resident population (in thousands) 1980 census; July 1 est. 1974-1979, 1981-1983	233,981	231,534	229,307	226,505	220,099	218,059	216,332	214,659	213,121	211,390
Amebiasis	6,658	7,304	6,632	5,271	4,107	3,937	3,044	2,906	2,775	2,743
Anthrax	-	-	-	1	-	6	-	2	2	2
Aseptic meningitis	12,696	9,680	9,547	8,028	8,754	6,573	4,789	3,510	4,475	3,197
Botulism, total (including wound and unsp.)	133	97	103	89	45	105	129	55	20	28
Brucellosis	200	173	185	183	215	179	232	296	310	240
Chancroid	847	1,392	850	788	840	521	455	628	700	945
Cholera	1	-	19	9	1	12	3	-	-	-
Diphtheria	5	2	5	3	59	76	84	128	307	272
Encephalitis, primary	1,761	1,464	1,492	1,362	1,504	1,351	1,414	1,651	4,064	1,164
Post-infectious	34	36	43	40	84	78	119	175	237	218
Gonorrhea	900,435	960,633	990,864	1,004,029	1,004,058	1,013,436	1,002,219	1,001,994	999,937	906,121
Granuloma inguinale	24	17	66	51	76	72	75	71	60	47
Hansen disease (leprosy)	259	250	256	223	185	168	151	145	162	118
Hepatitis A	21,532	23,403	25,802	29,087	30,407	29,500	31,153	33,288	35,855	40,358
Hepatitis B	24,318	22,177	21,152	19,015	15,452	15,016	16,831	14,973	13,121	10,631
Hepatitis, non-A, non-B	3,470	-	-	-	-	*	-	-	-	-
Hepatitis, unspecified	7,149	8,564	10,975	11,894	10,534	8,776	8,639	7,488	7,158	8,351
Legionellosis <sup>†</sup>	852	654	408	475	593	761	359	235	.....*	.....*
Leptospirosis	61	100	82	85	94	110	71	73	93	68
Lymphogranuloma venereum	335	235	263	199	250	284	348	365	353	394
Malaria	813	1,056	1,388	2,062	894	731	547	471	373	293
Measles (rubeola)	1,497	1,714	3,124	13,506	13,597	26,871	57,345	41,126	24,374	22,094
Meningococcal infections	2,736	3,056	3,525	2,840	2,724	2,505	1,828	1,605	1,478	1,346
Mumps	3,355	5,270	4,941	8,576	14,225	16,817	21,436	38,492	59,647	59,128
Murine typhus fever	62	58	61	81	69	46	75	69	41	26
Pertussis (whooping cough)	2,463	1,895	1,248	1,730	1,623	2,063	2,177	1,010	1,738	2,402
Plague	40	19	13	18	13	12	18	16	20	8
Poliomyelitis, total	15	8	6	9	34	15	18	14	8	7
Paralytic	15	8	6	8	26	9	17	12	8	7
Psittacosis	142	152	136	124	137	140	94	78	49	164
Rabies, animal	5,878	6,212	7,118	6,421	5,119	3,254	3,130	3,073	2,627	3,151
Rabies, human	2	-	2	-	4	4	1	2	2	-
Rheumatic fever, acute	88	137	264	432	629	851	1,738	1,865	2,854	2,431
Rocky Mountain spotted fever	1,126	976	1,192	1,163	1,070	1,063	1,153	937	844	754
Rubella (German measles)	970	2,325	2,077	3,904	11,795	18,269	20,395	12,491	16,652	11,917
Rubella, congenital syndrome	22	7	19	50	62	30	23	30	30	45
Salmonellosis, excluding typhoid fever	44,250	40,936	39,990	33,715	33,138	29,410	27,850	22,937	22,612	21,980
Shigellosis	19,719	18,129	19,859	19,041	20,135	19,511	16,052	13,140	16,584	22,600
Smallpox	..... Last documented case occurred in 1949 .....									
Syphilis, primary and secondary	32,698	33,613	31,266	27,204	24,874	21,656	20,399	23,731	25,561	25,385
Total, all stages	74,637	75,579	72,799	68,832	67,049	64,875	64,621	71,761	80,356	83,771
Tetanus	91	88	72	95	81	86	87	75	102	101
Toxic-shock syndrome	502	-	-	-	-	*	-	-	-	-
Trichinosis	45	115	206	131	157	67	143	115	252	120
Tuberculosis <sup>§</sup>	23,846	25,520	27,373	27,749	27,669	28,521	30,145	32,105	33,989	30,122
Tularemia	310	275	288	234	196	141	165	157	129	144
Typhoid fever	507	425	584	510	528	505	398	419	375	437
Varicella (chickenpox)	177,462	167,423	200,766	190,894	199,081	154,089	188,396	183,990	154,248	141,495
Yellow fever	..... Last indigenous case reported 1911, last imported 1924 .....									

\* Not previously notifiable nationally.

<sup>†</sup> Data for 1982-1983 recorded by date of report to the state health department. Data for 1976-1981 are from surveillance records reported by onset date.

<sup>§</sup> Case data subsequent to 1974 are not comparable to prior years due to changes in reporting criteria that became effective in 1975.

TABLE 4. NOTIFIABLE DISEASES — summary of reported cases, United States, 1964–1973

Disease	1973	1972	1971	1970	1969	1968	1967	1966	1965	1964
U.S. total resident population, July 1, estimate (in thousands)	209,851	208,232	206,256	203,805	201,385	199,399	197,457	195,576	193,526	191,141
Amebiasis	2,235	2,199	2,752	2,888	2,915	3,005	3,157	2,921	2,768	3,304
Anthrax	2	2	5	2	4	3	2	5	7	5
Aseptic meningitis	4,846	4,634	5,176	6,480	3,672	4,494	3,082	3,058	2,329	2,177
Botulism	34	22	25	12	16	7	5	9	19	23
Brucellosis	202	196	183	213	235	218	265	262	262	411
Chancroid	1,165	1,414	1,320	1,416	1,104	845	784	838	982	1,247
Cholera	1	—	1	—	—	—	—	—	2	—
Diphtheria	228	152	215	435	241	260	219	209	164	293
Encephalitis, primary	1,613	1,059	1,524	1,580	1,613	1,781	1,478	2,121	1,722	2,002
Post-infectious	354	243	439	370	304	502	1,060	964	981	1,585
Gonorrhoea	842,621	767,215	670,268	600,072	534,872	464,543	404,836	351,738	324,925	300,666
Granuloma inguinale	62	81	89	124	154	156	154	148	155	135
Hansen disease (leprosy)	146	130	131	129	98	123	81	109	96	97
Hepatitis A	50,749	54,074	59,606	56,797	48,416	45,893	38,909	32,859	33,856*	37,740*
Hepatitis B	8,451	9,402	9,556	8,310	5,909	4,829	2,458	1,497	—	—
Leptospirosis	57	41	62	47	89	69	67	72	84	142
Lymphogranuloma venereum	408	756	692	612	520	485	371	308	878	732
Malaria	237	742	2,375	3,051	3,102	2,317	2,022	565	147	93
Measles (rubeola)	26,690	32,275	75,290	47,351	25,826	22,231	62,705	204,136	261,904	458,083
Meningococcal infections	1,378	1,323	2,262	2,505	2,951	2,623	2,161	3,381	3,040	2,826
Mumps	69,612	74,215	124,939	104,953	90,918	152,209	..... †	..... †	..... †	..... †
Murine typhus fever	32	18	23	27	36	36	52	33	28	30
Pertussis (whooping cough)	1,759	3,287	3,036	4,249	3,285	4,810	9,718	7,717	6,799	13,005
Plague	2	1	2	13	5	3	3	5	8	—
Poliomyelitis, total	8	31	21	33	20	53	41	113	72	122
Paralytic	7	29	17	31	18	53	40	106	61	106
Psittacosis	33	52	32	35	57	43	41	50	60	53
Rabies, animal	3,640	4,369	4,310	3,224	3,490	3,591	4,481	4,178	4,574	4,780
Rabies, human	1	2	2	3	1	1	2	1	2	1
Rheumatic fever, acute	2,560	2,614	2,793	3,227	3,229	3,470	3,985	4,472	4,998	7,491
Rocky Mountain spotted fever	668	523	432	380	498	298	305	268	281	277
Rubella (German measles)	27,804	25,507	45,086	56,552	57,686	49,371	46,888	46,975	..... †	..... †
Rubella, congenital syndrome	35	42	68	77	31	14	10	11	..... †	..... †
Salmonellosis, excluding typhoid fever	23,818	22,151	21,928	22,096	18,419	16,514	18,120	16,841	17,161	17,144
Shigellosis	22,642	20,207	16,143	13,845	11,946	12,180	13,474	11,888	11,027	12,984
Smallpox	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
Streptococcal sore throat and scarlet fever	NN	NN	NN	433,405	450,008	435,013	453,351	427,752	395,168	402,334
Syphilis, primary and secondary	24,825	24,429	23,783	21,982	19,130	19,019	21,053	21,414	23,338	22,969
Total, all stages	87,469	91,149	95,997	91,382	92,162	96,271	102,581	105,159	112,842	114,325
Tetanus	101	128	116	148	192	178	263	235	300	289
Trichinosis	102	89	103	109	215	77	66	115	199	198
Tuberculosis <sup>§</sup>	30,998	32,882	35,217	37,137	39,120	42,623	45,647	47,767	49,016	50,874
Tularemia	171	152	187	172	149	186	184	208	264	342
Typhoid fever	680	398	407	346	364	395	396	378	454	501
Varicella (chickenpox)	182,927	164,114	..... †	..... †	..... †	..... †	..... †	..... †	..... †	..... †
Yellow fever	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....

\*Includes Hepatitis B.

† Not previously notifiable nationally.

§ Includes new active cases.

**TABLE 5. NOTIFIABLE DISEASES — summary of reported cases, United States, 1954-1963**

Disease	1963	1962	1961	1960	1959	1958	1957	1956	1955	1954
U.S. total resident population, July 1, estimate (in thousands)	188,483	185,771	182,992	179,979	176,513	173,320	170,371	167,306	164,308	161,164
Amebiasis	2,886	3,048	2,850	3,424	3,508	4,380	5,031	3,689	3,348	3,523
Anthrax	3	9	14	23	12	16	26	38	39	22
Aseptic meningitis	1,844	2,654	5,162*	1,593	.....	.....	.....	.....	.....	.....
Botulism	47	10	14	12	20	6	28	17	16	18
Brucellosis	407	409	636	751	892	924	983	1,300	1,444	1,823
Chancroid	1,220	1,344	1,438	1,680	1,537	1,595	1,637	2,135	2,649	3,003
Cholera	-	-	-	-	-	-	-	-	-	-
Dengue	NN	NN	-	-	-	-	-	2	1	6
Diphtheria	314	444	617	918	934	918	1,211	1,568	1,984	2,041
Encephalitis, acute infectious	1,993	2,094	2,248	2,341	2,347	2,587	2,135	2,624	2,166	2,606
Gonorrhea	278,289	263,714	264,158	258,933	240,254	232,386	214,496	224,346	236,197	242,050
Granuloma inguinale	173	207	241	296	265	314	348	357	490	618
Hansen disease (leprosy)	103	80	63	54	44	39	36	52	75	56
Hepatitis <sup>§</sup>	42,974	53,016	72,651	41,666	23,574	16,294	14,922	19,234	31,961	50,093
Leptospirosis	89	79	71	53	83	55	47	44	24	48
Lymphogranuloma venereum	586	590	787	835	604	434	448	500	762	875
Malaria	99	118	73	72	71	85	132	234	522	715
Measles (rubeola)	385,156	481,530	423,919	441,703	406,162	763,094	486,799	611,936	555,156	682,720
Meningococcal infections	2,470	2,150	2,232	2,259	2,180	2,581	2,691	2,735	3,455	4,436
Murine typhus fever	35	32	46	68	51	71	113	98	135	163
Pertussis (whooping cough)	17,135	17,749	11,468	14,809	40,005	32,148	28,295	31,732	62,786	60,886
Plague	1	-	3	2	4	-	1	1	-	-
Poliomyelitis, total	449	910	1,312	3,190	8,425	5,787	5,485	15,140	28,985	38,476
Paralytic	396	762	988	2,525	6,289	3,697	2,499	7,911	13,850	18,308
Psittacosis	76	79	102	113	147	158	278	568	334	563
Rabies, animal	3,929	3,732	3,599	3,567	4,177	4,787	4,542	5,681	5,799	7,297
Rabies, human <sup>¶</sup>	1	2	3	2	7	5	5	10	4	13
Rheumatic fever, acute	7,561	7,977	10,470	9,022	8,285	6,889	6,427	6,562	.....	.....
Rocky Mountain spotted fever	216	240	219	204	199	243	240	293	295	294
Salmonellosis, excluding typhoid fever	15,390	9,680	8,542	6,929	6,606	6,363	6,693	6,704	5,447	5,375
Shigellosis	13,009	12,443	12,571	12,487	12,888	11,861	9,822	10,306	13,912	13,846
Smallpox	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
Streptococcal sore throat and scarlet fever	342,161	315,809	338,410	315,173	334,715	264,097	226,973	176,392	147,502	147,785
Syphilis, primary and secondary	22,251	21,067	19,851	16,145	9,799	7,176	6,576	6,392	6,454	7,147
Total, all stages	124,137	126,245	124,658	122,538	120,824	113,884	123,758	130,201	122,392	130,697
Tetanus	325	322	379	368	445	445	447	468	462	524
Trichinosis	208	194	306	160	227	176	178	262	264	277
Tuberculosis**	54,042	53,315	53,726	55,494	57,535	63,534	67,149	69,895	77,368	79,775
Tularemia	327	328	365	390	459	587	601	522	584	681
Typhoid fever	566	608	814	816	859	1,043	1,231	1,700	1,704	2,169
Yellow fever	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....

\*Includes Meningitis, other, for some states.  
 †Not previously notifiable nationally.  
 §Data includes both hepatitis A and B.  
 ¶Registered deaths, 1954-1961.  
 \*\*Includes new active cases.

TABLE 6. NOTIFIABLE DISEASES — summary of reported cases, United States, 1944–1953

Disease	1953	1952	1951	1950	1949	1948	1947	1946	1945	1944
U.S. total resident population, July 1, estimate (in thousands)	158,242	155,687	153,310	151,235	148,665	146,093	143,446	140,054	132,481	132,885
Amebiasis	4,444	4,280	3,550	4,568	5,543	4,871	3,365	4,093	3,412	3,241
Anthrax	45	47	60	49	54	60	69	40	40	49
Botulism	18	18	33	20	24	39	44	.....	NA	.....
Brucellosis	2,032	2,537	3,139	3,510	4,235	4,991	6,321	5,887	5,049	4,436
Chancroid*	3,338	3,738	4,233	4,977	6,707	7,661	9,515	7,091	5,515	7,878
Cholera	-	-	-	-	-	-	-	-	-	-
Dengue	8	5	16	26	46	24	35	40	106	61
Diphtheria	2,355	2,960	3,983	5,796	7,969	9,493	12,262	16,354	18,675	14,150
Encephalitis, acute infectious	1,935	1,912	1,123	1,135	903	730	785	728	785	788
Gonorrhoea*	238,340	244,957	254,470	286,746	317,950	345,501	380,666	368,020	287,181	300,676
Granuloma inguinale*	667	951	1,352	1,783	2,402	2,469	2,330	2,232	1,857	1,759
Hansen disease (leprosy)	60	57	57	44	41	63	56	43	40	37
Hepatitis, infectious <sup>†</sup>	33,700	17,428	7,349	2,820	2,027	709	1,092	.....	NA	.....
Leptospirosis	42	62	9	30	17	18	14	.....	NA	.....
Lymphogranuloma venereum*	983	1,200	1,300	1,427	1,925	2,429	2,526	2,603	2,631	2,858
Malaria	1,310	7,023	5,600	2,184	4,151	9,606	15,116	48,610	62,763	57,626
Measles (rubeola)	449,146	683,077	530,118	319,124	625,281	615,104	222,375	659,843	146,013	630,291
Meningococcal infections	5,077	4,884	4,164	3,788	3,519	3,376	3,420	5,693	8,208	16,312
Murine typhus fever	221	205	378	685	985	1,171	2,050	3,365	5,193	5,401
Pertussis (whooping cough)	37,129	45,030	68,687	120,718	69,479	74,715	156,517	109,860	133,792	109,873
Plague	-	-	1	3	3	-	1	-	-	1
Poliomyelitis, total	35,592	57,879	28,386	33,300	42,033	27,726	10,827	25,698	13,624	19,029
Psittacosis	169	135	25	26	35	32	27	26	27	6
Rabies, animal <sup>§</sup>	8,903	8,445	8,008	7,901	7,587	8,495	8,920	10,850	9,928	10,487
Rabies, human <sup>  </sup>	12	24	18	18	10	24	26	34	43	56
Rocky Mountain spotted fever	313	327	347	464	570	547	596	587	472	470
Salmonellosis, excluding typhoid fever	3,946	2,596	1,773	1,233	1,243	882	951	723	649	712
Shigellosis	16,533	23,197	32,215	23,367	29,080	23,753	17,048	24,286	34,943	38,230
Smallpox	-	-	-	-	49	57	176	337	346	397
Streptococcal sore throat and scarlet fever	132,935	113,677	84,151	64,494	87,220	91,295	93,595	125,511	185,570	200,539
Syphilis, primary and secondary*	8,637	10,449	14,485	23,939	41,942	68,174	93,545	94,957	77,007	78,443
Total, all stages*	148,573	167,762	174,924	217,558	256,463	314,313	355,592	963,647	359,114	467,755
Tetanus	506	484	506	486	579	601	560	.....	NA	.....
Trichinosis	395	367	393	327	353	487	451	.....	NA	.....
Tuberculosis**	84,304	86,700	118,491	121,742	134,865	137,006	134,946	119,256	114,931	126,294
Tularemia	601	668	702	927	1,179	1,086	1,401	1,355	900	781
Typhoid fever	2,252	2,341	2,128	2,484	2,795	2,840	3,075	3,268	4,211	4,599
Yellow fever	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....

\*Data were reported for fiscal years 1944–1946; data were reported by calendar year beginning in 1947.

<sup>†</sup>Data for 1953 includes serum hepatitis.

<sup>§</sup>Data for 1944–1951 from Bureau of Animal Industry, U.S. Department of Agriculture, Agricultural Research Administration.

<sup>||</sup>Registered deaths.

\*\*Includes newly reported active and inactive cases, 1944–1951; new active cases 1952–1953.



**TABLE 7. NOTIFIABLE DISEASES — deaths from specified notifiable diseases, United States, 1982–1991**

Cause of Death	ICD*	1991	1990	1989	1988	1987	1986	1985	1984	1983	1982
AIDS <sup>†</sup>	*042–*044	29,555	25,188	22,082	16,602	13,468	10,900	6,040	2,943	1,141	NA
Amebiasis	006	5	5	4	7	9	8	10	10	21	7
Anthrax	022	–	–	–	–	–	–	–	–	–	–
Botulism, foodborne	005.1	2	4	2	1	–	1	4	4	7	4
Brucellosis	023	–	–	–	2	1	1	1	–	–	2
Chancroid	099.0	1	–	–	–	–	–	–	–	–	–
Cholera	001	2	2	–	–	1	–	1	–	–	–
Diphtheria	032	–	1	–	–	1	–	–	–	–	1
Encephalitis, acute infectious <sup>§</sup>	062–064,049	142	167	143	133	146	230	153	168	169	166
Gonococcal infections	098	3	3	4	3	7	7	2	3	4	6
Granuloma inguinale	099.2	–	–	–	–	–	–	–	–	–	–
<i>Haemophilus influenzae</i>	041.5	17	16	16	25	25	21	22	14	11	24
Hansen disease (leprosy)	030	–	3	4	–	1	1	2	6	3	3
Hepatitis, viral, infectious (Hep A)	070.0,070.1	71	76	88	70	77	65	80	77	82	83
Hepatitis, viral, serum (Hep B)	070.2,070.3	912	816	711	621	595	557	490	465	438	375
Hepatitis, viral, other and unsp.	070.4–070.9	857	686	717	599	510	384	372	327	343	356
Leptospirosis	100	1	2	–	2	1	–	4	–	5	4
Lymphogranuloma venereum	099.1	1	2	2	–	–	–	3	–	–	1
Malaria	084	4	3	11	7	5	5	13	7	3	2
Measles (rubeola)	055	27	64	32	3	2	2	4	1	4	2
Meningococcal infections	036	198	215	273	278	258	286	257	300	299	364
Mumps	072	1	1	3	2	2	–	–	1	2	2
Murine typhus fever	081.0	–	–	1	–	–	–	1	–	–	–
Pertussis (whooping cough)	033	–	12	12	4	1	6	4	7	5	4
Plague	020	–	–	–	–	1	–	1	3	5	3
Poliomyelitis, total	045.0–045.9	1	–	–	1	–	–	3	–	–	–
Psittacosis	073	–	2	1	1	2	–	1	–	1	–
Rabies, human	071	3	1	1	–	1	–	–	2	2	–
Rheumatic fever, acute	390–392	89	66	70	76	42	60	56	70	87	77
Rocky Mountain spotted fever	082.0	13	20	10	20	21	19	22	34	35	40
Rubella (German measles)	056	1	8	4	1	–	1	1	1	3	4
Salmonellosis, incl. paratyphoid fever	002.1–002.9,003	53	80	99	66	105	102	117	90	82	89
Shigellosis	004	10	10	16	8	13	4	17	8	9	9
Syphilis	090–097	93	106	105	85	98	80	80	105	121	126
Tetanus	037	11	11	9	17	16	22	23	20	22	22
Trichinosis	124	–	–	1	–	–	–	1	–	–	–
Tuberculosis (all forms)	010–018	1,713	1,810	1,970	1,921	1,755	1,782	1,752	1,729	1,779	1,807
Tularemia	021	2	1	1	2	4	4	3	2	1	2
Typhoid fever	002.0	1	1	–	–	2	2	–	–	3	2
Varicella (chickenpox)	052	81	120	89	83	89	47	68	53	57	61

\* Numbers in ICD column refer to the category numbers listed in the *Ninth Revision of the International Classification of Diseases, 1975*. (The asterisks in the ICD column pertain to the ICD code, not a footnote. They indicate that the numbers are not part of the ICD but were introduced for use in the United States.)

<sup>†</sup> For 1983–1986, deaths are estimated from death certificates with mention of conditions coded to deficiency of cell-mediated immunity (ICD-9 No. 279.1). Includes other human immunodeficiency virus (HIV)-related deaths and other diseases classifiable as deficiency of cell-mediated immunity.

<sup>§</sup> Arthropod-borne encephalitis and other nonarthropod-borne viral diseases of the central nervous system.

Source: National Center for Health Statistics System, 1982–1991. Deaths are classified according to the Ninth Revision, ICD.

## State and Territorial Epidemiologists and Laboratory Directors

State and Territorial Epidemiologists and Laboratory Directors are gratefully acknowledged for their contributions to this report. The epidemiologists listed below were in the positions shown as of July 1994, and the laboratory directors listed below were in the positions shown as of June 1994.

<b>State/Territory</b>	<b>Epidemiologist</b>	<b>Laboratory Director</b>
Alabama	Charles H. Woernle, MD, MPH	William J. Callan, PhD
Alaska	John P. Middaugh, MD	Katherine A. Kelley, DrPH
Arizona	Lawrence Sands, DO, MPH	Barbara J. Erickson, PhD
Arkansas	Thomas C. McChesney, DVM	Michael G. Foreman
California	George W. Rutherford, III, MD	Michael G. Volz, PhD
Colorado	Richard E. Hoffman, MD, MPH	Ronald L. Cada, DrPH
Connecticut	James L. Hadler, MD, MPH	Sanders F. Hawkins, PhD
Delaware	A. LeRoy Hathcock, Jr, PhD	Mahadeo P. Verma, PhD
District of Columbia	Martin E. Levy, MD, MPH	James B. Thomas, ScD
Florida	Richard S. Hopkins, MD, MSPH	E. Charles Hartwig, ScD
Georgia	Kathleen E. Toomey, MD, MPH	Elizabeth A. Franko, DrPH
Hawaii	Richard L. Vogt, MD	Vernon K. Miyamoto, PhD
Idaho	Jesse F. Greenblatt, MD, MPH	Richard H. Hudson, PhD
Illinois	Byron J. Francis, MD, MPH	David F. Carpenter, PhD
Indiana	Mary Lou Fleissner, DrPH	Barbara J. Wilder (MBA (Acting))
Iowa	Laverne A. Wintermeyer, MD	W. J. Hausler, Jr, PhD
Kansas	Andrew R. Pelletier, MD	Roger H. Carlson, PhD
Kentucky	Reginald Finger, MD, MPH	Thomas E. Maxson, DrPH
Louisiana	Louise McFarland, DrPH	Henry B. Bradford, Jr, PhD
Maine	Kathleen F. Gensheimer, MD, MPH	Philip W. Haines, DrPH
Maryland	Ebenezer Israel, MD, MPH	J. Mehzen Joseph, PhD
Massachusetts	Alfred DeMaria, Jr, MD	Ralph J. Timperi, MPH
Michigan	Kenneth R. Wilcox, Jr, MD, DrPH	Robert Martin, DrPH
Minnesota	Michael T. Osterholm, PhD, MPH	Pauline Bouchard, JD, MPH
Mississippi	Mary Currier, MD, MPH	Joe O. Graves, PhD
Missouri	H. Denny Donnell, Jr, MD, MPH	Eric C. Blank, DrPH
Montana	Todd D. Damrow, PhD, MPH	Douglas O. Abbott, PhD
Nebraska	Thomas J. Safraneck, MD	John D. Blosser
Nevada	Randall L. Todd, DrPH	Arthur F. DiSalvo, MD
New Hampshire	M. Geoffrey Smith, MD, MPH	Veronica C. Malmberg, MSN
New Jersey	Kenneth C. Spitalny, MD	Shahiedy I. Shahied, PhD
New Mexico	C. Mack Sewell, DrPH, MS	Loris W. Hughes, PhD
New York State	Susan Klitzman	Lawrence S. Sturman, MD, PhD
North Carolina	J. Newton MacCormack, MD, MPH	Samuel N. Merritt, DrPH
North Dakota	Larry A. Shireley, MS, MPH	James D. Anders, MPH
Ohio	Thomas J. Halpin, MD, MPH	Kathleen Meckstroth, DrPH
Oklahoma	James T. Rankin, Jr, DVM, PhD, MPH	Garry L. McKee, PhD
Oregon	David Fleming, MD	—
Pennsylvania	Maria E. Moll, MD	Bruce Kieger, DrPH
Rhode Island	Barbara A. DeBuono, MD, MPH	Walter Combs, PhD
South Carolina	James J. Gibson, MD, MPH	Harold Dowda, PhD
South Dakota	Susan Lance, DVM, MPH	Richard S. Steece, PhD
Tennessee	Kerry Gateley, MD	Michael W. Kimberly, DrPH
Texas	Diane M. Simpson, MD, PhD	David L. Maserang, PhD
Utah	Craig R. Nichols, MPA	Charles D. Brokopp, DrPH
Vermont	Robert O'Grady (Acting)	Burton W. Wilcke, Jr, PhD
Virginia	Grayson B. Miller, Jr, MD	James L. Pearson, DrPH
Washington	Paul Stehr-Green, DrPH	Jon M. Counts, DrPH
West Virginia	Loretta E. Haddy, MA, MS	Frank W. Lambert, Jr, DrPH
Wisconsin	Jeffrey P. Davis, MD	Ronald H. Laessig, PhD
Wyoming	Stanley I. Music, MD, DTPH	Carl H. Blank, DrPH
American Samoa	Julia L. Lyons, MD, MPH	—
Federated States of Micronesia	Steven B. Auerbach, MD, MPH	—
Guam	Robert L. Haddock, DVM, MPH	Jeff Benjamin (Acting)
Marshall Islands	Tony de Brum	—
Northern Mariana Islands	A. Mark Durand, MD, MPH	—
Palau	Jill McCready, MS, MPH	—
Puerto Rico	Carmen C. Deseda, MD	Adolpho Firpo Reforma, MD
Virgin Islands	Donna M. Green, MD	Norbert Mantor, PhD

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available on a paid subscription basis from the Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone (202) 783-3238.

The data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the succeeding Friday. Inquiries about the *MMWR* Series, including material to be considered for publication, should be directed to: Editor, *MMWR* Series, Mailstop C-08, Centers for Disease Control and Prevention, Atlanta, GA 30333; telephone (404) 332-4555.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without special permission; citation as to source, however, is appreciated.