

Executive Commentary

Executive Commentary

Highlights of 2007 Report

Since 1953, in cooperation with state and local health departments, the Centers for Disease Control and Prevention (CDC) have collected information on each newly reported case of tuberculosis (TB) disease in the United States. Currently, each individual TB case report (Report of Verified Case of Tuberculosis, or RVCT) is submitted electronically to CDC's Division of Tuberculosis Elimination. The following are the highlights of the 2007 report:

1. Updated case counts for each year from 1993 through 2006.
2. Case counts: 13,299 TB cases were reported to CDC from the 50 states and the District of Columbia (DC) for 2007, representing a 3.3% decrease from 2006 (Table 1).
 - Twenty-one states reported increased case counts from 2006 (Table 28).
 - California, Texas, New York, and Florida accounted for 48% of the national case total (Table 28).
 - For the fourth consecutive year, Hispanics (29%) exceeded all other racial or ethnic groups with the largest percentage of total cases (Table 2).
 - For the first time, Asians (26%) matched non-Hispanic blacks or African-Americans¹ (26%) as the second largest racial or ethnic group.
 - Blacks or African-Americans born in the United States represented 45% of TB cases in U.S.-born persons and accounted for approximately 18% of the national case total (Tables 17, 18).
 - Asians born outside the United States represented 43% of TB cases in foreign-born persons and accounted for approximately 25% of the national case total (Tables 17, 18).
3. Case rates: In 2007, the TB case rate declined from 4.6 to 4.4 per 100,000 persons, representing a 4.2% decrease from 2006.
 - Fifteen states and DC reported rates above the national average (Table 20).
 - Twenty-six states met the definition for low incidence, or ≤ 3.5 cases per 100,000 population (Table 20).
 - The TB case rate was 2.1 per 100,000 for U.S.-born persons and 20.7 for foreign-born persons (Table 5).
 - Asians continued to have the highest case rate (26.3 per 100,000 persons) among all racial or ethnic groups (Table 2).
4. Burden among the foreign-born: The percentage of cases occurring in foreign-born persons continued to increase and was 58% of the national case total.
 - Foreign-born Hispanics and Asians together represented 81% of TB cases in foreign-born persons, and accounted for 47% of the national case total (Tables 17, 18).
 - In 29 states, $\geq 50\%$ of TB cases occurred among foreign-born persons (Table 23).
 - In 13 states, $\geq 70\%$ of TB cases occurred among foreign-born persons (Table 23).
 - The top five countries of origin of foreign-born persons with TB were Mexico, Philippines, India, Vietnam, and China (Table 6).
5. Drug resistance: 1.1% of reported cases, compared to 1.0% in 2006, had primary multidrug resistance, which is defined as no previous history of TB disease and resistance to at least isoniazid and rifampin (Table 10).

¹Hispanic and non-Hispanic are ethnicities. All races are non-Hispanic. The category "non-Hispanic blacks or African-Americans" includes U.S.-born and foreign-born persons unless otherwise specified.

Tuberculosis in the United States

In 2007, the number of TB cases reported (13,299) and case rate (4.4 cases per 100,000) both decreased; this represented declines of 3.3% and 4.2%, respectively, compared to 2006. Since the 1992 TB resurgence peak in the United States, the number of TB cases reported annually has decreased by 50%. However, the trend of the declining annual case rate has slowed, from an annual average decline of 7.3% for 1993 through 2000 to an annual average decline of 3.8% for 2000 through 2007 (Table 1).

The proportion of total cases occurring in foreign-born persons has been increasing since 1993. In 2007, 58% of TB cases occurred in foreign-born persons. Foreign-born persons have accounted for the majority of TB cases in the United States every year since 2001. Moreover, the case rate among foreign-born persons in 2007 was nearly 10 times higher than among U.S.-born persons (Table 5).

Tuberculosis deaths decreased by 0.6%, from 648 deaths in 2005 to 644 deaths in 2006, the latest year for which complete data are available (Table 1).

Age

Since 1993, TB case rates have declined annually for all age groups. TB case rates vary by well-known factors such as age, race and ethnicity, and country of origin. In 2007, TB case rates declined or remained constant for all age groups except adults aged 15 to 24 years, which increased slightly compared to 2006. The highest burden of disease continues to be among older adults. In 2007, adults aged 65 years and older had a case rate of 6.8 cases per 100,000, while children aged <14 years had the lowest rate at 1.3 cases per 100,000 (Table 4).

Race and Ethnicity

In 2003, the race and ethnicity category, “non-

Hispanic, Asian or Pacific Islander,” was split into “non-Hispanic Asian” and “non-Hispanic Native Hawaiian or Other Pacific Islander.” In 2007, Asians had the highest TB case rate at 26.3 cases per 100,000, which was a slight increase from 25.9 in 2006. In 2007, Native Hawaiians or Other Pacific Islanders had the second-highest TB case rate at 23.0 cases per 100,000, which is a marked increase compared to 13.4 cases per 100,000 reported in 2006 (Table 2).

Since 1993, TB case rates have declined between 57% and 69% in these racial and ethnic groups: among Hispanic or Latinos from 19.9 to 8.5 cases per 100,000; among blacks or African-Americans from 28.5 to 9.4 cases per 100,000; and among non-Hispanic whites from 3.6 to 1.1 cases per 100,000. In 2007, the TB case rate for Asians was nearly three times higher than that for Hispanics or blacks or African-Americans (Table 2).

Origin of Birth

Since 1993, the TB case rate among U.S.-born persons has declined annually. In 2007, the TB case rate for U.S.-born persons was 2.1 cases per 100,000, representing a 72% decrease from 7.4 cases per 100,000 in 1993. The TB case rate among foreign-born persons also declined during the same interval but was less substantial. In 2007, the TB case rate among foreign-born persons was 20.7 cases per 100,000 representing a 39% decrease from 34.0 cases per 100,000 in 1993 (Table 5).

The proportion of TB cases among persons born in the United States also has declined annually since 1993. In 2007, 42% of TB cases were among U.S.-born persons compared to 69% in 1993 (Table 5). In 29 states, $\geq 50\%$ of TB cases occurred among foreign-born persons. In 13 states (California, Colorado, Delaware, Massachusetts, Minnesota, Nebraska, New Hampshire, New Jersey, New York, Oregon, Rhode Island, Virginia, and Washington), $\geq 70\%$ of TB cases occurred among foreign-born persons (Table 23).

Country of Origin and World Region

From 2001 through 2007, the top five countries of origin of foreign-born persons with TB were Mexico, Philippines, India, Vietnam, and China (Table 6). However, the changes in the distribution of TB cases by world region of origin reflect the changing immigration patterns among persons settling in the United States². Of the 7,750 TB cases reported among foreign-born persons in 2007, 43% occurred among persons born in the Americas region, and 31% occurred among persons born in the Western Pacific region (Table 19). From 1993 to 2007, the proportion of cases increased among persons born in the Eastern Mediterranean region (3% in 1993 to 4.5% in 2007), the Southeast Asia region (6% in 1993 to 12% in 2007), and the African region (2% in 1993 and 7% in 2007) (Table 19).

Multidrug-Resistant Tuberculosis

Since 1993, when the RVCT was expanded to include drug-susceptibility results, the proportion of patients with primary multidrug-resistant TB (MDR TB), which is defined as no previous history of TB disease and resistance to at least isoniazid and rifampin, has decreased from 2.5% to 1.1% in 2007. Since 1998, the percentage of U.S.-born patients with MDR TB has remained $\leq 0.7\%$. However, of the total number of reported primary MDR TB cases, the proportion occurring in foreign-born persons increased from 25.5% (103 of 407) in 1993 to 80% (78 of 98) in 2007 (Table 10).

Extensively Drug-Resistant Tuberculosis

CDC has included an updated case count of extensively drug-resistant TB (XDR TB) cases from 1993 to 2007 in the slide set that accompanies this report. XDR TB is defined as resistance to isoniazid and rifampin plus resistance to any fluoroquinolone and at least one of three injectable second-line anti-TB drugs (i.e., amikacin,

kanamycin, or capreomycin)^{3,4}. Two cases of XDR TB were reported during 2007, compared to four cases in 2006.

Tuberculosis Therapy

The proportion of TB patients prescribed an initial treatment regimen of three or more anti-TB drugs increased annually from 72% in 1993 to 88% in 2007. The proportion of patients who completed therapy within 1 year increased from 64% in 1993 to 83% in 2005. The proportion of persons receiving directly observed therapy at least for a portion of the treatment duration also increased from 35% in 1993 to 86% in 2005, the latest year for which complete outcome data are available (Table 12).

Summary

Essential elements for controlling TB in the United States include sufficient resources, interventions targeted to populations at high risk for TB, and collaborative efforts with the international community to reduce the burden of TB globally.

During 1993 through 2007, TB case rates in the United States decreased for U.S.-born and foreign-born persons; however, the decrease among foreign-born persons continues to be less substantial. Despite the decreasing case rate among foreign-born persons, well over half of the TB cases in the United States in 2007 occurred in this population, and the case rate was more than 10 times higher than among U.S.-born persons. To address the high TB case rates among foreign-born persons, CDC is collaborating with other national and international public health organizations to: 1) improve overseas screening of immigrants and refugees by systematically monitoring and evaluating the screening process; 2) strengthen the current notification system that alerts local health departments about the arrival of immigrants or refugees who have suspected

²World Health Organization (WHO). *Global Tuberculosis Control 2008: Surveillance, Planning, and Finance*. Geneva, Switzerland: World Health Organization, 2008 (WHO/HTM/TB/2008.393).

³Centers for Disease Control and Prevention (CDC). Notice to readers: revised definition of XDR-TB. *Morbidity and Mortality Weekly Report* 2006; 55:1176.

⁴WHO. Extensively drug-resistant tuberculosis (XDR-TB): recommendations for prevention and control. *Weekly Epidemiol Record* 2006; 81: 430–432.

TB in order to enhance the evaluation and treatment of such persons; 3) improve coordination of TB control activities between the United States and Mexico to ensure completion of treatment among TB patients who cross the border; 4) test recent arrivals from high-incidence countries for latent TB infection and monitor treatment completion; and 5) survey foreign-born TB patients in the United States to determine opportunities for improving prevention and control interventions. In addition, CDC continues to strengthen collaborations with international partners, including the World Health Organization's Stop TB Partnership, to improve TB control in high-incidence countries.

Accelerating progress in national TB elimination activities will require broader prevention efforts among high-risk population groups such as black or African-American and Asian communities, persons who are incarcerated, persons with excess alcohol and drug use, persons with human immunodeficiency virus infection, and persons living in poverty with limited access to medical care and stable housing.

In addition, low-incidence areas in the United States require continued support to maintain the capacity and expertise needed to respond to future TB cases⁵ especially in light of changing immigration patterns. CDC has updated the comprehensive national action plan to reflect the alignment of CDC priorities with the 2000 Institute of Medicine report on TB and to ensure that priority prevention activities are undertaken with optimal collaboration and coordination among national and international public health partners^{6,7}.

⁵CDC. Progressing toward tuberculosis elimination in low-incidence areas of the United States: Recommendations of the Advisory Council for the Elimination of Tuberculosis. *Morbidity and Mortality Weekly Report* 2002; 51 (No. RR-5): 1–20.

⁶Institute of Medicine. *Ending Neglect: The Elimination of Tuberculosis in the United States*. Washington, DC: National Academy Press, 2000.

⁷CDC. *CDC's Response to Ending Neglect: The Elimination of Tuberculosis from the United States*. Atlanta, GA: U.S. Department of Health and Human Services, CDC, 2000.