

Table 36. Culture-Positive Tuberculosis Cases and Percentages with Drug-Susceptibility Results, by Resistance to INH or Multidrug Resistance: Reporting Areas, 2007

Reporting Area	Total Culture Positive Cases	Cases with Initial Drug-Susceptibility Testing Performed ¹		Resistance ²			
				Isoniazid ¹		Isoniazid and Rifampin ¹	
		No.	(%)	No.	(%)	No.	(%)
United States	10,426	9,858	(94.6)	762	(7.7)	119	(1.2)
Alabama	146	134	(91.8)	1	(0.7)	0	(0.0)
Alaska	45	45	(100.0)	0	(0.0)	0	(0.0)
Arizona	222	217	(97.7)	16	(7.4)	3	(1.4)
Arkansas	70	68	(97.1)	7	(10.3)	3	(4.4)
California	2,189	2,121	(96.9)	232	(10.9)	27	(1.3)
Colorado	74	74	(100.0)	9	(12.2)	1	(1.4)
Connecticut	94	94	(100.0)	12	(12.8)	3	(3.2)
Delaware	13	13	(100.0)	0	(0.0)	0	(0.0)
District of Columbia	48	48	(100.0)	1	(2.1)	0	(0.0)
Florida	812	760	(93.6)	62	(8.2)	6	(0.8)
Georgia	372	360	(96.8)	21	(5.8)	2	(0.6)
Hawaii	98	95	(96.9)	5	(5.3)	1	(1.1)
Idaho	7	7	(100.0)	1	(14.3)	0	(0.0)
Illinois	406	383	(94.3)	31	(8.1)	4	(1.0)
Indiana	106	106	(100.0)	10	(9.4)	1	(0.9)
Iowa	33	32	(97.0)	3	(9.4)	1	(3.1)
Kansas	46	41	(89.1)	1	(2.4)	0	(0.0)
Kentucky	102	97	(95.1)	4	(4.1)	0	(0.0)
Louisiana	174	157	(90.2)	4	(2.5)	0	(0.0)
Maine	16	16	(100.0)	0	(0.0)	0	(0.0)
Maryland	215	215	(100.0)	10	(4.7)	3	(1.4)
Massachusetts	159	134	(84.3)	8	(6.0)	1	(0.7)
Michigan	175	171	(97.7)	11	(6.4)	3	(1.8)
Minnesota	177	176	(99.4)	17	(9.7)	3	(1.7)
Mississippi	96	95	(99.0)	3	(3.2)	0	(0.0)
Missouri	99	97	(98.0)	4	(4.1)	1	(1.0)
Montana	9	9	(100.0)	1	(11.1)	0	(0.0)
Nebraska	21	19	(90.5)	0	(0.0)	0	(0.0)
Nevada	83	79	(95.2)	6	(7.6)	1	(1.3)
New Hampshire	9	5	(55.6)	--	--	--	--
New Jersey	370	368	(99.5)	35	(9.5)	8	(2.2)
New Mexico	44	44	(100.0)	3	(6.8)	1	(2.3)
New York State ³	192	187	(97.4)	16	(8.6)	0	(0.0)
New York City	713	698	(97.9)	79	(11.3)	7	(1.0)
North Carolina	273	273	(100.0)	11	(4.0)	2	(0.7)
North Dakota	4	4	(100.0)	0	(0.0)	0	(0.0)
Ohio	197	194	(98.5)	10	(5.2)	1	(0.5)
Oklahoma	91	86	(94.5)	0	(0.0)	0	(0.0)
Oregon	79	79	(100.0)	7	(8.9)	2	(2.5)
Pennsylvania	205	170	(82.9)	14	(8.2)	2	(1.2)
Rhode Island	16	16	(100.0)	0	(0.0)	0	(0.0)
South Carolina	178	171	(96.1)	6	(3.5)	4	(2.3)
South Dakota	8	8	(100.0)	0	(0.0)	0	(0.0)
Tennessee	163	160	(98.2)	5	(3.1)	3	(1.9)
Texas	1,193	980	(82.1)	44	(4.5)	11	(1.1)
Utah	30	30	(100.0)	2	(6.7)	1	(3.3)
Vermont	3	3	(100.0)	0	(0.0)	0	(0.0)
Virginia	230	207	(90.0)	17	(8.2)	5	(2.4)
Washington	235	233	(99.1)	24	(10.3)	2	(0.9)
West Virginia	24	20	(83.3)	0	(0.0)	0	(0.0)
Wisconsin	61	58	(95.1)	9	(15.5)	6	(10.3)
Wyoming	1	1	(100.0)	0	(0.0)	0	(0.0)
American Samoa ⁴
Fed. States of Micronesia ⁴	5	1	(20.0)	--	--	--	--
Guam ⁴	37	33	(89.2)	1	(3.0)	0	(0.0)
Marshall Islands ⁴	22	18	(81.8)	1	(5.6)	1	(5.6)
N. Mariana Islands ⁴	10	7	(70.0)	--	--	--	--
Puerto Rico ⁴	88	87	(98.9)	6	(6.9)	3	(3.4)
Republic of Palau ⁴
U.S. Virgin Islands ⁴

¹Patients tested to at least isoniazid and rifampin

²Isolates may be resistant to other drugs. Overall U.S. percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for ≥75% of cases.

³Excludes New York City.

⁴Not included in U.S. totals.

Note: Ellipses indicate data not available.