

**Table 14. Tuberculosis Cases and Percentages by Reason TB Therapy Stopped: United States, 1993–2002**

Year	Total Cases <sup>1</sup>	Completed Therapy		Moved		Lost		Refused		Died <sup>2</sup>		Unknown <sup>3</sup>	
	No.	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
1993	23,741	18,042	(76.0)	1,121	(4.7)	1,087	(4.6)	223	(0.9)	3,051	(12.9)	217	(0.9)
1994	23,051	17,761	(77.1)	1,194	(5.2)	738	(3.2)	183	(0.8)	2,743	(11.9)	432	(1.9)
1995	21,711	17,285	(79.6)	970	(4.5)	563	(2.6)	155	(0.7)	2,390	(11.0)	348	(1.6)
1996	20,298	16,508	(81.3)	783	(3.9)	521	(2.6)	156	(0.8)	1,991	(9.8)	339	(1.7)
1997	18,930	15,653	(82.7)	668	(3.5)	434	(2.3)	119	(0.6)	1,755	(9.3)	301	(1.6)
1998	17,584	14,772	(84.0)	534	(3.0)	400	(2.3)	104	(0.6)	1,578	(9.0)	196	(1.1)
1999	16,863	14,237	(84.4)	457	(2.7)	356	(2.1)	105	(0.6)	1,438	(8.5)	270	(1.6)
2000	15,785	13,396	(84.9)	410	(2.6)	390	(2.5)	112	(0.7)	1,294	(8.2)	183	(1.2)
2001	15,405	13,182	(85.6)	373	(2.4)	364	(2.4)	95	(0.6)	1,112	(7.2)	279	(1.8)
2002	14,555	12,333	(84.7)	343	(2.4)	357	(2.5)	85	(0.6)	1,063	(7.3)	374	(2.6)

<sup>1</sup>Includes all cases in persons reported as alive at diagnosis and taking one or more TB drugs.

<sup>2</sup>Died = died of any cause (not only TB).

<sup>3</sup>Includes cases in persons reporting reason therapy stopped = Other, Missing, or Unknown.

**Note:** Data for all years are updated through April 1, 2005.

Data complete through 2002 only. See Technical Notes (page 9) for details.