

Table 36. Culture-Positive Tuberculosis Cases and Percentages with Drug-Susceptibility Results, by Resistance to INH or Multidrug Resistance: Reporting Areas, 2006

Reporting Area	Total Culture Positive Cases	Cases with Initial Drug-Susceptibility Testing Performed ¹		Resistance ²			
				Isoniazid ¹		Isoniazid and Rifampin ¹	
		No.	(%)	No.	(%)	No.	(%)
United States	10,783	9,946	(92.2)	818	(8.2)	111	(1.1)
Alabama	174	138	(79.3)	3	(2.2)	1	(0.7)
Alaska	61	61	(100.0)	4	(6.6)	2	(3.3)
Arizona	229	201	(87.8)	17	(8.5)	2	(1.0)
Arkansas	72	69	(95.8)	1	(1.4)	0	(0.0)
California	2,241	2,136	(95.3)	209	(9.8)	31	(1.5)
Colorado	86	86	(100.0)	11	(12.8)	1	(1.2)
Connecticut	72	71	(98.6)	3	(4.2)	1	(1.4)
Delaware	23	23	(100.0)	0	(0.0)	0	(0.0)
District of Columbia	65	62	(95.4)	6	(9.7)	2	(3.2)
Florida	845	795	(94.1)	72	(9.1)	5	(0.6)
Georgia	388	368	(94.8)	27	(7.3)	0	(0.0)
Hawaii	83	82	(98.8)	9	(11.0)	0	(0.0)
Idaho	19	16	(84.2)	0	(0.0)	0	(0.0)
Illinois	437	404	(92.4)	26	(6.4)	1	(0.2)
Indiana	98	92	(93.9)	2	(2.2)	0	(0.0)
Iowa	29	29	(100.0)	1	(3.4)	0	(0.0)
Kansas	64	57	(89.1)	4	(7.0)	2	(3.5)
Kentucky	71	66	(93.0)	3	(4.5)	0	(0.0)
Louisiana	168	132	(78.6)	4	(3.0)	1	(0.8)
Maine	13	12	(92.3)	1	(8.3)	0	(0.0)
Maryland	196	195	(99.5)	13	(6.7)	0	(0.0)
Massachusetts	200	197	(98.5)	26	(13.2)	4	(2.0)
Michigan	176	173	(98.3)	10	(5.8)	1	(0.6)
Minnesota	177	175	(98.9)	19	(10.9)	2	(1.1)
Mississippi	88	85	(96.6)	4	(4.7)	0	(0.0)
Missouri	82	81	(98.8)	9	(11.1)	2	(2.5)
Montana	11	10	(90.9)	1	(10.0)	1	(10.0)
Nebraska	19	19	(100.0)	1	(5.3)	0	(0.0)
Nevada	87	86	(98.9)	8	(9.3)	0	(0.0)
New Hampshire	12	12	(100.0)	2	(16.7)	0	(0.0)
New Jersey	386	373	(96.6)	43	(11.5)	5	(1.3)
New Mexico	44	41	(93.2)	2	(4.9)	0	(0.0)
New York State ³	239	238	(99.6)	34	(14.3)	3	(1.3)
New York City	717	691	(96.4)	70	(10.1)	15	(2.2)
North Carolina	302	290	(96.0)	22	(7.6)	0	(0.0)
North Dakota	7	7	(100.0)	0	(0.0)	0	(0.0)
Ohio	183	175	(95.6)	13	(7.4)	1	(0.6)
Oklahoma	77	71	(92.2)	3	(4.2)	0	(0.0)
Oregon	73	72	(98.6)	3	(4.2)	1	(1.4)
Pennsylvania	245	206	(84.1)	22	(10.7)	6	(2.9)
Rhode Island	13	13	(100.0)	2	(15.4)	0	(0.0)
South Carolina	173	165	(95.4)	7	(4.2)	0	(0.0)
South Dakota	11	11	(100.0)	1	(9.1)	0	(0.0)
Tennessee	207	196	(94.7)	2	(1.0)	0	(0.0)
Texas	1,218	918	(75.4)	43	(4.7)	9	(1.0)
Utah	22	22	(100.0)	3	(13.6)	0	(0.0)
Vermont	8	7	(87.5)	1	(14.3)	1	(14.3)
Virginia	267	228	(85.4)	21	(9.2)	4	(1.8)
Washington	216	210	(97.2)	26	(12.4)	5	(2.4)
West Virginia	22	14	(63.6)	--	--	--	--
Wisconsin	64	62	(96.9)	3	(4.8)	2	(3.2)
Wyoming	3	3	(100.0)	0	(0.0)	0	(0.0)
American Samoa ⁴
Fed. States of Micronesia ⁴	26	21	(80.8)	0	(0.0)	0	(0.0)
Guam ⁴	32	29	(90.6)	2	(6.9)	0	(0.0)
Marshall Islands ⁴	17	16	(94.1)	1	(6.3)	1	(6.3)
N. Mariana Islands ⁴	14	6	(42.9)	--	--	--	--
Puerto Rico ⁴	105	99	(94.3)	5	(5.1)	1	(1.0)
Republic of Palau ⁴	2	1	(50.0)	--	--	--	--
U.S. Virgin Islands ⁴

¹Patients tested to at least isoniazid and rifampin

²Isolates may be resistant to other drugs. Overall U.S. percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for ≥75% of cases.

³Excludes New York City.

⁴Not included in U.S. totals. Data for 2006 are interim as of April 6, 2007.

Note: Ellipses indicate data not available.