

TABLE 1**The Top 20 most frequently reported Salmonella serotypes
from Human sources reported to CDC in 2002**

Human 2002			
Rank	Serotype	Reported	Percent
1	S. Typhimurium *	7062	21.9
2	S. Enteritidis	5116	15.8
3	S. Newport	4204	13.0
4	S. Heidelberg	1957	6.1
5	S. Javiana	1188	3.7
6	S. Montevideo	717	2.2
7	S. Muenchen	591	1.8
8	S. Oranienburg	585	1.8
9	S. Saintpaul	535	1.7
10	S. Infantis	463	1.4
11	S. Thompson	440	1.4
12	S. Paratyphi B var. L(+) tartrate+ (Formerly Java)	429	1.3
13	S. Braenderup	384	1.2
14	S. Agona	337	1.0
15	S. Hadar	331	1.0
16	S. Mississippi	313	1.0
17	S. Berta	297	0.9
18	S. I 4,[5],12:i:-	291	0.9
19	S. Typhi	289	0.9
20	S. Poona	271	0.8
	Sub Total	25800	79.9
	All Other Serotyped	4879	15.1
	Unknown	449	1.4
	Partially serotyped isolates	1137	3.5
	Rough or nonmotile isolates	43	0.1
	Sub Total	6508	20.1
	Total	32308	100.0
NOTE: ----- * Typhimurium includes var. 5- (Formerly var. Copenhagen)			

TABLE 1a

**The Top 20 most frequently reported Salmonella serotypes from
Clinical and Non-Clinical Nonhuman sources reported to CDC and NVSL in 2002**

Clinical Nonhuman 2002			
Rank	Serotype	Reported	Percent
1	S. Typhimurium *	1453	21.6
2	S. Newport	1008	15.0
3	S. Agona	409	6.1
4	S. Heidelberg	327	4.9
5	S. Derby	261	3.9
6	S. Anatum	246	3.7
7	S. Choleraesuis **	204	3.0
8	S. Montevideo	195	2.9
9	S. Kentucky	157	2.3
10	S. Senftenberg	156	2.3
11	S. Dublin	137	2.0
12	S. Muenster	115	1.7
13	S. Uganda	115	1.7
14	S. Infantis	99	1.5
15	S. Oranienburg	87	1.3
16	S. I 4,[5],12:i:-	82	1.2
17	S. Mbandaka	79	1.2
18	S. Enteritidis	68	1.0
19	S. Bredeney	64	0.9
20	S. Hadar	64	0.9
	Sub Total	5326	79.1
	All Other Serotyped	1364	20.2
	Partially serotyped isolates	3	0.0
	Rough or nonmotile isolates	44	0.7
	Sub Total	1411	20.9
	Total	6737	100.0
NOTE: * Typhimurium includes var. 5- (Formerly var. Copenhagen) ** Choleraesuis includes var. Decatur and Kunzendorf			

Non-Clinical Nonhuman 2002			
Rank	Serotype	Reported	Percent
1	S. Heidelberg	2214	24.0
2	S. Typhimurium *	1134	12.3
3	S. Kentucky	799	8.7
4	S. Montevideo	583	6.3
5	S. Senftenberg	478	5.2
6	S. Derby	343	3.7
7	S. Mbandaka	246	2.7
8	S. Hadar	245	2.7
9	S. Muenster	232	2.5
10	S. Enteritidis	227	2.5
11	S. Agona	222	2.4
12	S. Anatum	216	2.3
13	S. Meleagridis	187	2.0
14	S. Newport	155	1.7
15	S. Cerro	131	1.4
16	S. Infantis	116	1.3
17	S. Saintpaul	105	1.1
18	S. Braenderup	100	1.1
19	S. Schwarzengrund	86	0.9
20	S. Reading	75	0.8
	Sub Total	7894	85.6
	All Other Serotyped	1291	14.0
	Partially serotyped isolates	1	0.0
	Rough or nonmotile isolates	33	0.4
	Sub Total	1325	14.4
	Total	9219	100.0
NOTE: * Typhimurium includes var. 5- (Formerly var. Copenhagen)			