

TABLE 1**The Top 20 most frequently reported Salmonella serotypes from Human sources reported to CDC in 2003**

Human 2003			
Rank	Serotype	Reported	Percent
1	S. Typhimurium *	6631	19.7
2	S. Enteritidis	4863	14.5
3	S. Newport	3847	11.5
4	S. Heidelberg	1810	5.4
5	S. Javiana	1659	4.9
6	S. Montevideo	849	2.5
7	S. Saintpaul	823	2.5
8	S. Muenchen	781	2.3
9	S. Oranienburg	554	1.6
10	S. Infantis	539	1.6
11	S. Braenderup	530	1.6
12	S. Agona	510	1.5
13	S. Thompson	494	1.5
14	S. I 4,[5],12:i:-	489	1.5
15	S. Mississippi	438	1.3
16	S. Typhi	359	1.1
17	S. Paratyphi B var. L(+) tartrate+	331	1.0
18	S. Hadar	280	0.8
19	S. Bareilly	234	0.7
20	S. Stanley	224	0.7
	Sub Total	26245	78.1
	All Other Serotyped	5239	15.6
	Unknown	735	2.2
	Partially serotyped isolates	1351	4.0
	Rough or nonmotile isolates	19	0.1
	Sub Total	7344	21.9
	Total	33589	100
NOTE:			
* Typhimurium includes var. 5- (Formerly var. Copenhagen)			

TABLE 1a

**The Top 20 most frequently reported Salmonella serotypes from
Clinical and Non-Clinical Nonhuman sources reported to CDC and NVSL in 2003**

Clinical Nonhuman 2003			
Rank	Serotype	Reported	Percent
1	S. Typhimurium *	1239	23.1
2	S. Newport	888	16.6
3	S. Agona	240	4.5
4	S. Choleraesuis **	173	3.2
5	S. Derby	157	2.9
6	S. Heidelberg	147	2.7
7	S. Anatum	141	2.6
8	S. Montevideo	141	2.6
9	S. Kentucky	133	2.5
10	S. Senftenberg	107	2.0
11	S. Muenster	99	1.8
12	S. Dublin	92	1.7
13	S. Infantis	87	1.6
14	S. 14,[5],12:i:-	74	1.4
15	S. Oranienburg	67	1.3
16	S. Uganda	66	1.2
17	S. Hadar	62	1.2
18	S. Enteritidis	59	1.1
19	S. Saintpaul	55	1.0
20	S. Braenderup	51	1.0
	Sub Total	4078	76.1
	All Other Serotyped	1225	22.9
	Rough or nonmotile isolates	56	1.0
	Sub Total	1281	23.9
	Total	5359	100

NOTE:

* Typhimurium includes var. 5- (Formerly var. Copenhagen)
** Choleraesuis includes var. Decatur and Kunzendorf

Non-Clinical Nonhuman 2003			
Rank	Serotype	Reported	Percent
1	S. Typhimurium *	806	14.2
2	S. Heidelberg	786	13.8
3	S. Senftenberg	641	11.3
4	S. Hadar	467	8.2
5	S. Kentucky	467	8.2
6	S. Newport	290	5.1
7	S. Mbandaka	195	3.4
8	S. Muenster	183	3.2
9	S. Agona	134	2.4
10	S. Bredeney	129	2.3
11	S. Montevideo	124	2.2
12	S. Braenderup	90	1.6
13	S. Enteritidis	86	1.5
14	S. Saintpaul	63	1.1
15	S. Thompson	63	1.1
16	S. Schwarzengrund	62	1.1
17	S. Anatum	58	1.0
18	S. Worthington	55	1.0
19	S. Derby	51	0.9
20	S. Reading	51	0.9
	Sub Total	4801	84.6
	All Other Serotyped	854	15.1
	Rough or nonmotile isolates	21	0.4
	Sub Total	875	15.4
	Total	5676	100

NOTE:

* Typhimurium includes var. 5- (Formerly var. Copenhagen)