

TABLE 1
THE 20 MOST FREQUENTLY REPORTED SALMONELLA SEROTYPES
FROM HUMAN SOURCES REPORTED TO CDC IN 1998 AND FROM
NONHUMAN SOURCES REPORTED TO CDC AND USDA IN 1998

HUMAN 1998				NONHUMAN 1998			
RANK	SEROTYPE	NUMBER	PERCENT	RANK	SEROTYPE	NUMBER	PERCENT
1	TYPHIMURIUM *	8818	26.0	1	TYPHIMURIUM *	4837	22.9
2	ENTERITIS	6029	17.7	2	HEIDELBERG	2282	10.8
3	NEWPORT	2272	6.7	3	KENTUCKY	1265	6.0
4	HEIDELBERG	1900	5.6	4	DERBY	1061	5.0
5	JAVIANA	1167	3.4	5	SENFTENBERG	1008	4.8
6	AGONIA	991	2.9	6	ENTERITIS	850	4.0
7	MONTEVIDEO	828	2.4	7	MONTEVIDEO	659	3.1
8	ORANIENBURG	693	2.0	8	ANATUM	649	3.1
9	MUENCHEN	639	1.9	9	HADAR	633	3.0
10	INFANTIS	600	1.8	10	AGONIA	625	3.0
11	THOMPSON	571	1.7	11	INFANTIS	469	2.2
12	HADAR	544	1.6	12	SCHWARZENGROUND	416	2.0
13	BRAENDERUP	497	1.5	13	CHOLERAESUIS **	407	1.9
14	SAINTPAUL	479	1.4	14	MUENSTER	381	1.8
15	TYPHI	382	1.1	15	MBANDAKA	379	1.8
16	POONA	346	1.0	16	BREDENEY	349	1.7
17	MISSISSIPPI	314	0.9	17	NEWPORT	297	1.4
18	JAVA	248	0.7	18	SAINTPAUL	289	1.4
19	STANLEY	193	0.6	19	CERRO	278	1.3
20	PARATYPHI B	189	0.6	20	READING	259	1.2
	SUB TOTAL	27700	81.5			17393	82.4
	TOTAL	33971				21097	

* TYPHIMURIUM INCLUDES VAR. COPENHAGEN
** CHOLERAESUIS INCLUDES VAR. KUNZENDORF