

TABLE 1
THE 20 MOST FREQUENTLY REPORTED SALMONELLA SEROTYPES
FROM HUMAN SOURCES REPORTED TO CDC IN 2001 AND FROM
CLINICAL AND NON-CLINICAL NONHUMAN SOURCES REPORTED TO CDC AND NVSL IN 2001

HUMAN 2001				CLINICAL NONHUMAN 2001				NON-CLINICAL NONHUMAN 2001			
RANK	SEROTYPE	NUMBER	PERCENT	RANK	SEROTYPE	NUMBER	PERCENT	RANK	SEROTYPE	NUMBER	PERCENT
1	TYPHIMURIUM *	6999	22.1	1	TYPHIMURIUM *	2043	28.2	1	HEIDELBERG	2455	23.5
2	ENTERITIS	5614	17.7	2	NEWPORT	986	13.6	2	TYPHIMURIUM *	1477	14.1
3	NEWPORT	3158	10.0	3	HEIDELBERG	436	6.0	3	MONTEVERDEO	647	6.2
4	HEIDELBERG	1884	5.9	4	AGONA	363	5.0	4	KENTUCKY	615	5.9
5	JAVIANA	1067	3.4	5	SENFTENBERG	282	3.9	5	SENFTENBERG	597	5.7
6	MONTEVIDEO	626	2.0	6	CHOLERAESUIS **	256	3.5	6	AGONA	410	3.9
7	ORANIENBURG	595	1.9	7	MUENSTER	201	2.8	7	MUENSTER	329	3.1
8	MUENCHEN	583	1.8	8	MONTEVERDEO	191	2.6	8	ENTERITIDS	273	2.6
9	THOMPSON	514	1.6	9	ENTERITIDS	159	2.2	9	DERBY	268	2.6
10	SAINTPAUL	469	1.5	10	DUBLIN	146	2.0	10	HADAR	266	2.5
11	JAVA	466	1.5	11	DERBY	144	2.0	11	ANATUM	218	2.1
12	INFANTIS	440	1.4	12	ANATUM	143	2.0	12	NEWPORT	184	1.8
13	BRAENDERUP	388	1.2	13	INFANTIS	135	1.9	13	MBANDAKA	179	1.7
14	AGONA	370	1.2	14	KENTUCKY	109	1.5	14	INFANTIS	175	1.7
15	TYPHI	343	1.1	15	UGANDA	89	1.2	15	SCHWARZENGRUND	156	1.5
16	MISSISSIPPI	336	1.1	16	BREDENEY	81	1.1	16	ORANIENBURG	142	1.4
17	BERTA	330	1.0	17	MUENCHEN	68	0.9	17	BERTA	139	1.3
18	POONA	330	1.0	18	HADAR	64	0.9	18	WORTHINGTON	117	1.1
19	HADAR	307	1.0	19	MBANDAKA	64	0.9	19	CERRO	109	1.0
20	BAREILLY	205	0.6	20	CERRO	63	0.9	20	LIVINGSTONE	102	1.0
SUB TOTAL		25024	79.0	SUB TOTAL		6023	83.2	SUB TOTAL		8858	84.8
TOTAL		31675		TOTAL		7243		TOTAL		10451	

* TYPHIMURIUM INCLUDES VAR. COPENHAGEN
** CHOLERAESUIS INCLUDES VAR. KUNZENDORF