

**TABLE 1**

**The Top 20 most frequently reported Salmonella serotypes from Human sources reported to CDC in 2004**

Human 2004			
Rank	Serotype	Reported	Percent
1	S. Typhimurium *	6842	19.2
2	S. Enteritidis	5012	14.1
3	S. Newport	3325	9.3
4	S. Javiana	1772	5.0
5	S. Heidelberg	1757	4.9
6	S. Montevideo	870	2.4
7	S. I 4,[5],12:i:-	739	2.1
8	S. Muenchen	739	2.1
9	S. Saintpaul	692	1.9
10	S. Braenderup	684	1.9
11	S. Infantis	588	1.6
12	S. Mississippi	558	1.6
13	S. Oranienburg	495	1.4
14	S. Thompson	493	1.4
15	S. Berta	409	1.1
16	S. Agona	406	1.1
17	S. Paratyphi B var. L(+) tartrate+	354	1.0
18	S. Typhi	306	0.9
19	S. Hadar	277	0.8
20	S. Anatum	250	0.7
	<b>Sub Total</b>	<b>26568</b>	<b>74.5</b>
	All Other Serotyped	5651	15.8
	Unknown	2053	5.8
	Partially serotyped isolates	1328	3.7
	Rough or nonmotile isolates	61	0.2
	<b>Sub Total</b>	<b>9093</b>	<b>25.5</b>
	<b>Total</b>	<b>35661</b>	<b>100</b>
NOTE:			
* Typhimurium includes var. 5- (Formerly var. Copenhagen)			

**TABLE 1a**

**The Top 20 most frequently reported Salmonella serotypes from  
Clinical and Non-Clinical Nonhuman sources reported to CDC and NVSL in 2004**

Clinical Nonhuman 2004			
Rank	Serotype	Reported	Percent
1	S. Typhimurium *	1558	23.7
2	S. Newport	1067	16.3
3	S. Agona	267	4.1
4	S. Heidelberg	215	3.3
5	S. Choleraesuis **	213	3.2
6	S. Derby	207	3.2
7	S. I 4,[5],12:i:-	154	2.3
8	S. Infantis	148	2.3
9	S. Dublin	139	2.1
10	S. Senftenberg	138	2.1
11	S. Montevideo	136	2.1
12	S. Uganda	134	2.0
13	S. Muenster	128	2.0
14	S. Anatum	126	1.9
15	S. Reading	91	1.4
16	S. Kentucky	89	1.4
17	S. Muenchen	89	1.4
18	S. Enteritidis	63	1.0
19	S. Mbandaka	53	0.8
20	S. Cerro	52	0.8
	<b>Sub Total</b>	<b>5067</b>	<b>77.2</b>
	All Other Serotyped	1413	21.5
	Rough or nonmotile isolates	82	1.2
	<b>Sub Total</b>	<b>1495</b>	<b>22.8</b>
	<b>Total</b>	<b>6562</b>	<b>100</b>

NOTE:

\* Typhimurium includes var. 5- (Formerly var. Copenhagen)  
\*\* Choleraesuis includes var. Decatur and Kunzendorf

Non-Clinical Nonhuman 2004			
Rank	Serotype	Reported	Percent
1	S. Typhimurium *	672	13.0
2	S. Heidelberg	526	10.2
3	S. Senftenberg	508	9.9
4	S. Hadar	413	8.0
5	S. Kentucky	292	5.7
6	S. Derby	211	4.1
7	S. Newport	187	3.6
8	S. Enteritidis	175	3.4
9	S. Agona	170	3.3
10	S. Montevideo	142	2.8
11	S. Muenster	124	2.4
12	S. Cerro	114	2.2
13	S. Mbandaka	110	2.1
14	S. Schwarzengrund	110	2.1
15	S. Worthington	97	1.9
16	S. Reading	78	1.5
17	S. Inverness	74	1.4
18	S. I 4,[5],12:i:-	60	1.2
19	S. Anatum	53	1.0
20	S. Dublin	52	1.0
	<b>Sub Total</b>	<b>4168</b>	<b>80.9</b>
	All Other Serotyped	957	18.6
	Rough or nonmotile isolates	25	0.5
	<b>Sub Total</b>	<b>982</b>	<b>19.1</b>
	<b>Total</b>	<b>5150</b>	<b>100</b>

NOTE:

\* Typhimurium includes var. 5- (Formerly var. Copenhagen)