Disease Burden from Hepatitis A, B, and C in the United States

	Hepatitis A							
	2006	2005	2004	2003	2002	2001		
Number of Acute Clinical Cases Reported ¹	3,579	4,488	5,683	7,653	8,795	10,616		
Estimated Number of Acute Clinical Cases ²	15,000 19,000		24,000	33,000	38,000	45,000		
Estimated Number of New Infections ² (current)	32,000 42,000		56,000	61,000	73,000	93,000		
Estimated Number of New Infections ² (historical)			mean	min	max			
	1990-1999		301,000	181,000	373,000			
	1980-1989		254,000	221,000	380,000			
Number of Persons with Chronic Infection ³	no chronic infection							
Estimated Annual Number of Chronic Liver Disease Deaths †	no chronic infection							
Percent Ever Infected ‡	33%							

	Hepatitis B						
	2006	2006 2005 2004 2003 2002					
Number of Acute Clinical Cases Reported ¹	4,758	5,494	6,212	7,526	8,064	7,844	
Estimated Number of Acute Clinical Cases ²	13,000	15,000	17,000	22,000	no data	no data	
Estimated Number of New Infections ² (current)	46,000	51,000	60,000	73,000	79,000	78,000	

Estimated Number of New Infections ² (historical)		mean	min	max
	1990-1999	140,000	79,000	232,000
	1980-1989	259,000	208,000	287,000

Number of Persons with Chronic Infection ³	1.25 million persons
Estimated Annual Number of Chronic Liver Disease Deaths †	5,000
Percent Ever Infected ‡	4.9%

	Hepatitis C						
	2006	2005	2004	2003	2002	2001	
Number of Acute Clinical Cases Reported ¹		no data	no data		no data		
Estimated Number of Acute Clinical Cases ²	3,200	3,400	4,200	4,500	4,800	3,900	
Estimated Number of New Infections ² (current)	19,000	21,000	26,000	28,000	29,000	24,000	
Estimated Number of New Infections ² (historical)			mean	min	max		
	1990-1999 1982-1989		67,000	36,000	179,000		
			232,000	180,000	291,000		
Number of Persons with Chronic Infection ³	3.2 million persons						
Estimated Annual Number of Chronic Liver Disease Deaths †	8,000-10,000						
Percent Ever Infected ‡	1.6%						

¹ Number of Acute Clinical Cases Reported: For hepatitis A and hepatitis B, the number of cases reported to the National Notifiable Disease Surveillance System(NNDSS). Cases of hepatitis C/NANB are also reported to NNDSS (758 in 2004) but are unreliable for monitoring trends in hepatitis C because these reports include cases based only on a positive lab test for anti-HCV, most of which represent chronic HCV infection. (CDC. Summary of Notifiable Diseases, United States, 2002, MMWR 2003;51(53).


² Estimated Number of Acute Clinical Cases and New Infections: Incidence estimates are derived from catalytic modeling of seroprevalence data from the Third National Health and Nutrition Examination Survey (NHANES III) applied to cases reported to the Nationally Notifiable Disease Surveillance System (NNDSS) (for hepatitis A and hepatitis B) or to the Sentinel Counties Study of Viral Hepatitis (for hepatitis C).

³ Number of Persons with Chronic Infection: HBV: Margolis HS, Coleman PJ, Brown RE, et al. Prevention of hepatitis B virus transmission by immunization: An economic analysis of current recommendations. JAMA 1995; 274(15): 1201 - 1208. HCV: Armstrong GL et al. Prevalence of hepatitis C virus infection in the United States, 1999 through 2002. Ann Int Med 2006 (in press).


† Estimated Annual Number of Chronic Liver Disease Deaths: HBV: Margolis HS, Coleman PJ, Brown RE, et al. Prevention of hepatitis B virus transmission by immunization: An economic analysis of current recommendations. JAMA 1995; 274(15): 1201 - 1208.HCV: Centers for Disease Control and Prevention. Recommendations for the prevention and control of hepatitis C virus infection and HCV-related chronic disease. MMWR 1998; 47(RR-19):1-39.

‡ Percent Ever Infected: Seroprevalence estimates for HAV, HBV and HCV come from the National Health and Nutrition Examination Survey HAV: Bell BP, Kruszon-Moran D, Shapiro CN et al. Hepatitis A virus infection in the United States: Serologic results from the Third National Health and Nutrition Examination Survey. Vaccine 23(2005):5798-5806. HBV: McQuillan GM, et al. Prevalence of hepatitis B virus infection in the United States: The National Health and Nutrition and Examination Surveys, 1976 through 1994. AJPH 1999; 89(1)14-18. HCV: Armstrong GL et al. Prevalence of hepatitis C virus infection in the United States, 1999 through 2002. Ann Int Med 2006; 144:705-71.

Acute Disease Burden Data 1980 - 2006



		Year	Reported Acute Cases	Estimated Acute Cases	Estimated Total New Infections
		1980	19,014	53,000	208,000
		1981	21,151	59,000	229,000
		1982	22,176	62,000	239,000
		1983	24,319	68,000	267,000
	Incidence of acute hepatitis B,	1984	26,116	73,000	281,000
;	United States	1985	26,612	74,000	287,000
	\sim	1986	26,106	73,000	283,000
	70000 - Reported Acute Cases	1987	25,915	72,000	287,000
	BOODO - Estimated Acute Cases	1988	23,175	65,000	253,000
20		1989	23,421	65,000	255,000
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	50000 -	1990	21,277	59,000	232,000
Number of cases	10000 -	1991	17,911	50,000	193,000
		1992	16,126	45,000	175,000
	30000 -	1993	13,361	37,000	144,000
1	20000	1994	12,517	35,000	133,000
		1995	10,805	30,000	113,000
	10000 -	1996	10,637	30,000	112,000
	0 +	1997	10,416	29,000	110,000
	2006 2004 2004 1998 1996 1986 1986 1986 1986	1998	10,258	29,000	109,000
	ਕੇ ਕੇ 6 ਕੇ ਕੇ 6 ਕੇ 7 6 6 ਕੇ 7 6	1999	7,694	21,000	79,000
		2000	8,036	22,000	81,000
		2001	7,844	22,000	78,000
		2002	8,064	23,000	79,000
		2003	7,526	21,000	73,000
		2004	6,212	17,000	60,000
		2005	5,494	15,000	51,000
		2006	4,758	13,000	46,000



2003

2004

2005

2006

4,500

4,200

3,400

3,200

28,000

26,000

21,000

19,000