

**Table 35. Tuberculosis Cases and Percentages by Initial Drug Regimen: 59 Reporting Areas, 2003**

Reporting Area	Total Cases	Cases in Persons Alive at Diagnosis	Cases with Information on Initial Drug Regimen		Percentage of Cases in Persons with Initial Drug Regimen <sup>1,2</sup>		
			No.	(%)	IR	IRZ	IRZ,E/S
<b>United States</b>	<b>14,874</b>	<b>14,534</b>	<b>14,476</b>	<b>(99.6)</b>	<b>(1.4)</b>	<b>(8.1)</b>	<b>(81.1)</b>
Alabama	258	254	254	(100.0)	(0.8)	(58.7)	(35.8)
Alaska	57	55	55	(100.0)	(0.0)	(10.9)	(85.5)
Arizona	295	284	280	(98.6)	(1.1)	(5.0)	(90.0)
Arkansas	127	124	122	(98.4)	(7.4)	(51.6)	(35.2)
California	3,227	3,177	3,172	(99.8)	(1.5)	(4.2)	(86.0)
Colorado	111	110	110	(100.0)	(0.9)	(4.5)	(82.7)
Connecticut	111	108	104	(96.3)	(4.8)	(4.8)	(82.7)
Delaware	33	31	31	(100.0)	(3.2)	(3.2)	(93.5)
District of Columbia	79	79	79	(100.0)	(0.0)	(2.5)	(65.8)
Florida	1,046	1,016	1,016	(100.0)	(0.6)	(6.8)	(81.7)
Georgia	526	515	513	(99.6)	(0.8)	(8.8)	(82.3)
Hawaii	117	116	116	(100.0)	(4.3)	(12.1)	(75.9)
Idaho	13	13	13	(100.0)	(0.0)	(23.1)	(76.9)
Illinois	633	624	617	(98.9)	(0.6)	(7.8)	(79.4)
Indiana	143	142	142	(100.0)	(0.0)	(7.7)	(85.2)
Iowa	40	38	37	(97.4)	(2.7)	(13.5)	(81.1)
Kansas	75	73	73	(100.0)	(2.7)	(5.5)	(79.5)
Kentucky	138	133	133	(100.0)	(1.5)	(12.0)	(77.4)
Louisiana	260	252	249	(98.8)	(2.8)	(7.2)	(82.7)
Maine	25	24	24	(100.0)	(4.2)	(16.7)	(75.0)
Maryland	268	262	262	(100.0)	(0.4)	(3.4)	(91.6)
Massachusetts	261	258	257	(99.6)	(1.2)	(5.1)	(83.3)
Michigan	243	233	233	(100.0)	(3.4)	(21.9)	(66.5)
Minnesota	214	213	213	(100.0)	(0.5)	(7.5)	(83.6)
Mississippi	128	124	121	(97.6)	(0.0)	(9.9)	(83.5)
Missouri	131	126	126	(100.0)	(1.6)	(7.1)	(79.4)
Montana	7	7	7	(100.0)	(0.0)	(28.6)	(57.1)
Nebraska	28	27	27	(100.0)	(0.0)	(14.8)	(63.0)
Nevada	107	102	101	(99.0)	(0.0)	(1.0)	(95.0)
New Hampshire	15	14	14	(100.0)	(0.0)	(0.0)	(92.9)
New Jersey	495	485	484	(99.8)	(1.4)	(9.1)	(79.1)
New Mexico	49	47	47	(100.0)	(0.0)	(27.7)	(68.1)
New York State <sup>3</sup>	340	332	332	(100.0)	(0.3)	(3.9)	(87.7)
New York City	1,140	1,111	1,111	(100.0)	(0.7)	(5.1)	(84.2)
North Carolina	374	365	365	(100.0)	(0.3)	(4.7)	(83.3)
North Dakota	6	5	5	(100.0)	(0.0)	(0.0)	(100.0)
Ohio	229	220	220	(100.0)	(1.8)	(14.5)	(75.5)
Oklahoma	163	160	160	(100.0)	(3.8)	(11.3)	(60.6)
Oregon	106	103	103	(100.0)	(1.0)	(2.9)	(90.3)
Pennsylvania	336	327	327	(100.0)	(1.8)	(9.2)	(57.5)
Rhode Island	46	45	45	(100.0)	(0.0)	(4.4)	(80.0)
South Carolina	254	242	242	(100.0)	(0.8)	(9.9)	(78.5)
South Dakota	20	19	19	(100.0)	(0.0)	(21.1)	(78.9)
Tennessee	285	271	271	(100.0)	(1.5)	(10.3)	(82.7)
Texas	1,594	1,557	1,538	(98.8)	(2.4)	(7.3)	(80.8)
Utah	39	37	37	(100.0)	(0.0)	(8.1)	(83.8)
Vermont	9	9	9	(100.0)	(0.0)	(33.3)	(66.7)
Virginia	332	328	327	(99.7)	(0.6)	(6.1)	(89.9)
Washington	250	247	243	(98.4)	(0.4)	(2.9)	(90.9)
West Virginia	21	21	21	(100.0)	(4.8)	(14.3)	(52.4)
Wisconsin	66	65	65	(100.0)	(1.5)	(7.7)	(83.1)
Wyoming	4	4	4	(100.0)	(0.0)	(0.0)	(75.0)
American Samoa <sup>4</sup>	...	...	...	...	...	...	...
Fed. States of Micronesia <sup>4</sup>	...	...	...	...	...	...	...
Guam <sup>4</sup>	61	61	60	(98.4)	(0.0)	(0.0)	(90.0)
N. Mariana Islands <sup>4</sup>	45	45	45	(100.0)	(0.0)	(0.0)	(97.8)
Puerto Rico <sup>4</sup>	115	104	104	(100.0)	(1.0)	(0.0)	(91.3)
Republic of Palau <sup>4</sup>	9	9	9	(100.0)	(0.0)	(11.1)	(77.8)
U.S. Virgin Islands <sup>4</sup>	...	...	...	...	...	...	...

<sup>1</sup>Percentage based on 52 reporting areas (50 states, New York City, and the District of Columbia). Counts and percentages shown only for reporting areas with information reported for  $\geq 75\%$  of cases.

<sup>2</sup>I=isoniazid; R=rifampin; Z=pyrazinamide; E=ethambutol; S=streptomycin.

<sup>3</sup>Excludes New York City.

<sup>4</sup>Not included in U.S. totals.

**Note:** Excluding cases with no information on drug regimen, 159 (0.9%) persons were not started on any drugs, 27 (0.2%) were started on one drug, and 1,463 (8.6%) had an initial multidrug regimen other than IR, IRZ, or IRZ,E/S.

Ellipses indicate data not available.