

Table 10. Tuberculosis Cases and Percentages, by Monodrug (INH) or Multidrug Resistance in Persons with No Previous TB, by Origin: United States, 1993-2003

Year	Resistance to Isoniazid ¹						Resistance to Isoniazid and Rifampin ¹					
	Total Cases ²		U.S.-born		Foreign-born ³		Total Cases ²		U.S.-born		Foreign-born ³	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
1993	1,400	8.4	804	6.8	580	12.4	410	2.5	302	2.6	105	2.3
1994	1,353	8.3	708	6.4	631	12.1	352	2.2	238	2.2	109	2.1
1995	1,173	7.3	554	5.4	618	11.0	253	1.6	168	1.6	85	1.5
1996	1,136	7.4	494	5.2	639	11.3	206	1.3	104	1.1	101	1.8
1997	1,079	7.5	436	5.0	639	11.2	155	1.1	76	0.9	79	1.4
1998	1,011	7.5	366	4.7	643	11.3	130	1.0	54	0.7	75	1.3
1999	899	7.1	283	4.0	614	11.0	127	1.0	39	0.6	88	1.6
2000	894	7.6	269	4.4	622	11.1	121	1.0	38	0.6	83	1.5
2001	797	7.0	241	4.4	555	9.5	113	1.0	34	0.6	79	1.4
2002	801	7.6	200	4.1	598	10.7	124	1.2	35	0.7	89	1.6
2003	784	7.9	204	4.6	577	10.6	90	0.9	25	0.6	65	1.2

¹Isolates may be resistant to other drugs.

²Includes persons of unknown country of birth.

³Includes persons born outside the United States, American Samoa, the Federated States of Micronesia, Guam, the Republic of the Marshall Islands, Midway Island, the Commonwealth of the Northern Mariana Islands, Puerto Rico, the Republic of Palau, the U.S. Virgin Islands, and U.S. minor and outlying Pacific islands.

Note: Data for all years updated through March 30, 2004.

More than 85% of all persons in each group had drug susceptibility test results reported for an initial isolate.

See Surveillance Slides #19 through #22.

Table 11. Tuberculosis Cases and Percentages, by Monodrug (INH) or Multidrug Resistance in Persons with Previous TB, by Origin: United States, 1993-2003

Year	Resistance to Isoniazid ¹						Resistance to Isoniazid and Rifampin ¹					
	Total Cases ²		U.S.-born		Foreign-born ³		Total Cases ²		U.S.-born		Foreign-born ³	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
1993	164	16.6	85	12.7	76	25.0	75	7.7	30	4.5	45	15.0
1994	177	17.1	81	11.7	95	28.1	75	7.3	35	5.1	39	11.6
1995	168	17.6	77	13.0	91	25.1	70	7.3	28	4.7	42	11.6
1996	142	16.5	67	12.1	74	24.4	43	5.0	20	3.6	22	7.3
1997	109	14.7	35	7.7	74	25.9	44	5.9	12	2.6	32	11.2
1998	98	13.1	38	7.8	60	22.8	23	3.1	6	1.2	17	6.5
1999	82	12.3	25	6.5	55	19.4	28	4.2	6	1.6	22	7.8
2000	82	13.0	22	6.1	60	22.2	24	3.8	2	0.6	22	8.2
2001	82	13.2	27	8.4	55	18.5	29	4.7	7	2.2	22	7.4
2002	77	14.1	23	7.9	54	21.3	24	4.4	3	1.0	21	8.3
2003	63	12.6	17	7.1	46	17.6	18	3.6	3	1.3	15	5.8

¹Isolates may be resistant to other drugs.

²Includes persons of unknown country of birth.

³Includes persons born outside the United States, American Samoa, the Federated States of Micronesia, Guam, the Republic of the Marshall Islands, Midway Island, the Commonwealth of the Northern Mariana Islands, Puerto Rico, the Republic of Palau, the U.S. Virgin Islands, and U.S. minor and outlying Pacific islands.

Note: Data for all years updated through March 30, 2004.

More than 85% of all persons in each group had drug susceptibility test results reported for an initial isolate.