

**Table 9. Percentage of Reported TB Cases by Initial Drug Regimen, Use of Directly Observed Therapy, and Completion of Therapy (COT): United States, 1993-2000**

Year	Initial Drug Regimen <sup>1,2</sup>			Directly Observed Therapy <sup>3</sup>		Therapy ≤1 Year Indicated <sup>4</sup>	
	IR	IRZ	IRZ,E/S	DOT Only	Both DOT and Self-Administered	COT ≤1 Year	COT
1993	13.0	31.2	40.9	21.7	14.4	63.6	87.5
1994	7.0	23.3	56.3	28.1	20.5	68.5	87.9
1995	5.2	20.3	63.3	37.2	21.5	72.9	89.6
1996	4.2	17.5	67.9	42.5	22.4	75.5	90.2
1997	3.2	15.1	72.4	46.9	23.8	77.7	91.1
1998	2.6	12.9	74.7	51.7	22.4	79.1	90.7
1999	2.2	11.2	77.2	...	...	...	...
2000	1.9	10.3	78.2	...	...	...	...

1. Includes cases in persons alive at diagnosis.

2. I=isoniazid; R=rifampin; Z=pyrazinamide; E=ethambutol; S=streptomycin. Excluding cases with no information on initial drug regimen, 1% were not started on any drugs, less than 1% were started on one drug, and approximately 10% had an initial multidrug regimen other than IR, IRZ, and IRZ,E/S.

3. Includes cases in persons alive at diagnosis with initial drug regimen of one or more drugs prescribed.

4. Includes cases in persons alive at diagnosis, with initial drug regimen of one or more drugs prescribed, who did not die during therapy. Excludes persons with initial isolate resistant to rifampin and pediatric (aged <15) cases with meningeal, bone or joint, or miliary disease. See Technical Notes (Appendix A) for description of COT calculation.

Ellipses indicate data not available.

Note: Data for all years updated through April 17, 2001.

See Surveillance Slides #17 and #18.

**Table 10. Number and Percentage of Reported TB Cases with HIV Test Results and with HIC Coinfection by Age Group: United States, 1993-1999**

Year	25-44 Years Old				All Ages			
	HIV Test Results <sup>1</sup>		HIV Positive <sup>2</sup>		HIV Test Results <sup>1</sup>		HIV Positive <sup>2</sup>	
	No.	%	No.	%	No.	%	No.	%
1993	4,375	46	2,788	29	7,451	30	3,680	15
1994	4,433	49	2,660	29	7,871	33	3,590	15
1995	4,269	52	2,169	26	8,173	36	3,035	13
1996	4,347	57	1,857	25	8,800	41	2,617	12
1997	4,134	60	1,469	21	8,756	44	2,088	11
1998	3,856	61	1,238	20	8,274	45	1,828	10
1999	3,785	62	1,171	19	8,349	48	1,721	10

1. Rhode Island reported HIV test results in 1998 and 1999. HIV test results were not reported from California. California did provide HIV status for TB cases reported during 1993-1999 in persons also reported with AIDS (i.e., HIV-positive). Includes cases with positive, negative, or indeterminate HIV test results and California cases also reported with AIDS. Percentages based on all reported TB cases.

2. Includes cases with HIV-positive test results and California cases also reported with AIDS. Percentages based on all reported TB cases.

Note: Data for all years updated through April 17, 2000.

See Surveillance Slide #16.