

Table 41. Tuberculosis Cases and Percentages by Completion of Tuberculosis Therapy (COT): 60 Reporting Areas, 2002

Reporting Area	Total Cases	Therapy ≤1 Year Indicated ¹		Therapy >1 Year Indicated ²		All Drug Therapy		
		No. ³	COT ≤1 Year(%)	COT(%)	No. ³	COT(%)	No. ³	COT(%)
United States	15,057	13,276	(80.9)	(91.6)	216	(77.8)	13,492	(91.4)
Alabama	233	203	(88.7)	(97.0)	2	(100.0)	205	(97.1)
Alaska	49	45	(73.3)	(100.0)	1	(100.0)	46	(100.0)
Arizona	263	225	(80.4)	(90.2)	2	(50.0)	227	(89.9)
Arkansas	136	114	(86.8)	(91.2)	2	(100.0)	116	(91.4)
California	3,173	2,822	(81.2)	(92.0)	52	(75.0)	2,874	(91.7)
Colorado	102	95	(85.3)	(90.5)	1	(100.0)	96	(90.6)
Connecticut	105	90	(70.0)	(88.9)	2	(100.0)	92	(89.1)
Delaware	25	23	(91.3)	(100.0)	0	...	23	(100.0)
District of Columbia	82	77	(97.4)	(100.0)	0	...	77	(100.0)
Florida	1,084	945	(81.5)	(92.7)	16	(93.8)	961	(92.7)
Georgia	535	464	(76.7)	(89.4)	3	(100.0)	467	(89.5)
Hawaii	148	137	(69.3)	(87.6)	2	(100.0)	139	(87.8)
Idaho	14	12	—	—	0	...	12	—
Illinois	680	583	(76.2)	(88.2)	7	(57.1)	590	(87.8)
Indiana	128	107	(96.3)	(97.2)	0	...	107	(97.2)
Iowa	34	33	(81.8)	(97.0)	0	...	33	(97.0)
Kansas	86	76	(80.3)	(86.8)	2	(0.0)	78	(84.6)
Kentucky	146	129	(85.3)	(93.8)	1	(100.0)	130	(93.8)
Louisiana	230	210	—	—	1	—	211	—
Maine	23	18	(77.8)	(83.3)	1	(100.0)	19	(84.2)
Maryland	305	265	(86.4)	(94.7)	5	(100.0)	270	(94.8)
Massachusetts	271	255	(76.9)	(90.6)	2	(100.0)	257	(90.7)
Michigan	315	263	(81.4)	(93.5)	2	(50.0)	265	(93.2)
Minnesota	237	220	(85.5)	(95.0)	8	(87.5)	228	(94.7)
Mississippi	134	113	(83.2)	(97.3)	1	(100.0)	114	(97.4)
Missouri	136	115	(85.2)	(93.9)	0	...	115	(93.9)
Montana	12	10	(90.0)	(90.0)	0	...	10	(90.0)
Nebraska	28	24	(66.7)	(87.5)	0	...	24	(87.5)
Nevada	85	71	(97.2)	(98.6)	2	(50.0)	73	(97.3)
New Hampshire	19	18	(88.9)	(94.4)	0	...	18	(94.4)
New Jersey	530	475	(77.9)	(88.2)	7	(85.7)	482	(88.2)
New Mexico	57	50	(78.0)	(100.0)	1	(100.0)	51	(100.0)
New York State ⁴	347	318	(76.4)	(88.7)	5	(80.0)	323	(88.5)
New York City	1,071	927	(85.5)	(94.9)	28	(78.6)	955	(94.5)
North Carolina	434	369	(90.5)	(96.5)	6	(83.3)	375	(96.3)
North Dakota	6	6	(100.0)	(100.0)	0	...	6	(100.0)
Ohio	257	219	(80.4)	(93.2)	7	(85.7)	226	(92.9)
Oklahoma	190	169	(84.6)	(91.1)	3	(66.7)	172	(90.7)
Oregon	111	104	(86.5)	(94.2)	2	(50.0)	106	(93.4)
Pennsylvania	353	306	(73.9)	(92.2)	3	(33.3)	309	(91.6)
Rhode Island	49	45	(68.9)	(95.6)	0	...	45	(95.6)
South Carolina	256	222	(80.2)	(91.0)	2	(50.0)	224	(90.6)
South Dakota	13	9	(88.9)	(100.0)	0	...	9	(100.0)
Tennessee	308	276	(83.3)	(96.4)	1	(100.0)	277	(96.4)
Texas	1,540	1,362	(75.7)	(86.4)	26	(76.9)	1,388	(86.2)
Utah	31	29	(96.6)	(100.0)	1	(100.0)	30	(100.0)
Vermont	8	8	(87.5)	(87.5)	0	...	8	(87.5)
Virginia	315	289	(86.5)	(92.7)	4	(50.0)	293	(92.2)
Washington	252	235	(83.4)	(94.5)	4	(75.0)	239	(94.1)
West Virginia	30	26	(65.4)	(88.5)	0	...	26	(88.5)
Wisconsin	78	67	(82.1)	(94.0)	1	(100.0)	68	(94.1)
Wyoming	3	3	(100.0)	(100.0)	0	...	3	(100.0)
American Samoa ⁵
Fed. States of Micronesia ⁵
Guam ⁵	65	53	—	—	0	...	53	—
Marshall Islands ⁵
N. Mariana Islands ⁵
Puerto Rico ⁵	129	93	(79.6)	(86.0)	3	(66.7)	96	(85.4)
Republic of Palau ⁵
U.S. Virgin Islands ⁵

¹Initial isolate susceptible to rifampin (n=9,889) or susceptibility unknown (n=415); culture negative (n=2,180); culture status unknown (n=789); age unknown (n=4).

²Initial isolate rifampin resistant, or pediatric (aged <15) patients with meningeal, bone, joint, or miliary disease.

³Number of cases in persons alive at diagnosis, with an initial regimen of one or more drugs prescribed, who did not die during therapy. Percentage for U.S. based on 52 reporting areas (50 states, New York City, and the District of Columbia). Percentages shown only for reporting areas with information reported for ≥ 90% of cases.

⁴Excludes New York City.

⁵Not included in U.S. totals.

Note: Ellipses indicate data not available. See Technical Notes for description of COT calculation (page 9).