

Table 44. Tuberculosis Cases and Percentages in Persons Completing Therapy for Whom Therapy Was Indicated for One Year or Less: 60 Reporting Areas, 1998–2002¹

Reporting Area	Year									
	1998		1999		2000		2001		2002	
	No. ²	(%) ³	No. ²	(%) ³	No. ²	(%) ³	No. ²	(%) ³	No. ²	(%) ³
United States	15,782	(80.2)	15,207	(79.9)	14,258	(80.8)	14,057	(80.9)	13,276	(80.9)
Alabama	330	(83.9)	260	(80.4)	263	(88.2)	224	(79.5)	203	(88.7)
Alaska	54	(87.0)	59	(88.1)	104	(83.7)	50	(80.0)	45	(73.3)
Arizona	211	(84.4)	231	(82.3)	228	(78.1)	254	(81.5)	225	(79.1)
Arkansas	140	(85.0)	156	(82.1)	176	(84.1)	138	(91.3)	114	(86.8)
California	3,400	(79.3)	3,183	(79.8)	2,931	(79.5)	2,968	(80.1)	2,822	(81.1)
Colorado	64	(96.9)	80	(88.8)	83	(92.8)	128	(93.0)	95	(85.3)
Connecticut	109	(82.6)	108	(78.7)	90	(68.9)	113	(70.8)	90	(70.0)
Delaware	29	(75.9)	30	(80.0)	23	(78.3)	30	(93.3)	23	(91.3)
District of Columbia	90	(73.3)	57	(84.2)	71	(80.3)	63	(85.7)	77	(97.4)
Florida	1,096	(81.6)	1,075	(84.0)	1,004	(82.7)	1,018	(80.7)	945	(81.5)
Georgia	555	(79.5)	594	(77.4)	599	(80.6)	507	(81.3)	464	(76.7)
Hawaii	155	(83.9)	168	(67.3)	119	(73.1)	138	(73.9)	137	(69.3)
Idaho	14	(42.9)	14	(71.4)	13	(53.8)	7	(71.4)	12	(58.3)
Illinois	722	(78.1)	700	(80.9)	636	(83.0)	637	(77.1)	583	(76.2)
Indiana	154	(83.1)	133	(79.7)	119	(89.9)	97	(92.8)	107	(96.3)
Iowa	46	(73.9)	54	(83.3)	38	(89.5)	36	(77.8)	33	(81.8)
Kansas	49	(93.9)	62	(80.6)	70	(82.9)	70	(72.9)	76	(80.3)
Kentucky	145	(82.8)	177	(80.2)	117	(88.0)	131	(87.8)	129	(85.3)
Louisiana	320	(75.3)	310	(77.4)	285	(77.5)	259	(74.5)	210	(72.4)
Maine	10	(70.0)	19	(94.7)	22	(72.7)	19	(94.7)	18	(77.8)
Maryland	278	(84.9)	256	(86.7)	254	(83.1)	234	(89.3)	265	(86.4)
Massachusetts	252	(83.3)	248	(80.6)	261	(83.9)	250	(77.2)	255	(76.9)
Michigan	323	(83.0)	293	(83.6)	248	(82.7)	275	(81.1)	263	(81.4)
Minnesota	146	(82.9)	184	(87.5)	172	(84.9)	221	(84.6)	220	(85.5)
Mississippi	198	(90.4)	187	(84.5)	153	(83.7)	137	(83.9)	113	(83.2)
Missouri	150	(78.0)	174	(79.3)	183	(81.4)	138	(84.8)	115	(85.2)
Montana	14	(100.0)	10	(90.0)	19	(89.5)	17	(94.1)	10	(90.0)
Nebraska	26	(69.2)	18	(55.6)	22	(63.6)	31	(83.9)	24	(66.7)
Nevada	114	(83.3)	81	(77.8)	91	(84.6)	87	(80.5)	71	(97.2)
New Hampshire	13	(100.0)	17	(94.1)	20	(75.0)	19	(89.5)	18	(88.9)
New Jersey	546	(74.5)	490	(71.8)	480	(74.2)	458	(75.5)	475	(77.9)
New Mexico	58	(87.9)	48	(87.5)	38	(71.1)	46	(84.8)	50	(78.0)
New York State ⁴	382	(70.9)	332	(75.9)	358	(79.6)	375	(81.6)	318	(76.4)
New York City	1,312	(81.6)	1,252	(78.8)	1,122	(83.9)	1,065	(86.7)	927	(85.5)
North Carolina	424	(92.2)	431	(88.6)	391	(90.3)	340	(88.5)	369	(90.5)
North Dakota	8	(62.5)	7	(57.1)	5	(100.0)	5	(100.0)	6	(100.0)
Ohio	194	(77.3)	262	(71.4)	298	(74.2)	258	(79.1)	219	(80.4)
Oklahoma	167	(78.4)	180	(73.9)	128	(82.8)	159	(80.5)	169	(84.6)
Oregon	138	(78.3)	114	(78.1)	110	(80.9)	113	(87.6)	104	(86.5)
Pennsylvania	379	(72.3)	370	(76.2)	337	(74.2)	280	(76.4)	306	(73.9)
Rhode Island	58	(82.8)	44	(68.2)	43	(74.4)	60	(73.3)	45	(68.9)
South Carolina	240	(74.2)	264	(69.7)	245	(76.3)	227	(85.0)	222	(80.2)
South Dakota	13	(92.3)	17	(76.5)	12	(75.0)	12	(66.7)	9	(88.9)
Tennessee	371	(82.2)	319	(84.0)	322	(80.4)	265	(84.2)	276	(83.3)
Texas	1,569	(78.1)	1,441	(79.8)	1,323	(78.9)	1,442	(75.7)	1,362	(75.6)
Utah	48	(75.0)	34	(73.5)	41	(73.2)	33	(81.8)	29	(96.6)
Vermont	4	(100.0)	2	(50.0)	2	(100.0)	7	(100.0)	8	(87.5)
Virginia	304	(83.6)	292	(85.6)	254	(86.2)	272	(80.9)	289	(86.5)
Washington	236	(84.3)	235	(81.3)	233	(82.0)	241	(84.6)	235	(83.4)
West Virginia	30	(66.7)	36	(69.4)	24	(50.0)	25	(64.0)	26	(65.4)
Wisconsin	90	(82.2)	96	(86.5)	75	(78.7)	75	(81.3)	67	(82.1)
Wyoming	4	(100.0)	3	(100.0)	3	(100.0)	3	(100.0)	3	(100.0)
American Samoa ⁵
Fed. States of Micronesia ⁵
Guam ⁵	83	(88.0)	65	(53.8)	47	(66.0)	56	(37.5)	53	(37.7)
Marshall Islands ⁵
N. Mariana Islands ⁵	107	(79.4)	62	(82.3)	72	(83.3)	50	(88.0)	51	(68.6)
Puerto Rico ⁵	149	(81.2)	148	(82.4)	128	(87.5)	92	(92.4)	93	(79.6)
Republic of Palau ⁵
U.S. Virgin Islands ⁵

¹Most recent year for which data are available.

²Total cases for which therapy less than 1 year indicated in persons alive at diagnosis, with an initial regimen of one or more drugs prescribed, who did not die during therapy.

³Percentage of total cases in persons who completed therapy for whom therapy less than 1 year was indicated.

⁴Excludes New York City.

⁵Not included in U.S. totals.

See Technical Notes for description of COT calculation (page 9).