
Asbestosis: Proportionate mortality ratio (PMR) adjusted for age, sex, and race by usual industry, U.S. residents age 15 and over, selected states and years, 1990–1999

CIC	Industry	Number of Deaths	PMR	95% Confidence Interval	
				LCL	UCL
262	Miscellaneous nonmetallic mineral and stone products	75	16.4	13.0	20.7
360	Ship and boat building and repairing	171	15.7	13.5	18.3
502	Lumber and construction materials	20	7.0	4.2	10.8
192	Industrial and miscellaneous chemicals	124	4.8	4.0	5.7
211	Other rubber products, and plastics footwear and belting	40	4.3	3.1	5.9
462	Electric and gas, and other combinations	14	3.1	1.7	5.1
180	Plastics, synthetics, and resins	12	2.8	1.4	4.9
200	Petroleum refining	31	2.7	1.9	3.9
272	Primary aluminum industries	16	2.7	1.5	4.3
460	Electric light and power	55	2.7	2.0	3.5
250	Glass and glass products	30	2.6	1.7	3.7
881	Membership organizations	13	2.5	1.3	4.2
060	Construction	702	2.4	2.2	2.6
282	Fabricated structural metal products	28	2.3	1.5	3.3
420	Water transportation	24	2.3	1.5	3.4
210	Tires and inner tubes	15	2.2	1.3	3.7
400	Railroads	89	1.6	1.3	2.0
270	Blast furnaces, steelworks, rolling and finishing mills	67	1.3	1.0	1.7

CIC - Census Industry Code LCL - lower confidence limit UCL - upper confidence limit

NOTE: The total number of deaths with asbestosis reported was 2,859 in these same selected states and years, and the comparable number of asbestosis deaths in the entire United States for this same time period was 10,914. See selected limitations for general cautions regarding inferences based on small numbers of deaths, and see appendices for source description, methods, ICD codes, industry and occupation codes, and list of selected states and years.

SOURCE: National Center for Health Statistics multiple cause-of-death data.