

6 FOCUS ON MINING

This section presents a detailed overview of mining and injuries. Historically, mining has been the industry sector with the highest fatal and nonfatal injury rates. Mining still has the highest fatal injury rate—more than five times the national average. Because of the accompanying social toll and public pressures for action, data on fatalities and injuries became available far earlier in the mining industry than in many others. State and Federal agencies began collecting data in the 1870s, and reliable information has been available for an entire century. Examination of the history of coal mining in this century shows important relationships between adverse outcomes (such as fatality rates) and regulatory actions such as enactment of Federal legislation and establishment of enforcement and consultation agencies.

Fatal Injuries

Historical Perspective

More than 103,000 workers died in the mining industry (including all commodities and work locations) during the 85-year period from 1911 to 1995 (Figure 6–1). From 1911 through 1915 alone, 16,646 fatalities occurred, with an annual average of 2,517 deaths in coal mining and 813 in metal and nonmetal mining. The corresponding annual average fatality rates during this 5-year period were 340 and 300 per 100,000 workers in coal mining and metal and nonmetal mining, respectively. The U.S. Bureau of Mines, established in 1910, focused on coal mine fires and explosions. In the first decade after its creation, disaster-related fatalities* in coal mines decreased substantially, with a 62% reduction in deaths from mine fires and a 45% reduction from explosions. Disaster-related fatalities in metal and nonmetal mining increased during the same period. The number and rate of fatalities decreased again during the Great Depression, a period accompanied by reductions in both the labor force and production, as measured by tonnage mined. Rates increased during the economic mobilization required during World War II. For the years 1941–1945, 694 million tons of coal were mined annually, compared with 520 million tons for the 5 preceding years—a 33% increase. Coal mining

*Five or more fatalities resulting from a single incident.

fatality rates increased during the decade preceding the Coal Mine Health and Safety Act of 1969 [Public Law 91–173]. Rates rose similarly for metal and nonmetal mining in the decade before the Federal Mine Safety and Health Act of 1977 [Public Law 95–164]. Fatality rates decreased following the passage of these two Federal mine acts.

Fatal Injuries during 1988–1997

Mining operations are located in every State as well as in Puerto Rico and the Virgin Islands. In 1997, 13,682 mining operations reported employment to the Mine Safety and Health Administration (MSHA)—2,609 were coal mining operations (average employment size of 35 full-time workers); 374 were metal mining operations (average employment size of 122 full-time workers); 786 were nonmetal mining operations (average employment size of 32 full-time workers); 3,712 were stone mining operations (average employment size of 19 full-time workers); and 6,201 were sand and gravel mines (average employment size of 5 full-time workers).




Figure 6–1. Number of fatalities (5-year aggregates) and annual average fatality rates in the mining industry by commodity, 1911–1995. Annual average fatality rates are calculated per 100,000 workers aggregated over 5-year periods. Metal and nonmetal includes metal, nonmetal, stone, and sand and gravel. (Source: MSHA [1999]; Adams and Wrenn [1941]; Adams and Kolhos [1941]; Reese et al. [1955]; MSHA [1984].)

At least one mine operator fatality occurred in each State but Delaware, Maine, and Rhode Island during the period 1988–1997 (Figure 6–2). The national annual average mine operator fatality rate during this period was 28.5 per 100,000 miners, which is more than five times the national annual average occupational fatality rate of 5.3 per 100,000 workers from 1980 through 1995 [NTOF 1999] (see Figure 2–1 and preceding text). Nine states had very high mine operator fatality rates that exceeded the national mining fatality rate by 50% or more. Nine other States and Puerto Rico had rates greater than the national rate.




Figure 6–2. Number of mine operator fatalities (as numbers within State boundaries) and corresponding annual average fatality rates (shading within States) for each State and Puerto Rico, 1988–1997. (Source: MSHA [1999].)

FOCUS ON MINING

During the 10-year period 1988–1997, the number of fatalities was highest in 1990 (n=122) and lowest in 1994 (n=82) (Figure 6–3). The overall downward trend since 1990 is attributable primarily to decreases in fatalities of mine operator workers in the coal and metal commodities. During 1988–1997, mine operator workers accounted for 806 (81.2%) of the total number of fatalities (n=993), with the remaining 187 (18.8%) being independent contractor workers. Coal operator workers accounted for 432 (43.5%) of the total, followed by mine operator workers in the commodities of stone (n=157, 15.8%), metal (n=98, 9.9%), sand and gravel (n=83, 8.4%), and nonmetal (n=36, 3.6%). Independent contractor workers in metal and nonmetal mining made up 11.3% (112) of the total, with 7.6% (75) attributed to independent contractors in coal.

Despite annual fluctuations, overall fatality rates have decreased for mine operator workers in the coal and metal commodities (Figure 6–4). Although fatality rates for independent contractor workers are the highest overall, reduced employment reporting requirements for this sector of workers compromise any direct comparison with rates for mine operator workers.




Figure 6–3. Number of fatalities, by type of employer (mine operator versus independent contractor) and commodity, 1988–1997. Metal and nonmetal includes metal, nonmetal, stone, and sand and gravel. (Source: MSHA [1999].)




Figure 6-4. Fatality rates by type of employer (mine operator versus independent contractor) and commodity, 1988-1997. Fatality rates are calculated per 100,000 full-time workers or 200 million employee hours. Metal and nonmetal includes metal, nonmetal, stone, and sand and gravel. (Source: MSHA [1999].)

Both the number and annual average rate of fatalities from 1988 through 1997 varied by work location and by type of employer and commodity (Table 6–1). Overall, underground work locations exhibited both the highest numbers and rates of fatalities, and preparation plants and mills exhibited the lowest fatality rates. Among mine operator workers, the combination of high numbers and rates of fatalities is most conspicuous for those working in underground coal mines, underground metal mines, stone surface mines (or quarries), and sand and gravel operations. Although independent contractor workers accounted for about one-fifth (19%) of all fatalities during this 10-year period, they accounted for almost one-third (30%) of the fatalities at the surface areas of underground mines, at surface mines, and at mills or preparation plants.

A trend toward decreasing fatality rates with increasing mine size is apparent in underground coal and metal mining operations (Figure 6–5). This trend is particularly significant for underground coal mining operations, which also accounted for 83% of all underground employee hours and 77% of all underground fatalities reported during the 10-year period 1988–1997. In addition, coal accounted for 90% of the employee hours worked underground in small mines (fewer than 50 workers), followed by stone mines (5.5%), metal mines (3%), and nonmetal mines (less than 1%).




Figure 6–5. Fatality rates at underground mining operations, by commodity and employment size of operation, 1988–1997. Fatality rates are computed per 100,000 full-time workers or 200 million employee hours. (Note: There are no sand and gravel underground operations.) (Source: MSHA [1999].)

Table 6–1. Number and annual average rate* of fatalities associated with various types of employers and commodities by work location, 1988–1997

Type of employer and commodity	Work location													
	Underground mines						Surface mines							
	All		Underground		Surface areas		Strip/ open pit/ quarry		Dredge		Other surface operations [†]		Mills/plants	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
All	993	31.9	388	56.9	51	46.3	383	29.4	21	38.3	10	37.2	140	15.0
Mine operator:														
Coal	432	37.9	298	54.2	27	49.5	74	19.1	0	0.0	4	25.7	29	22.0
Metal	98	21.6	53	83.8	6	36.5	28	16.6	0	0.0	0	0.0	11	5.7
Nonmetal	36	13.4	14	40.0	0	0.0	11	18.5	0	0.0	NA [‡]	NA	11	6.6
Stone	157	23.5	15	93.5	3	77.1	95	33.7	0	0.0	0	0.0	44	12.1
Sand and gravel	83	28.2	NA	NA	NA	NA	65	26.3	18	38.3	NA	NA	NR [§]	NR
Independent contractor:														
Coal	75	59.1	2	17.6	12	52.3	40	62.5	0	0.0	5	253.2	16	60.3
Metal and nonmetal**	112	71.3	6	102.0	3	49.0	70	75.0	3	278.1	1	232.7	29	57.7

Source: MSHA [1999].

*Computed per 100,000 full-time workers or 200 million employee hours.

[†]Includes culm banks, auger mining, independent shops and yards, and surface mining n.e.c.

[‡]NA=Not applicable for this commodity.

[§]NR=Not reported separately. Sand and gravel operators report mill employment under strip or dredge operations.

**Includes metal, nonmetal, stone, and sand and gravel.




Figure 6-6. Fatality rates at surface mining operations, by commodity and employment size of operation, 1988-1997. Fatality rates are computed per 100,000 full-time workers or 200 million employee hours. (Source: MSHA [1999].)

Fatality rates from 1988 through 1997 at surface work locations were highest at the smallest mining operations (fewer than 10 workers) in every commodity (Figure 6-6). The proportion of employee hours worked at the surface locations of these mines was highest in sand and gravel (43%), followed by stone (8%), coal (6%), nonmetal (5%), and metal (less than 1%). Elevated rates persist for small operations with 10 to 49 workers, although these rates are only slightly elevated in coal. No sand and gravel operations employed more than 250 workers.

The major types of incidents associated with fatal injuries are shown in Table 6-2 by commodity and type of employer. Overall, powered haulage incidents accounted for the largest percentage of fatalities (30.8%), followed by fall of ground (18.9%) and machinery (16.3%). Some types of incidents, such as falls of ground, are substantially more frequent among mine operators than among independent contractors. The rate differences between type of employer and among commodities suggest that different strategies are needed to reduce fatality rates in various sectors of the mining industry.

Table 6–2. Number* and annual average rate† of fatalities associated with various types of employers and commodities, by type of incident,‡ 1988–1997

Type of employer and commodity	Type of incident											
	All		Powered haulage		Fall of ground (from in place)		Machinery		Electrical		Slip or fall of person	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
All	993	31.9	306	9.8	188	6.0	162	5.2	80	2.6	60	1.9
Mine operator:												
Coal	432	37.9	101	8.9	135	11.8	76	6.7	33	2.9	9	0.8
Metal	98	21.6	30	6.6	18	4.0	6	1.3	11	2.4	8	1.8
Nonmetal	36	13.4	11	4.1	11	4.1	4	1.5	5	1.9	2	0.7
Stone	157	23.5	64	9.6	16	2.4	18	2.7	12	1.8	13	1.9
Sand and gravel	83	28.2	38	12.9	1	0.3	16	5.4	5	1.7	7	2.4
Independent contractor:												
Coal	75	59.1	34	26.8	2	1.6	10	7.9	8	6.3	8	6.3
Metal and nonmetal§	112	71.3	28	17.8	5	3.2	32	20.4	6	3.8	13	8.3

See footnotes at end of table.

(Continued)

Table 6–2 (Continued). Number* and annual average rate† of fatalities associated with various types of employers and commodities, by type of incident,‡ 1988–1997

Type of employer and commodity	Type of incident									
	Falling, rolling, or sliding rock or material		Ignition/ explosion of gas/dust		Explosives and breaking agents		Unknown or n.e.c.		Hand tools	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
All	57	1.8	35	1.1	31	1.0	27	0.9	15	0.5
Mine operator:										
Coal	13	1.1	28	2.5	12	1.1	9	0.8	6	0.5
Metal	13	2.9	1	0.2	4	0.9	1	0.2	0	0.0
Nonmetal	0	0.0	0	0.0	0	0.0	1	0.4	1	0.4
Stone	16	2.4	2	0.3	7	1.0	3	0.4	3	0.4
Sand and gravel	4	1.4	1	0.3	0	0.0	6	2.0	1	0.3
Independent contractor:										
Coal	4	3.2	3	2.4	2	1.6	2	1.6	1	0.8
Metal and nonmetal [§]	7	4.5	0	0.0	6	3.8	5	3.2	3	1.9

See footnotes at end of table.

(Continued)

Table 6–2 (Continued). Number* and annual average rate† of fatalities associated with various types of employers and commodities, by type of incident,‡ 1988–1997

Type of employer and commodity	Type of incident									
	Exploding vessels under pressure		Handling materials		Fire		Hoisting		Inundation	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
All	11	0.4	8	0.3	7	0.2	4	0.1	2	0.1
Mine operator:										
Coal	3	0.3	1	0.1	4	0.4	0	0.0	2	0.2
Metal	3	0.7	0	0.0	2	0.4	1	0.2	0	0.0
Nonmetal	1	0.4	0	0.0	0	0.0	0	0.0	0	0.0
Stone	1	0.1	1	0.1	1	0.1	0	0.0	0	0.0
Sand and gravel	1	0.3	3	1.0	0	0.0	0	0.0	0	0.0
Independent contractor:										
Coal	0	0.0	0	0.0	0	0.0	1	0.8	0	0.0
Metal and nonmetal§	2	1.3	3	1.9	0	0.0	2	1.3	0	0.0

Source: MSHA [1999].

*Note: See Appendix A, *Mining Injury and Employment Statistics*, for selection of fatalities.

†Computed per 100,000 full-time workers or 200 million employee hours.

‡MSHA's accident/injury/illness classification. See Appendix A, *Mining Injury and Employment Statistics*, for modifications.

§Includes metal, nonmetal, stone, and sand and gravel.

Lost-Workday Injuries

Lost-workday injury rates decreased between 1988 and 1997 in all five commodities for mine operator workers and independent contractor workers in metal and nonmetal mining (Figure 6–7). The lost-workday injury rates for independent contractor workers in coal follow a somewhat different trend, with a gradual increase from 1988 through 1992, then a slow decrease for the remainder of the period. Over the 10-year period, the highest lost-workday injury rate was observed in mine operator workers in coal (7.9 cases per 100 full-time workers), followed by mine operator workers in stone (4.7 cases per 100 full-time workers), sand and gravel (4.1 cases per 100 full-time workers), metal (3.9 cases per 100 full-time workers), and nonmetal mining (3.7 cases per 100 full-time workers). Among mine operator workers, the largest percentage of decrease in the lost-workday injury rate occurred in metal mining, dropping from a rate of 5.4 cases per 100 full-time workers in 1988 to 3.1 cases per 100 full-time workers in 1997 (a 43% decrease). Following metal mining, the decreases in lost-workday injury rates were as follows: coal (35%), nonmetal (31%), stone (30%), and sand and gravel mining (24%). A 46% decrease was observed for independent contractor workers in metal and nonmetal mining, and a 6% decrease was observed for independent contractor workers in coal mining.

The number and rate of lost-workday cases from 1988 to 1997 for various work locations are shown by type of employer and commodity in Table 6–3. Overall, underground work locations exhibited both the highest numbers and rates of lost-workday cases, whereas surface strip/open pit/quarry operations exhibited the lowest rates. Among mine operator workers, a combination of high numbers and rates of lost-workday cases are most conspicuous for workers in underground coal mines, stone surface mines (or quarries) and mills, and sand and gravel operations. Independent contractor workers accounted for less than 6% of all lost-workday cases but 19% of all fatalities (see Table 6–2).

The leading types of incidents associated with lost-workday cases (Table 6–4) are handling materials (34.4%), slip or fall of person (20.9%), powered haulage (10.9%), machinery (10.9%), and hand tools (9.5%). These incidents accounted for 86.6% of all cases between 1988 and 1997. Incidents involving handling materials accounted for more than one-third of all lost-workday cases, compared with fewer than 1% of the fatalities. Furthermore, these incidents account for 29.2% of the 6,840,987 lost workdays, followed by slip or fall of person (20.8%) and powered haulage (17.3%).

Sprains to the back region accounted for the largest proportion of lost workdays during the 10-year period (Figure 6–8). Sprains to the lower extremities (primarily the knee), amputations of the arms or hands (primarily the fingers), and fractures to the lower extremities also accounted for a substantial proportion of lost workdays.




Figure 6–7. Incidence rates for lost-workday cases, by type of employer and commodity, 1988–1997. Incident rates are calculated per 100 full-time workers or 200,000 employee hours. Metal and nonmetal includes metal, nonmetal, stone, and sand and gravel. (Source: MSHA [1999].)

Table 6–3. Number and annual average rate* of lost-workday cases associated with various types of employers and commodities, by work location, 1988–1997

Type of employer and commodity	Work location													
	All		Underground mines				Surface mines							
			Underground		Surface areas		Strip/ open pit/ quarry		Dredge		Other surface operations [†]		Mills/plants	
Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	
All	170635	5.5	74264	10.9	6272	5.7	46396	3.6	2270	4.1	1250	4.7	40183	4.3
Mine operator:														
Coal	89895	7.9	65668	11.9	4348	8.0	12453	3.2	9	1.7	848	5.4	6569	5.0
Metal	17622	3.9	4534	7.2	641	3.9	5152	3.1	193	4.5	214	3.3	6888	3.5
Nonmetal	9855	3.7	1528	4.4	308	5.0	1623	2.7	31	5.6	NA [‡]	NA	6365	3.8
Stone	31642	4.7	748	4.7	270	6.9	13426	4.8	66	5.6	82	3.3	17050	4.7
Sand and gravel	12059	4.1	NA	NA	NA	NA	10106	4.1	1953	4.2	NA	NA	NR [§]	NR
Independent contractor:														
Coal	4363	3.4	1367	12.1	499	2.2	1385	2.2	2	1.0	96	4.9	1014	3.8
Metal and nonmetal**	5199	3.3	419	7.1	206	3.4	2251	2.4	16	1.5	10	2.3	2297	4.6

Source: MSHA [1999].

*Computed per 100 full-time workers or 200,000 employee hours.

[†]Includes culm banks, auger mining, independent shops and yards, and surface mining n.e.c.

[‡]NA=Not applicable for this commodity.

[§]NR=Not reported separately. Sand and gravel operators report mill employment under strip or dredge operations.

**Includes metal, nonmetal, stone, and sand and gravel.

Table 6–4. Number and annual average rate* of lost-workday cases and mean days lost associated with various types of employers and commodities, by type of incident,[†] 1988–1997

Type of employer and commodity	Type of incident															
	All			Handling materials			Slip or fall of person			Powered haulage			Machinery			
	Number	Rate	MDL [‡]	Number	Rate	MDL	Number	Rate	MDL	Number	Rate	MDL	Number	Rate	MDL	
All	170,635	5.49	40.1	58,661	1.89	34.0	35,679	1.15	39.8	18,676	0.60	63.5	18,647	0.60	43.0	
Mine operator:																
Coal	89,895	7.89	41.7	31,072	2.73	39.0	16,337	1.43	42.5	10,859	0.95	57.7	9,078	0.80	39.2	
Metal	17,622	3.89	39.1	5,975	1.32	32.0	4,067	0.90	40.4	1,591	0.35	59.1	1,950	0.43	41.4	
Nonmetal	9,855	3.67	35.7	3,910	1.46	27.7	2,294	0.85	33.3	912	0.34	77.9	939	0.35	37.9	
Stone	31,642	4.74	33.6	11,108	1.66	26.3	7,588	1.14	33.1	2,825	0.42	63.4	3,752	0.56	36.1	
Sand and gravel	12,059	4.10	39.9	3,760	1.28	23.7	3,077	1.05	34.6	1,512	0.51	90.7	1,334	0.45	60.6	
Independent contractor:																
Coal	4,363	3.44	49.6	1,238	0.98	44.4	1,054	0.83	57.0	515	0.41	51.2	696	0.55	52.3	
Metal and nonmetal [§]	5,199	3.31	55.8	1,598	1.02	30.3	1,262	0.80	54.6	462	0.29	113.2	898	0.57	85.2	

See footnotes at end of table.

(Continued)

Table 6-4 (Continued). Number and annual average rate* of lost-workday cases and mean days lost associated with various types of employers and commodities, by type of incident,[†] 1988-1997

Type of employer and commodity	Type of incident															
	Hand tools			Fall of ground (from in place)			Stepping or kneeling on object			Unknown or n.e.c.			Electrical			
	Number	Rate	MDL [‡]	Number	Rate	MDL	Number	Rate	MDL	Number	Rate	MDL	Number	Rate	MDL	
All	16,134	0.52	28.7	10,522	0.34	45.0	3,446	0.11	25.7	3,203	0.10	38.8	1,571	0.05	53.4	
Mine operator:																
Coal	6,905	0.61	28.9	9,142	0.80	44.5	1,919	0.17	29.8	1,477	0.13	48.3	916	0.08	43.4	
Metal	1,835	0.40	33.6	806	0.18	43.9	404	0.09	21.7	403	0.09	37.5	92	0.02	46.2	
Nonmetal	931	0.35	26.4	166	0.06	45.5	178	0.07	20.7	234	0.09	49.7	74	0.03	23.2	
Stone	4,039	0.60	26.5	123	0.02	79.9	589	0.09	19.7	664	0.10	25.8	251	0.04	46.4	
Sand and gravel	1,554	0.53	26.4	6	0.00	41.5	193	0.07	18.5	256	0.09	22.6	135	0.05	64.2	
Independent contractor:																
Coal	366	0.29	30.0	206	0.16	52.1	66	0.05	23.7	61	0.05	25.3	33	0.03	251.4	
Metal and nonmetal [§]	504	0.32	35.2	73	0.05	45.4	97	0.06	21.1	108	0.07	15.8	70	0.04	136.1	

See footnotes at end of table.

(Continued)

Table 6–4 (Continued). Number and annual average rate* of lost-workday cases and mean days lost associated with various types of employers and commodities, by type of incident,† 1988–1997

Type of employer and commodity	Type of incident															
	Striking or bumping			Nonpowered haulage			Exploding vessels under pressure			Fire			Falling, rolling, or sliding rock or material			
	Number	Rate	MDL‡	Number	Rate	MDL	Number	Rate	MDL	Number	Rate	MDL	Number	Rate	MDL	
All	1,504	0.05	32.8	722	0.02	44.0	398	0.01	43.6	388	0.01	33.1	363	0.01	86.6	
Mine operator:																
Coal	1,130	0.10	35.3	308	0.03	57.5	181	0.02	27.0	144	0.01	27.0	154	0.01	80.1	
Metal	91	0.02	28.7	126	0.03	38.1	53	0.01	36.0	39	0.01	29.5	46	0.01	32.7	
Nonmetal	55	0.02	14.0	46	0.02	36.2	22	0.01	233.1	24	0.01	28.2	22	0.01	35.8	
Stone	139	0.02	26.6	170	0.03	37.5	81	0.01	27.1	100	0.01	49.6	75	0.01	190.1	
Sand and gravel	57	0.02	30.4	38	0.01	12.7	34	0.01	79.4	42	0.01	18.1	36	0.01	42.2	
Independent contractor:																
Coal	18	0.01	27.4	12	0.01	26.7	14	0.01	10.3	24	0.02	28.2	9	0.01	33.9	
Metal and nonmetal§	14	0.01	4.6	22	0.01	20.3	13	0.01	29.4	15	0.01	49.4	21	0.01	34.9	

See footnotes at end of table.

(Continued)

Table 6–4 (Continued). Number and annual average rate* of lost-workday cases and mean days lost associated with various types of employers and commodities, by type of incident,† 1988–1997

Type of employer and commodity	Type of incident														
	Ignition/explosion of gas/dust			Explosives and breaking agents			Hoisting			Entrapment			Inundation		
	Number	Rate	MDL‡	Number	Rate	MDL	Number	Rate	MDL	Number	Rate	MDL	Number	Rate	MDL
All	307	0.01	45.3	219	0.01	139.1	160	0.01	99.5	23	0.00	46.9	12	0.00	114.7
Mine operator:															
Coal	131	0.01	45.3	101	0.01	25.6	33	0.00	64.2	4	0.00	115.0	4	0.00	72.0
Metal	34	0.01	22.4	48	0.01	349.6	55	0.01	54.6	3	0.00	32.0	4	0.00	228.3
Nonmetal	21	0.01	36.2	4	0.00	151.5	20	0.01	63.1	1	0.00	15.0	2	0.00	7.0
Stone	67	0.01	59.5	44	0.01	97.7	17	0.00	436.4	8	0.00	42.9	2	0.00	80.5
Sand and gravel	18	0.01	23.6	2	0.00	1.5	3	0.00	35.3	2	0.00	5.5	0	0.00	0.0
Independent contractor:															
Coal	22	0.02	53.5	12	0.01	102.9	16	0.01	55.7	1	0.00	58.0	0	0.00	0.0
Metal and nonmetal§	14	0.01	60.8	8	0.01	620.8	16	0.01	70.1	4	0.00	23.8	0	0.00	0.0

Source: MSHA [1999].

*Computed per 100 full-time workers or 200,000 employee hours.

†MSHA's accident/injury/illness classification. See Appendix A, *Mining Injury and Employment Statistics*, for modifications.

‡MDL = mean days lost. MDL is average number of days lost (including restricted workdays) per lost-workday case.

§Includes metal, nonmetal, stone, and sand and gravel.




Figure 6-8. Lost workdays in mining, by part of body affected and nature of injury, 1988-1997. (Source: MSHA [1999].)

References

REFERENCES

ABLES [1999]. The Adult Blood-Lead Epidemiology and Surveillance Program (ABLES). Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Database. [www.cdc.gov/niosh/ables.html].

Adams WW, Kolhos ME [1941]. Metal- and nonmetal-mine accidents in the United States during the calendar year 1939 (excluding coal mines). Washington, DC: U.S. Department of the Interior, Bureau of Mines, Bulletin 440.

Adams WW, Wrenn VE [1941]. Quarry accidents in the United States during the calendar year 1939. Washington, DC: U.S. Department of the Interior, Bureau of Mines, Bulletin 438.

BLS [1998]. Employment and wages annual averages, 1997. Washington, DC: U.S. Department of Labor, Bureau of Labor Statistics, Bulletin 2511, p. 535.

BLS [1999]. Current population survey. Washington, DC: U.S. Department of Labor, Bureau of Labor Statistics.

BLS [2000]. A special issue: charting the projections; 1998–2008. *Occup Outlook Q* 43(4):2–38. Winter 1999–2000.

Bureau of the Census [1992]. 1990 census of population and housing: alphabetical index of industries and occupations. Washington, DC: U.S. Government Printing Office, Publication CPH-R-3.

California Department of Health Services [1999]. Sentinel events notification system for occupational risks (SENSOR): occupational carpal tunnel syndrome—California year 2; second report, #60/CCU902990-13. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health.

CDC (Centers for Disease Control and Prevention) [1999]. HIV/AIDS *Surveill Rep* 11(1):26. [www.cdc.gov/hiv/stats/hasr1101.pdf].

CDPR [1999]. Pesticide illness surveillance program. Sacramento, CA: California Environmental Protection Agency, California Department of Pesticide Regulation, Worker Health and Safety Branch. Database.

CFOI [1999]. Census of fatal occupational injuries, 1992–1997. Washington, DC: U.S. Department of Labor, Bureau of Labor Statistics. Database. [www.bls.gov/iif/oshcfoi1.htm].

CWXSP [1999]. Coal workers' X-ray surveillance program 1970–1995. Morgantown, WV: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Database. [aspe.os.dhhs.gov/DATACNCL/datadir/cdc5.htm#cwvsp].

Fullerton HN Jr. [1999]. Labor force projections to 2008: steady growth and changing composition. *Mon Labor Rev* 122(11):19–32.

ILO [1980]. Guidelines for the use of ILO international classification of radiographs of pneumoconioses. Rev. ed. Geneva, Switzerland: International Labour Office, Occupational Safety and Health Series No. 22 (Rev.).

MSHA [1984]. Summary of selected injury experience and worktime for the mining industry in the United States, 1931–77. Denver, CO: U.S. Department of Labor, Mine Safety and Health Administration, IR 1132.

MSHA [1999]. Quarterly employment and coal production: accidents/injuries/illnesses reported to MSHA under 30 CFR Part 50, 1986–1997. Denver, CO: U.S. Department of Labor, Mine Safety and Health Administration, Office of Injury and Employment Information.

NaSH [1999]. National Surveillance System for Hospital Health Care Workers, 1995–1999. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Infectious Diseases. Database. [www.cdc.gov/ncidod/hip/SURVEILL/nash.htm].

NCHS [1999]. Mortality data, multiple-cause-of-death public-use data files, 1968–1996. Hyattsville, MD: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Health Statistics. Database. [www.cdc.gov/nchs/products/elec_prods/subject/mortmcd.htm].

NCID [1999]. Sentinel counties study of acute viral hepatitis, 1985–1995. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Infectious Diseases. Database.

NEISS [1999]. National Electronic Injury Surveillance System. (Data collected by the Consumer Product Safety Commission; work-related case records maintained by the National Institute for Occupational Safety and Health.) Washington, DC: Consumer Product Safety Commission, Division of Hazard and Injury Data Systems; and Morgantown, WV: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Database.

New York State Department of Health [1999]. New York State pesticide poisoning registry, 1992–1996. Albany, NY: New York State Department of Health, Bureau of Occupational Health.

NHAMCS [1999]. National hospital ambulatory medical care survey, 1995–1997. Hyattsville, MD: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Health Statistics. Database. [www.cdc.gov/nchs/products/elec_prods/subject/nhamcs.htm].

NHANES III [1999]. Third National Health and Nutrition Examination Survey. Hyattsville, MD: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Health Statistics. Database. [www.cdc.gov/nchs/about/major/nhanes/datalink.htm].

NIOSH [1997]. Mortality by occupation, industry, and cause of death: 24 reporting States (1984–1988). Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 97-114. [www.cdc.gov/niosh/97-114.html].

NIOSH [1999]. Work-related lung disease surveillance report 1999 (WoRLD). Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 2000-105. [www.cdc.gov/niosh/docs/2000-105/pdfs/2000-105.pdf].

NOMS [1999]. National Occupational Mortality Surveillance System. Cincinnati, OH: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Database. [aspe.os.dhhs.gov/datacncl/datadir/cdc5.htm#noms].

NSSPM [1999]. National Surveillance System for Pneumoconiosis Mortality, 1968–1996. Morgantown, WV: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Database.

NTOF [1999]. National Traumatic Occupational Fatalities Surveillance System. Morgantown, WV: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Database.

Oregon Health Division [1999]. Pesticide Poisoning Prevention Program, 1992–1996. Portland, OR: Oregon Health Division; Center for Disease Prevention and Epidemiology; Environmental and Occupational Epidemiology Section. Database.

PEST [1999]. Pesticide Exposure Surveillance in Texas Program, 1992–1996. Austin, TX: Texas Department of Health, Bureau of Epidemiology, Division of Environmental Epidemiology and Toxicology. Database.

Ramazzini B [1713]. Diseases of workers. *De Morbis Artificum*. (translation by WC Wright, 1964). New York: Hafner Publishing Co.

Reese ST, Wrenn VE, Reid EJ [1955]. Injury experience in coal mining, 1952: analysis of mine safety factors, related employment, and production data. Washington, DC: U.S. Department of the Interior, Bureau of Mines, Bulletin 559.

Rosenman KD, Reilly MJ [1999]. Unpublished data, 1992–1998. East Lansing, MI: Michigan State University, Department of Medicine.

Rosenman KD, Reilly MJ, Deliefde B, Kalinowski DJ [1999]. 1998 annual report on occupational noise-induced hearing loss in Michigan. East Lansing, MI: Michigan State University, Department of Medicine; and Michigan Department of Consumer and Industry Services, Occupational Health Division.

SOII [1999]. Survey of occupational injuries and illnesses. Washington, DC: U.S. Department of Labor, Bureau of Labor Statistics. Database. [www.bls.gov/iif/].

staffTRAK-TB [1999]. Surveillance for tuberculosis infection in health care workers, 1994–1998. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for HIV, STD, and TB Prevention. Database.

TESS [1998]. AAPCC Toxic Exposure Surveillance System. Washington, DC: American Association of Poison Control Centers. Database. [www.aapcc.org/poison1.htm].

VHSP [1999]. Viral Hepatitis Surveillance Program. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Infectious Diseases. Database. [www.cdc.gov/ncidod/osr/site/surv_resources/surv_sys.htm].

Washburn AE, LeBlanc PR, Fahy RF [1999]. Firefighter fatalities: National Fire Protection Association. *NFPA Journal*, July/August, pp. 56–70.

WHO [1977]. Manual of the international statistical classification of diseases, injuries, and causes of death, based on the recommendations of the Ninth Revision Conference, 1975. Geneva, Switzerland: World Health Organization.