§ 180.435

parent compound in or on the following commodities when present therein as a result of application of propiconazole to growing crops in paragraphs (a) and (c) of this section:

Commodity	Parts per million
Alfalfa, forage	0.1 0.1

[71 FR 55306, Sept. 22, 2006, as amended at 72 FR 20439, Apr. 25, 2007]

§ 180.435 Deltamethrin; tolerances for

(a) General. (1) Tolerances are established for the combined residues of the pesticide chemical deltamethrin [(1R,3R)-3-(2,2-dibromoviny1)-2,2-

dimethylcyclopropanecarboxylic acid (S)-alpha-cyano-3-phenoxybenzyl ester and its major metabolites, trans deltamethrin [(S)-alpha-cyano-m-phenoxybenzyl(1R,3S)-3-(2,2-

dibromovinyl)-2,2dimethylcyclopropanecarboxylate] and alpha-R-deltamethrin [(R)-alpha-cyano-m-phenoxybenzyl-(1R,3R)-3-(2,2-

dibromovinyl)-2,2-

dimethylcyclopropanecarboxylate] in or on the following agricultural commodities:

Commodity	Parts per million
Almond, hulls	2.5
Apples, wet pomace	1.0
Artichoke, globe	0.5
Barley, bran	5.0
Cattle, fat	0.05
Cattle, meat	0.02
Cattle, meat byproducts	0.05
Corn, field, forage	0.7
Corn, field, refined oil	2.5
Corn, field, stover	5.0
Corn, pop, stover	5.0
Corn, sweet, forage	10
Corn, sweet, kernel plus cob with husks re-	
moved	0.03
Corn, sweet, stover	15
Cotton, undelinted seed	0.04
Cotton, oil	0.2
Egg	0.02
Fruit, pome, Group 11	0.2
Goat, fat	0.05
Goat, meat	0.02
Goat, meat byproducts	0.05
Grain, aspirated fractions	65
Grain, cereal, Group 15, except sweet corn	1.0
Hog, fat	0.05
Horse, fat	0.05
Horse, meat	0.02
Horse, meat byproducts	0.05
Lvchee*	0.2
Milk, fat (reflecting 0.02 ppm in whole milk)	0.1
Nut, tree, Group 14	

Commodity	Parts per million
Onion, dry bulb	0.1
Onion, green	1.5
Poultry, fat	0.05
Poultry, meat	0.02
Poultry, meat byproducts	0.02
Radish tops	4.0
Rapeseed	0.2
Rice, hulls	2.5
Rye, bran	5.0
Sheep, fat	0.05
Sheep, meat	0.02
Sheep, meat byproducts	0.05
Sorghum, grain forage	0.5
Sorghum, grain stover	1.0
Soybean, seed	0.1
Soybean, hulls	0.2
Starfruit*	0.2
Sunflower seed	0.1
Tomato	0.2
Tomato, concentrated products	1.0
Vegetable, cucurbit, Group 9	0.2
Vegetable, fruiting, Group 8	0.3
Vegetable, root, except sugar beet, Subgroup IB	0.2
Vegetable, tuberous and corm, Subgroup IC	0.04
Wheat, bran	5.0

*There are no U.S. registrations for use of deltamethrin on starfruit and lychee.

(2) A tolerance of 0.05 ppm is established for residues of the insecticide deltamethrin (1R,3R)-3-(2,2-dibromovinyl)-2,2-

dimethylcyclopropanecarboxylic acid (S)-alpha-cyano-3-phenoxybenzyl ester and its major metabolites, trans deltamethrin (S)-alpha-cyano-m-phenoxybenzyl-(1R,3R)-3-(2,2-dibromovinyl)-2,2-

dimethylcyclopropanecarboxylate and alpha-*R*-deltamethrin[(*R*)-alpha-cyano-*m*-phenoxybenzyl-(1*R*,3*R*)-3-(2,2-dibromovinyl)-2,2-

dimethylcyclopropanecarboxylate] as follows:

- (i) In or on all food/feed items (other than those covered by a higher tolerance as a result of use on growing crops) in food/feed handling establishments.
- (ii) The insecticide may be present as a residue from application of deltamethrin in food handling establishments, including food service, manufacturing and processing establishments, such as restaurants, cafeterias, supermarkets, bakeries, breweries, dairies, meat slaughtering and packing plants, and canneries, feed handling establishments including feed manufacturing and processing establishments, in accordance with the following prescribed conditions:

Environmental Protection Agency

- (A) Application shall be limited to general surface and spot and/or crack and crevice treatment in food/feed handling establishments where food/feed and food/feed products are held, processed, prepared and served. General surface application may be used only when the facility is not in operation provided exposed food/feed has been covered or removed from the area being treated. Spot and/or crack and crevice application may be used while the facility is in operation provided exposed food/feed is covered or removed from the area being treated prior to application. Spray concentration shall be limited to a maximum of 0.06 percent active ingredient. Contamination of food/feed or food/feed contact surfaces shall be avoided.
- (B) To assure safe use of the insecticide, its label and labeling shall conform to that registered with the U.S. Environmental Protection Agency and shall be used in accordance with such label and labeling.
- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[62 FR 63001, Nov. 26, 1997, as amended at 63 FR 45414, Aug. 26, 1998; 69 FR 62614, Oct. 27, 2004]

§180.436 Cyfluthrin; tolerances for residues.

(a) General. (1) Tolerances are established for residues of the insecticide cyfluthrin (cyano(4-fluoro-3-phenoxyphenyl)methyl-3-(2,2-dichloroethenyl)-2,2dimethyl-cyclopropane-carboxylate; CAS No. 68359-37-5) in or on the following raw agricultural commodities:

Commodity	Parts per million
Alfalfa	5.0
Alfalfa, hay	10.0
Almond, hulls	0.5
Barley, bran	5.0
Brassica, head and stem, subgroup 5A	2.5
Brassica, leafy greens, subgroup 5B	7.0
Carrot, roots	0.20
Cattle, fat	10.0
Cattle, meat	0.40
Cattle, meat byproducts	0.40
Citrus, dried pulp	0.3
Citrus, oil	0.3
Corn, field, forage	3.0

Commodity	Parts per million
Corn, field, milled byproducts	7.0
Corn, field, refined oil	30.0
Corn, field, stover	6.0
Corn, pop, stover	6.0 15.00
Corn, sweet, kernel plus cob with husks re-	10.00
moved	0.05
Corn, sweet, stover	30.00
Cotton, hulls	2.0
Cotton, refined oil Cotton, undelinted seed	2.0 1.0
Egg	0.01
Fruit, citrus, group 10	0.2
Fruit, pome, group 11	0.5
Fruit, stone, group 12	0.3
Goat, fat	10.0 0.40
Goat, meat byproducts	0.40
Grain, aspirated fractions	600
Grain, cereal, group 15	4.0
Grape	1.0
Grape, raisin	3.5
Hog, fat	10.0 0.40
Hog, meatHog, meat byproducts	0.40
Hop, dried cones	20.0
Hop, vines	4.0
Horse, fat	10.0
Horse, meat	0.40
Horse, meat byproducts Lettuce, head	0.40 2.0
Lettuce, leaf	3.0
Milk	1.0
Milk, fat	30.0
Mustard greens	7.0
Nut, tree, group 14	0.01
Oat, bran Pea and bean, dried shelled, except soybean,	5.0
subgroup 6C	0.15
Pea, dry, seed	0.15
Pea, southern, succulent	0.25
Peanut	0.01
Peanut, hay	6.0
Pepper Pistachio	0.50 0.01
Poultry, fat	0.01
Poultry, meat	0.01
Poultry, meat byproducts	0.01
Radish, roots	1.0
Rice, bran	6.0 18.0
Rye, bran	5.0
Sheep, fat	10.0
Sheep, meat	0.40
Sheep, meat byproducts	0.40
Sorghum, grain, forage	2.0
Sorghum, grain, stover	5.0 8.0
Soybean, hay	4.0
Soybean, seed	0.03
Sugarcane, cane	0.05
Sugarcane, molasses	0.20
Sunflower, forage	5.0
Sunflower, seed Tomato	0.02 0.20
Tomato, paste	0.20
Tomato, pomace	5.0
Turnips, greens	7.0
Vegetable, cucurbit, group 9	0.1
Vegetable, fruiting, group 8 Vegetable, leafy greens, except Brassica, group	0.5
Vegetable, tuberous and corm, subgroup 1C	6.0 0.01