

are therefore subject to the same comment as applies to those intended for oral administration.

§ 201.301 Notice to manufacturers, packers, and distributors of estrogenic hormone preparations.

Some drug preparations fabricated wholly or in part from estradiol and labeled as to potency in terms of international units or in terms of international units of estrone activity have been marketed. The international unit of the estrus-producing hormone was established by the International Conference on the Standardization of Sex Hormones at London, England, on August 1, 1932. This unit was defined as "the specific estrus-producing activity contained in 0.1 gamma (=0.0001 mg.) of the standard" hydroxyketonic hormone found in urine (estrone). The International Conference declared that it did not recommend the determination of the activity of nonhydroxyketonic forms of estrogenic hormones in units of estrone because of the varying ratios between the activity of such nonhydroxyketonic estrogenic hormones and estrone, when measured by different methods on test animals. There is no international unit for measuring the activity of estradiol and no accepted relationship between its activity and that of estrone, either in test animals or in humans. The declaration of potency of estradiol in terms of international units or in terms of international units of estrone activity is therefore considered misleading, within the meaning of 21 U.S.C. 352(a). The declaration of the estradiol content of an estrogenic hormone preparation in terms of weight is considered appropriate.

§ 201.302 Notice to manufacturers, packers, and distributors of drugs for internal use which contain mineral oil.

(a) In the past few years research studies have altered medical opinion as to the usefulness and harmfulness of mineral oil in the human body. These studies have indicated that when mineral oil is used orally near mealtime it interferes with absorption from the digestive tract of provitamin A and the fat-soluble vitamins A, D, and K, and consequently interferes with the utili-

zation of calcium and phosphorus, with the result that the user is left liable to deficiency diseases. When so used in pregnancy it predisposes to hemorrhagic disease of the newborn.

(b) There is accumulated evidence that the indiscriminate administration of mineral oil to infants may be followed by aspiration of the mineral oil and subsequent "lipoid pneumonia."

(c) In view of these facts, the Department of Health and Human Services will regard as misbranded under the provisions of the Federal Food, Drug, and Cosmetic Act a drug for oral administration consisting in whole or in part of mineral oil, the labeling of which encourages its use in pregnancy or indicates or implies that such drug is for administration to infants.

(d) It is also this Department's view that the act requires the labelings of such drugs to bear a warning against consumption other than at bedtime and against administration to infants. The following form of warning is suggested: "Caution: To be taken only at bedtime. Do not use at any other time or administer to infants, except upon the advice of a physician."

(e) This statement of interpretation does not in any way exempt mineral oil or preparations containing mineral oil from complying in all other respects with the requirements of the Federal Food, Drug, and Cosmetic Act.

§ 201.303 Labeling of drug preparations containing significant proportions of wintergreen oil.

(a) Because methyl salicylate (wintergreen oil) manifests no toxicity in the minute amounts in which it is used as a flavoring, it is mistakenly regarded by the public as harmless even when taken in substantially larger amounts. Actually, it is quite toxic when taken in quantities of a teaspoonful or more. Wintergreen oil and preparations containing it have caused a number of deaths through accidental misuse by both adults and children. Children are particularly attracted by the odor and are likely to swallow these products when left within reach.

(b) To safeguard against fatalities from this cause, the Department of Health and Human Services will regard as misbranded under the provisions of

§ 201.304

the Federal Food, Drug, and Cosmetic Act any drug containing more than 5 percent methyl salicylate (wintergreen oil), the labeling of which fails to warn that use otherwise than as directed therein may be dangerous and that the article should be kept out of reach of children to prevent accidental poisoning.

(c) This statement of interpretation in no way exempts methyl salicylate (wintergreen oil) or its preparations from complying in all other respects with the requirements of the Federal Food, Drug, and Cosmetic Act.

§ 201.304 Tannic acid and barium enema preparations.

(a) It has become a widespread practice for tannic acid to be added to barium enemas to improve X-ray pictures. Tannic acid is capable of causing diminished liver function and severe liver necrosis when absorbed in sufficient amounts. The medical literature reports a number of deaths associated with the addition of tannic acid to barium enemas. There is a lack of scientific evidence to establish the conditions, if any, under which tannic acid is safe and effective for use in enemas. Tannic acid for rectal use to enhance X-ray visualization is regarded as a new drug within the meaning of section 201(p) of the Federal Food, Drug, and Cosmetic Act.

(b) In view of the hazards involved when tannic acid is used in barium enemas, any shipments of tannic acid labeled to come within the exemptions under 502(f) of the Act containing such phrases as: "Caution: For manufacturing, processing, or repackaging," "For prescription compounding," or "Diagnostic reagent—For professional use only" will be regarded by the Commissioner of Food and Drugs as misbranded within the meaning of section 502(f) of the Federal Food, Drug, and Cosmetic Act unless the label and the labeling bear conspicuously a warning to the effect: "*Warning*—Not for use in enemas."

(c) Any tannic acid intended for use by man and found within the jurisdiction of the Federal Food, Drug, and Cosmetic Act labeled contrary to this section after 60 days from the date of its publication in the FEDERAL REG-

21 CFR Ch. I (4-1-01 Edition)

ISTER may be made the subject of regulatory proceedings.

§ 201.305 Isoproterenol inhalation preparations (pressurized aerosols, nebulizers, powders) for human use; warnings.

(a) Accumulating reports have been received by the Food and Drug Administration and have appeared in the medical literature of severe paradoxical bronchoconstriction associated with repeated, excessive use of isoproterenol inhalation preparations in the treatment of bronchial asthma and other chronic bronchopulmonary disorders. The cause of this paradoxical reaction is unknown; it has been observed, however, that patients have not responded completely to other forms of therapy until use of the isoproterenol inhalation preparation was discontinued. In addition, sudden unexpected deaths have been associated with the excessive use of isoproterenol inhalation preparations. The mechanism of these deaths and their relationship, if any, to the cases of severe paradoxical bronchospasm are not clear. Cardiac arrest was noted in several of these cases of sudden death.

(b) On the basis of the above information and after discussion with and concurrence of the Respiratory and Anesthetic Drugs Advisory Committee for Food and Drug Administration, the Commissioner of Food and Drugs concludes that in order for the labeling of such drugs to bear adequate information for their safe use, as required by § 201.100, such labeling must include the following:

Warning: Occasional patients have been reported to develop severe paradoxical airway resistance with repeated, excessive use of isoproterenol inhalation preparations. The cause of this refractory state is unknown. It is advisable that in such instances the use of this preparation be discontinued immediately and alternative therapy instituted, since in the reported cases the patients did not respond to other forms of therapy until the drug was withdrawn.

Deaths have been reported following excessive use of isoproterenol inhalation preparations and the exact cause is unknown. Cardiac arrest was noted in several instances.

(c)(1) The Commissioner also concludes that in view of the manner in