

NOTE: The indented statement below is required by the Federal government's Clean Air Act for all products containing or manufactured with chlorofluorocarbons (CFC's) [or name of other class I substance, if applicable]:

WARNING: Contains [or Manufactured with, if applicable] *[insert name of substance]*, a substance which harms public health and the environment by destroying ozone in the upper atmosphere.

A notice similar to the above WARNING has been placed in the information for the patient [or patient information leaflet, if applicable] of this product under the Environmental Protection Agency's (EPA's) regulations. The patient's warning states that the patient should consult his or her physician if there are questions about alternatives.

(c)(1) For over-the-counter drug products for human use, the following alternative warning statement may be used:

NOTE: The indented statement below is required by the Federal government's Clean Air Act for all products containing or manufactured with chlorofluorocarbons (CFC's) [or other class I substance, if applicable]:

WARNING: Contains [or Manufactured with, if applicable] *[insert name of substance]*, a substance which harms public health and environment by destroying ozone in the upper atmosphere.

CONSULT WITH YOUR PHYSICIAN OR HEALTH PROFESSIONAL IF YOU HAVE ANY QUESTION ABOUT THE USE OF THIS PRODUCT.

(2) The warning statement shall be clearly legible and conspicuous on the product, its immediate container, its outer packaging, or other labeling in accordance with the requirements of 40 CFR part 82 and appear with such prominence and conspicuousness as to render it likely to be read and understood by consumers under normal conditions of purchase.

(d) This section does not replace or relieve a person from any requirements imposed under 40 CFR part 82.

[61 FR 20100, May 3, 1996]

§ 201.322 Over-the-counter drug products containing internal analgesic/antipyretic active ingredients; required alcohol warning.

(a) People who regularly consume large quantities of alcohol (three or more drinks every day) have an increased risk of adverse effects (possible liver damage or gastrointestinal bleed-

ing). OTC drug products containing internal analgesic/antipyretic active ingredients may cause similar adverse effects. FDA concludes that the labeling of OTC drug products containing internal analgesic/antipyretic active ingredients should advise consumers with a history of heavy alcohol use to consult a physician. Accordingly, any OTC drug product, labeled for adult use, containing any internal analgesic/antipyretic active ingredients (including, but not limited to, acetaminophen, aspirin, carbaspirin calcium, choline salicylate, ibuprofen, ketoprofen, magnesium salicylate, naproxen sodium, and sodium salicylate) alone or in combination shall bear an alcohol warning statement in its labeling as follows:

(1) *Acetaminophen*. "Alcohol Warning" [heading in boldface type]: "If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage."

(2) *Nonsteroidal anti-inflammatory analgesic/antipyretic active ingredients—including but not limited to aspirin, carbaspirin calcium, choline salicylate, ibuprofen, ketoprofen, magnesium salicylate, naproxen sodium, and sodium salicylate*. "Alcohol Warning" [heading in boldface type]: "If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take [insert one nonsteroidal anti-inflammatory analgesic/antipyretic active ingredient] or other pain relievers/fever reducers. [Insert one nonsteroidal anti-inflammatory analgesic/antipyretic active ingredient] may cause stomach bleeding."

(3) *Combinations of acetaminophen with nonsteroidal anti-inflammatory analgesic/antipyretic active ingredients—including but not limited to aspirin, carbaspirin calcium, choline salicylate, ibuprofen, ketoprofen, magnesium salicylate, naproxen sodium, and sodium salicylate*. "Alcohol Warning" [heading in boldface type]: "If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take [insert acetaminophen and one nonsteroidal anti-inflammatory analgesic/antipyretic active ingredient—including, but not limited to aspirin, carbaspirin calcium, choline salicylate, magnesium

salicylate, or sodium salicylate] or other pain relievers/fever reducers. [Acetaminophen and (insert one nonsteroidal anti-inflammatory analgesic/antipyretic ingredient—including, but not limited to aspirin, carbaspirin calcium, choline salicylate, magnesium salicylate, or sodium salicylate] may cause liver damage and stomach bleeding.”

(b) *Requirements to supplement approved application.* Holders of approved applications for OTC drug products that contain internal analgesic/antipyretic active ingredients that are subject to the requirements of paragraph (a) of this section must submit supplements under §314.70(c) of this chapter to include the required warning in the product’s labeling. Such labeling may be put into use without advance approval of FDA provided it includes the exact information included in paragraph (a) of this section.

(c) Any drug product subject to this section that is not labeled as required and that is initially introduced or initially delivered for introduction into interstate commerce after April 23, 1999, is misbranded under section 502 of the Federal Food, Drug, and Cosmetic Act (21 U.S.C. 352) and is subject to regulatory action.

[63 FR 56801, Oct. 23, 1998]

§ 201.323 Aluminum in large and small volume parenterals used in total parenteral nutrition.

(a) The aluminum content of large volume parenteral (LVP) drug products used in total parenteral nutrition (TPN) therapy must not exceed 25 micrograms per liter ($\mu\text{g/L}$).

(b) The package insert of LVP’s used in TPN therapy must state that the drug product contains no more than 25 $\mu\text{g/L}$ of aluminum. This information must be contained in the “Precautions” section of the labeling of all large volume parenterals used in TPN therapy.

(c) The maximum level of aluminum present at expiry must be stated on the immediate container label of all small volume parenteral (SVP) drug products and pharmacy bulk packages (PBP’s) used in the preparation of TPN solutions. The aluminum content must be stated as follows: “Contains no more

than $_\mu\text{g/L}$ of aluminum.” The immediate container label of all SVP’s and PBP’s that are lyophilized powders used in the preparation of TPN solutions must contain the following statement: “When reconstituted in accordance with the package insert instructions, the concentration of aluminum will be no more than $_\mu\text{g/L}$.” This maximum level of aluminum must be stated as the highest of:

(1) The highest level for the batches produced during the last 3 years;

(2) The highest level for the latest five batches, or

(3) The maximum historical level, but only until completion of production of the first five batches after January 26, 2001.

(d) The package insert for all LVP’s, all SVP’s, and PBP’s used in TPN must contain a warning statement. This warning must be contained in the “Warnings” section of the labeling. The warning must state:

WARNING: This product contains aluminum that may be toxic. Aluminum may reach toxic levels with prolonged parenteral administration if kidney function is impaired. Premature neonates are particularly at risk because their kidneys are immature, and they require large amounts of calcium and phosphate solutions, which contain aluminum.

Research indicates that patients with impaired kidney function, including premature neonates, who receive parenteral levels of aluminum at greater than 4 to 5 $\mu\text{g/kg/day}$ accumulate aluminum at levels associated with central nervous system and bone toxicity. Tissue loading may occur at even lower rates of administration.

(e) Applicants and manufacturers must use validated assay methods to determine the aluminum content in parenteral drug products. The assay methods must comply with current good manufacturing practice requirements. Applicants must submit to the Food and Drug Administration validation of the method used and release data for several batches. Manufacturers of parenteral drug products not subject to an approved application must make assay methodology available to FDA during inspections. Holders of pending applications must submit an amendment under §314.60 or §314.96 of this chapter.

[65 FR 4110, Jan. 26, 2000]