

waive the requirement for the submission of evidence obtained in vivo demonstrating the bioavailability of the drug product if the drug product meets one of the following criteria:

(1) [Reserved]

(2) The drug product is in the same dosage form, but in a different strength, and is proportionally similar in its active and inactive ingredients to another drug product for which the same manufacturer has obtained approval and the conditions in paragraphs (d)(2)(i) through (d)(2)(iii) of this section are met:

(i) The bioavailability of this other drug product has been demonstrated;

(ii) Both drug products meet an appropriate in vitro test approved by FDA; and

(iii) The applicant submits evidence showing that both drug products are proportionally similar in their active and inactive ingredients.

(iv) This subparagraph does not apply to enteric coated or controlled release dosage forms.

(3) The drug product is, on the basis of scientific evidence submitted in the application, shown to meet an in vitro test that has been correlated with in vivo data.

(4) The drug product is a reformulated product that is identical, except for a different color, flavor, or preservative that could not affect the bioavailability of the reformulated product, to another drug product for which the same manufacturer has obtained approval and the following conditions are met:

(i) The bioavailability of the other product has been demonstrated; and

(ii) Both drug products meet an appropriate in vitro test approved by FDA.

(e) FDA, for good cause, may waive a requirement for the submission of evidence of in vivo bioavailability if waiver is compatible with the protection of the public health. For full new drug applications, FDA may defer a requirement for the submission of evidence of in vivo bioavailability if deferral is compatible with the protection of the public health.

(f) FDA, for good cause, may require evidence of in vivo bioavailability or bioequivalence for any drug product if

the agency determines that any difference between the drug product and a listed drug may affect the bioavailability or bioequivalence of the drug product.

[57 FR 17998, Apr. 28, 1992]

§ 320.23 Basis for demonstrating in vivo bioavailability or bioequivalence.

(a)(1) The in vivo bioavailability of a drug product is demonstrated if the product's rate and extent of absorption, as determined by comparison of measured parameters, e.g., concentration of the active drug ingredient in the blood, urinary excretion rates, or pharmacological effects, do not indicate a significant difference from the reference material's rate and extent of absorption. For drug products that are not intended to be absorbed into the bloodstream, bioavailability may be assessed by measurements intended to reflect the rate and extent to which the active ingredient or active moiety becomes available at the site of action.

(2) Statistical techniques used shall be of sufficient sensitivity to detect differences in rate and extent of absorption that are not attributable to subject variability.

(3) A drug product that differs from the reference material in its rate of absorption, but not in its extent of absorption, may be considered to be bioavailable if the difference in the rate of absorption is intentional, is appropriately reflected in the labeling, is not essential to the attainment of effective body drug concentrations on chronic use, and is considered medically insignificant for the drug product.

(b) Two drug products will be considered bioequivalent drug products if they are pharmaceutical equivalents or pharmaceutical alternatives whose rate and extent of absorption do not show a significant difference when administered at the same molar dose of the active moiety under similar experimental conditions, either single dose or multiple dose. Some pharmaceutical equivalents or pharmaceutical alternatives may be equivalent in the extent of their absorption but not in their rate of absorption and yet may be considered bioequivalent because such differences in the rate of absorption are

§ 320.24

21 CFR Ch. I (4-1-01 Edition)

intentional and are reflected in the labeling, are not essential to the attainment of effective body drug concentrations on chronic use, and are considered medically insignificant for the particular drug product studied.

[57 FR 17999, Apr. 28, 1992]

§ 320.24 Types of evidence to establish bioavailability or bioequivalence.

(a) Bioavailability or bioequivalence may be determined by several *in vivo* and *in vitro* methods. FDA may require *in vivo* or *in vitro* testing, or both, to establish the bioavailability of a drug product or the bioequivalence of specific drug products. Information on bioequivalence requirements for specific products is included in the current edition of FDA's publication "Approved Drug Products with Therapeutic Equivalence Evaluations" and any current supplement to the publication. The selection of the method used to meet an *in vivo* or *in vitro* testing requirement depends upon the purpose of the study, the analytical methods available, and the nature of the drug product. Applicants shall conduct bioavailability and bioequivalence testing using the most accurate, sensitive, and reproducible approach available among those set forth in paragraph (b) of this section. The method used must be capable of demonstrating bioavailability or bioequivalence, as appropriate, for the product being tested.

(b) The following *in vivo* and *in vitro* approaches, in descending order of accuracy, sensitivity, and reproducibility, are acceptable for determining the bioavailability or bioequivalence of a drug product.

(1)(i) An *in vivo* test in humans in which the concentration of the active ingredient or active moiety, and, when appropriate, its active metabolite(s), in whole blood, plasma, serum, or other appropriate biological fluid is measured as a function of time. This approach is particularly applicable to dosage forms intended to deliver the active moiety to the bloodstream for systemic distribution within the body; or

(ii) An *in vitro* test that has been correlated with and is predictive of human *in vivo* bioavailability data; or

(iii) An *in vivo* test in animals that has been correlated with and is predictive of human bioavailability data.

(2) An *in vivo* test in humans in which the urinary excretion of the active moiety, and, when appropriate, its active metabolite(s), are measured as a function of time. The intervals at which measurements are taken should ordinarily be as short as possible so that the measure of the rate of elimination is as accurate as possible. Depending on the nature of the drug product, this approach may be applicable to the category of dosage forms described in paragraph (b)(1)(i) of this section. This method is not appropriate where urinary excretion is not a significant mechanism of elimination.

(3) An *in vivo* test in humans in which an appropriate acute pharmacological effect of the active moiety, and, when appropriate, its active metabolite(s), are measured as a function of time if such effect can be measured with sufficient accuracy, sensitivity, and reproducibility. This approach is applicable to the category of dosage forms described in paragraph (b)(1)(i) of this section only when appropriate methods are not available for measurement of the concentration of the moiety, and, when appropriate, its active metabolite(s), in biological fluids or excretory products but a method is available for the measurement of an appropriate acute pharmacological effect. This approach may be particularly applicable to dosage forms that are not intended to deliver the active moiety to the bloodstream for systemic distribution.

(4) Well-controlled clinical trials in humans that establish the safety and effectiveness of the drug product, for purposes of establishing bioavailability, or appropriately designed comparative clinical trials, for purposes of demonstrating bioequivalence. This approach is the least accurate, sensitive, and reproducible of the general approaches for determining bioavailability or bioequivalence. For dosage forms intended to deliver the active moiety to the bloodstream for systemic distribution, this approach may be considered acceptable only when analytical methods cannot be developed to permit use of one of the approaches