

samples of blood should be taken to define adequately the maximum (C_{max}) and minimum (C_{min}) blood concentrations on 2 or more consecutive days to establish that steady-state conditions are achieved.

(2) Whenever comparison of the test product and the reference material is to be based on cumulative urinary excretion-time curves at steady-state, sufficient samples of urine should be taken to define the rate and extent of urinary excretion on 2 or more consecutive days to establish that steady-state conditions are achieved.

(3) A more complete characterization of the blood concentration or urinary excretion rate during the absorption and elimination phases of a single dose administered at steady-state is encouraged to permit estimation of the total area under concentration-time curves or cumulative urinary excretion-time curves and to obtain pharmacokinetic information, e.g., half-life or blood clearance, that is essential in preparing adequate labeling for the drug product.

(e) *Steady-state parameters.* (1) In certain instances, e.g., in a study involving a new drug entity, blood clearances at steady-state obtained in a multiple-dose study should be compared to blood clearances obtained in a single-dose study to support adequate dosage recommendations.

(2) In a linear system, the area under the blood concentration-time curve during a dosing interval in a multiple-dose steady-state study is directly proportional to the fraction of the dose absorbed and is equal to the corresponding "zero to infinity" area under the curve for a single-dose study. Therefore, when steady-state conditions are achieved, a comparison of blood concentrations during a dosing interval may be used to define the fraction of the active drug ingredient or therapeutic moiety absorbed.

(3) Other methods based on valid scientific reasons should be used to determine the bioavailability of a drug product having dose-dependent kinetics (non-linear system).

(f) *Measurement of an acute pharmacological effect.* When comparison of the test product and the reference material is to be based on acute pharma-

cological effect-time curves, measurements of this effect should be made with sufficient frequency to demonstrate a maximum effect and a lack of significant difference between the test product and the reference material.

§ 320.28 Correlation of bioavailability with an acute pharmacological effect or clinical evidence.

Correlation of in vivo bioavailability data with an acute pharmacological effect or clinical evidence of safety and effectiveness may be required if needed to establish the clinical significance of a special claim, e.g., in the case of a controlled release preparation.

§ 320.29 Analytical methods for an in vivo bioavailability study.

(a) The analytical method used in an in vivo bioavailability study to measure the concentration of the active drug ingredient or therapeutic moiety, or its metabolite(s), in body fluids or excretory products, or the method used to measure an acute pharmacological effect shall be demonstrated to be accurate and of sufficient sensitivity to measure, with appropriate precision, the actual concentration of the active drug ingredient or therapeutic moiety, or its metabolite(s), achieved in the body.

(b) When the analytical method is not sensitive enough to measure accurately the concentration of the active drug ingredient or therapeutic moiety, or its metabolite(s), in body fluids or excretory products produced by a single dose of the test product, two or more single doses may be given together to produce higher concentration if the requirements of § 320.31 are met.

§ 320.30 Inquiries regarding bioavailability and bioequivalence requirements and review of protocols by the Food and Drug Administration.

(a) The Commissioner of Food and Drugs strongly recommends that, to avoid the conduct of an improper study and unnecessary human research, any person planning to conduct a bioavailability or bioequivalence study submit the proposed protocol for the study to FDA for review prior to the initiation of the study.

§ 320.31

21 CFR Ch. I (4-1-01 Edition)

(b) FDA may review a proposed protocol for a bioavailability or bioequivalence study and will offer advice with respect to whether the following conditions are met:

(1) The design of the proposed bioavailability or bioequivalence study is appropriate.

(2) The reference material to be used in the bioavailability or bioequivalence study is appropriate.

(3) The proposed chemical and statistical analytical methods are adequate.

(c)(1) General inquiries relating to in vivo bioavailability requirements and methodology shall be submitted to the Food and Drug Administration, Center for Drug Evaluation and Research, Division of Biopharmaceutics (HFD-420), 5600 Fishers Lane, Rockville, MD 20857.

(2) General inquiries relating to bioequivalence requirements and methodology shall be submitted to the Food and Drug Administration, Center for Drug Evaluation and Research, Division of Bioequivalence (HFD-650), 5600 Fishers Lane, Rockville, MD 20857.

[57 FR 18000, Apr. 28, 1992]

§ 320.31 Applicability of requirements regarding an "Investigational New Drug Application."

(a) Any person planning to conduct an in vivo bioavailability or bioequivalence study in humans shall submit an "Investigational New Drug Application" (IND) if:

(1) The test product contains a new chemical entity as defined in § 314.108(a) of this chapter; or

(2) The study involves a radioactively labeled drug product; or

(3) The study involves a cytotoxic drug product.

(b) Any person planning to conduct a bioavailability study in humans using a drug product that contains an already approved, non-new chemical entity shall submit an IND if the study is one of the following:

(1) A single-dose study in normal subjects or patients where either the maximum single or total daily dose exceeds that specified in the labeling of the drug product that is the subject of an approved new drug application or abbreviated new drug application.

(2) A multiple-dose study in normal subjects or patients where either the

single or total daily dose exceeds that specified in the labeling of the drug product that is the subject of an approved new drug application or abbreviated new drug application.

(3) A multiple-dose study on a controlled release product on which no single-dose study has been completed.

(c) The provisions of parts 50, 56, and 312 of this chapter are applicable to any bioavailability or bioequivalence study in humans conducted under an IND.

(d) A bioavailability or bioequivalence study in humans other than one described in paragraphs (a) through (c) of this section is exempt from the requirements of part 312 of this chapter if the following conditions are satisfied:

(1) If the study is one described under § 320.38(b) or § 320.63, the person conducting the study, including any contract research organization, shall retain reserve samples of any test article and reference standard used in the study and release the reserve samples to FDA upon request, in accordance with, and for the period specified in, § 320.38; and

(2) An in vivo bioavailability or bioequivalence study in humans shall be conducted in compliance with the requirements for institutional review set forth in part 56 of this chapter, and informed consent set forth in part 50 of this chapter.

[57 FR 18000, Apr. 28, 1992, as amended at 58 FR 25927, Apr. 28, 1993]

§ 320.32 Procedures for establishing or amending a bioequivalence requirement.

(a) The Food and Drug Administration, on its own initiative or in response to a petition by an interested person, may propose and promulgate a regulation to establish a bioequivalence requirement for a product not subject to section 505(j) of the act if it finds there is well-documented evidence that specific pharmaceutical equivalents or pharmaceutical alternatives intended to be used interchangeably for the same therapeutic effect:

(1) Are not bioequivalent drug products; or