

Substances	Limitations
<p>Polyethylenimine reaction product with 1,2-dichloroethane (CAS Reg.No. 68130-97-2) is the reaction product of homopolymerization of ethylenimine in aqueous hydrochloric acid at 100 °C and of cross-linking with 1,2-dichloroethane. The finished polymer has an average molecular weight of 50,000 to 70,000 as determined by gel permeation chromatography. The analytical method is entitled "Methodology for Molecular Weight Detection of Polyethylenimine," which is incorporated by reference in accordance with 5 U.S.C. 552(a) and 1 CFR part 51. Copies may be obtained from the Division of Petition Control, Center for Food Safety and Applied Nutrition (HFS-200), 200 C St. SW., Washington, DC 20204, and may be examined at the Center for Food Safety and Applied Nutrition's Library, 200 C St. SW., rm. 3321, Washington, DC, or at the Office of the Federal Register, 800 North Capitol St. NW., suite 700, Washington, DC.</p>	<p>May be used as a fixing material in the immobilization of glucoamylase enzyme preparations from <i>Aspergillus niger</i> for use in the manufacture of beer.</p> <p>May be used as a fixing material in the immobilization of:</p> <ol style="list-style-type: none"> 1. Glucose isomerase enzyme preparations for use in the manufacture of high fructose corn syrup, in accordance with § 184.1372 of this chapter. 2. Glucoamylase enzyme preparations from <i>Aspergillus niger</i> for use in the manufacture of beer. Residual ethylenimine in the finished polyethylenimine polymer will be less than 1 part per million as determined by gas chromatography-mass spectrometry. The residual ethylenimine is determined by an analytical method entitled "Methodology for Ethylenimine Detection in Polyethylenimine," which is incorporated by reference in accordance with 5 U.S.C. 552(a) and 1 CFR part 51. Residual 1,2-dichloroethane in the finished polyethylenimine polymer will be less than 1 part per million as determined by gas chromatography. The residual 1,2-dichloroethane is determined by an analytical method entitled, "Methodology for Ethylenedichloride Detection in Polyethylenimine," which is incorporated by reference in accordance with 5 U.S.C. 552(a) and 1 CFR part 51. Copies may be obtained from the Division of Petition Control, Center for Food Safety and Applied Nutrition (HFS-215), 200 C St. SW., Washington, DC 20204, or may be examined at the Center for Food Safety and Applied Nutrition's Library, 200 C St. SW., rm. 3321, Washington, DC, or the Office of the Federal Register, 800 North Capitol St. NW., suite 700, Washington, DC.

(b) The fixed enzyme preparation is washed to remove residues of the fixing materials.

[48 FR 5716, Feb. 8, 1983, as amended at 52 FR 39512, Oct. 22, 1987; 55 FR 12172, Apr. 2, 1990; 59 FR 36937, July 20, 1994; 61 FR 4873, Feb. 9, 1996; 61 FR 14245, Apr. 1, 1996]

§ 173.360 Octafluorocyclobutane.

The food additive octafluorocyclobutane may be safely used as a propellant and aerating agent in foamed or sprayed food products in accordance with the following conditions:

(a) The food additive meets the following specifications:

99.99 percent octafluorocyclobutane.
Less than 0.1 part per million fluoroolefins, calculated as perfluoroisobutylene.

(b) The additive is used or intended for use alone or with one or more of the following substances: Carbon dioxide, nitrous oxide, and propane, as a propellant and aerating agent for foamed or sprayed food products, except for those standardized foods that do not provide for such use.

(c) To assure safe use of the additive:

(1) The label of the food additive container shall bear, in addition to the other information required by the act, the following:

(i) The name of the additive, octafluorocyclobutane.

(ii) The percentage of the additive present in the case of a mixture.

(iii) The designation "food grade".

(2) The label or labeling of the food additive container shall bear adequate directions for use.

§ 173.370 Peroxyacids.

Peroxyacids may be safely used in accordance with the following prescribed conditions:

(a) The additive is a mixture of peroxyacetic acid, octanoic acid, acetic acid, hydrogen peroxide, peroxyoctanoic acid, and 1-hydroxyethylidene-1,1-diphosphonic acid.

(b) The additive is used as an antimicrobial agent on red meat carcasses in accordance with current industry practice where the maximum concentration of peroxyacids is 220 parts per million (ppm) as peroxyacetic acid and the maximum concentration of hydrogen peroxide is 75 ppm.

(c) The concentrations of peroxyacids and hydrogen peroxide in the additive are determined by a method entitled "Hydrogen Peroxide and Peracid (as Peracetic Acid) Content," dated July 26, 2000, developed by Ecolab, Inc., which is incorporated by reference. The Director of the Office of the Federal Register approves this incorporation by reference in accordance with 5