

a temperature between 1 and 6 °C immediately after the platelets are separated.

[38 FR 32089, Nov. 20, 1973, as amended at 42 FR 59878, Nov. 22, 1977; 43 FR 34460, Aug. 4, 1978; 49 FR 23834, June 8, 1984; 50 FR 4138, Jan. 29, 1985; 55 FR 11013, Mar. 26, 1990; 64 FR 45372, Aug. 19, 1999; 66 FR 1836, Jan. 10, 2001]

#### § 640.5 Testing the blood.

All laboratory tests shall be made on a pilot sample specimen of blood taken from the donor at the time of collecting the unit of blood, and these tests shall include the following:

(a) *Serological test for syphilis.* Whole Blood shall be negative to a serological test for syphilis.

(b) *Determination of blood group.* Each container of Whole Blood shall be classified as to ABO blood group. At least two blood group tests shall be made and the unit shall not be issued until grouping tests by different methods or with different lots of antisera are in agreement. Only those Anti-A and Anti-B Blood Grouping Reagents licensed under, or that otherwise meet the requirements of, the regulations of this subchapter shall be used, and the technique used shall be that for which the serum is specifically designed to be effective.

(c) *Determination of the Rh factors.* Each container of Whole Blood shall be classified as to Rh type on the basis of tests done on the pilot sample. The label shall indicate the extent of typing and the results of all tests performed. If the test, using Anti-D Blood Grouping Reagent, is positive, the container may be labeled "Rh Positive". If this test is negative, the results shall be confirmed by further testing which may include tests for the Rh<sub>0</sub> variant (D<sup>u</sup>) and for other Rh-Hr factors. Blood may be labeled "Rh Negative" if negative to tests for the Rh<sub>0</sub> (D) and Rh<sub>0</sub> variant (D<sup>u</sup>) factors. If the test using Anti-D Blood Grouping Reagent is negative, but not tested for the Rh<sub>0</sub> variant (D<sup>u</sup>), the label must indicate that this test was not done. Only Anti-Rh Blood Grouping Reagents licensed under, or that otherwise meet the requirements of, the regulations of this

subchapter shall be used, and the technique used shall be that for which the serum is specifically designed to be effective.

(d) *Sterility test.* Whole Blood intended for transfusion shall not be tested for sterility by a method that entails entering the final container before the blood is used for transfusion.

(e) *Inspection.* Whole Blood shall be inspected visually during storage and immediately prior to issue. If the color or physical appearance is abnormal or there is any indication or suspicion of microbial contamination the unit of Whole Blood shall not be issued for transfusion.

(f) *Test for antibody to HIV.* Whole Blood shall be tested for antibody to HIV as prescribed in §610.45 of this chapter.

[38 FR 32089, Nov. 20, 1973, as amended at 50 FR 4138, Jan. 29, 1985; 53 FR 117, Jan. 5, 1988; 53 FR 12764, Apr. 19, 1988; 64 FR 45372, Aug. 19, 1999; 66 FR 1836, Jan. 10, 2001]

#### § 640.6 Modifications of Whole Blood.

Upon approval by the Director, Center for Biologics Evaluation and Research, of a supplement to the biologics license application for Whole Blood a manufacturer may prepare Whole Blood from which the antihemophilic factor has been removed, provided the Whole Blood meets the applicable requirements of this subchapter and the following conditions are met:

(a) The antihemophilic factor shall be removed in accordance with paragraphs (a), (b), and (c) of § 640.52.

(b) Although the closed system between the red blood cells and plasma shall be maintained, the red blood cells shall be maintained between 1 and 6° C at all times, including that time when the plasma is being frozen for removal of the antihemophilic factor.

[38 FR 32089, Nov. 20, 1973, as amended at 49 FR 23834, June 8, 1984; 50 FR 4138, Jan. 29, 1985; 55 FR 11013, Mar. 26, 1990; 59 FR 49351, Sept. 28, 1994; 64 FR 45372, Aug. 19, 1999; 64 FR 56453, Oct. 20, 1999]