

Monogenic Forms of Diabetes: Neonatal Diabetes Mellitus and Maturity-onset Diabetes of the Young

National Diabetes Information Clearinghouse



National
Institute of
Diabetes and
Digestive
and Kidney
Diseases

NATIONAL
INSTITUTES
OF HEALTH

The most common forms of diabetes, type 1 and type 2, are polygenic, meaning the risk of developing these forms of diabetes is related to multiple genes. Environmental factors, such as obesity in the case of type 2 diabetes, also play a part in the development of polygenic forms of diabetes. Polygenic forms of diabetes often run in families. Doctors diagnose polygenic forms of diabetes by testing blood glucose in individuals with risk factors or symptoms of diabetes.

Genes provide the instructions for making proteins within the cell. If a gene has a mutation, the protein may not function properly. Genetic mutations that cause diabetes affect proteins that play a role in the ability of the body to produce insulin or in the ability of insulin to lower blood glucose. People have two copies of most genes; one gene is inherited from each parent.

Monogenic Forms of Diabetes

Some rare forms of diabetes result from mutations in a single gene and are called monogenic. Monogenic forms of diabetes account for about 1 to 5 percent of all cases of diabetes in young people. In most cases of monogenic diabetes, the gene mutation is inherited; in the remaining cases the gene

mutation develops spontaneously. Most mutations in monogenic diabetes reduce the body's ability to produce insulin, a protein produced in the pancreas that helps the body use glucose for energy. Neonatal diabetes mellitus (NDM) and maturity-onset diabetes of the young (MODY) are the two main forms of monogenic diabetes. MODY is much more common than NDM. NDM first occurs in newborns and young infants; MODY usually first occurs in children or adolescents but may be mild and not detected until adulthood.

Genetic testing can diagnose most forms of monogenic diabetes. If genetic testing is not performed, people with monogenic diabetes may appear to have one of the polygenic forms of diabetes. When hyperglycemia is first detected in adulthood, type 2 is often diagnosed instead of monogenic diabetes. Some monogenic forms of diabetes can be treated with oral diabetes medications while other forms require insulin injections. A correct diagnosis that allows the proper treatment to be selected should lead to better glucose control and improved health in the long term. Testing of other family members may also be indicated to determine whether they are at risk for diabetes.


What is neonatal diabetes mellitus (NDM)?

NDM is a monogenic form of diabetes that occurs in the first 6 months of life. It is a rare condition occurring in only one in 100,000 to 500,000 live births. Infants with

For more information about diabetes, see *Diabetes Overview*, which is available from the National Diabetes Information Clearinghouse. See page 5 for contact information.



U.S. Department
of Health and
Human Services



Genes affect a person's risk of developing diabetes.

NDM do not produce enough insulin, leading to an increase in blood glucose. NDM can be mistaken for the much more common type 1 diabetes, but type 1 diabetes usually occurs later than the first 6 months of life. In about half of those with NDM, the condition is lifelong and is called permanent neonatal diabetes mellitus (PNDM). In the rest of those with NDM, the condition is transient and disappears during infancy but can reappear later in life; this type of NDM is called transient neonatal diabetes mellitus (TNDM). Specific genes that can cause NDM have been identified. More information about each type of NDM is provided in the appendix.


Symptoms of NDM include thirst, frequent urination, and dehydration. NDM can be diagnosed by finding elevated levels of glucose in blood or urine. In severe cases, the deficiency of insulin may cause the body to produce an excess of acid, resulting in a

potentially life-threatening condition called ketoacidosis. Most fetuses with NDM do not grow well in the womb and newborns are much smaller than those of the same gestational age, a condition called intrauterine growth restriction. After birth, some infants fail to gain weight and grow as rapidly as other infants of the same age and sex. Appropriate therapy improves and may normalize growth and development.

What is maturity-onset diabetes of the young (MODY)?

MODY is a monogenic form of diabetes that usually first occurs during adolescence or early adulthood. However, MODY sometimes remains undiagnosed until later in life. A number of different gene mutations have been shown to cause MODY, all of which limit the ability of the pancreas to produce insulin. This process leads to the high blood glucose levels characteristic of diabetes and, in time, may damage body tissues, particularly the eyes, kidneys, nerves, and blood vessels. MODY accounts for about 1 to 5 percent of all cases of diabetes in the United States. Family members of people with MODY are at greatly increased risk for the condition.

People with MODY may have only mild or no symptoms of diabetes and their hyperglycemia may only be discovered during routine blood tests. MODY may be confused with type 1 or type 2 diabetes. People with MODY are generally not overweight and do not have other risk factors for type 2 diabetes, such as high blood pressure or abnormal blood fat levels. While both type 2 diabetes and MODY can run in families, people with MODY typically have a family history of diabetes in multiple successive generations, meaning that MODY is



Each child of a parent with MODY has a 50 percent chance of inheriting the disease.

present in a grandparent, a parent, and a child. Unlike people with type 1 diabetes who always require insulin, people with MODY can often be treated with oral diabetes medications. Treatment varies depending on the genetic mutation that has caused the MODY. More information about each type of MODY is provided in the appendix.

What do I need to know about genetic testing and counseling?

Testing for monogenic diabetes involves providing a blood sample from which DNA is isolated. The DNA is analyzed for changes in the genes that cause monogenic diabetes. Abnormal results can determine the gene responsible for diabetes in a particular individual or show whether someone is likely to develop a monogenic form of

diabetes in the future. Genetic testing can also be helpful in selecting the most appropriate treatment for individuals with monogenic diabetes. Prenatal testing can diagnose these conditions in unborn children.

Most forms of monogenic diabetes are caused by dominant mutations, meaning that the condition can be passed on to children when only one parent is affected. In contrast, if the mutation is a recessive mutation, a disease gene must be inherited from both parents for diabetes to occur. For recessive forms of monogenic diabetes, testing can indicate whether parents or siblings without disease are carriers for recessive genetic conditions that could be inherited by their children.

If you suspect that you or a member of your family may have a monogenic form of diabetes, you should seek help from health care professionals—physicians and genetic counselors—who have specialized knowledge and experience in this area. They can determine whether genetic testing is appropriate, select the genetic tests that should be performed, and provide information about the basic principles of genetics, genetic testing options, and confidentiality issues. They also can review the test results with the patient or parent after testing, make recommendations about how to proceed, and discuss testing options for other family members.

Hope Through Research

Researchers are studying the genetic causes of and metabolic processes related to diabetes. Discoveries about monogenic forms of diabetes may contribute to the search for the causes of and treatments for type 1 and type 2 diabetes. For information about clinical trials related to diabetes and genetics, see www.ClinicalTrials.gov.

Points to Remember

- Mutations in single genes can cause rare forms of diabetes.
- Genetic testing can identify many forms of monogenic diabetes.
- A physician evaluates whether genetic testing is appropriate.
- A correct diagnosis aided by genetic testing can lead to optimal treatment.
- Recent research results show that people with certain forms of monogenic diabetes can be treated with oral diabetes medications instead of insulin injections.

For More Information

Information About Genetic Testing, Evaluation, and Counseling, Funded by the National Institutes of Health (NIH)

The GeneTests website (www.genetests.org) provides information about medical genetics, an international directory of genetic testing laboratories, and an international directory of genetics clinics providing genetic evaluation and genetic counseling.

Information About Genetics

National Human Genome Research Institute
Genetic and Rare Diseases Information Center
P.O. Box 8126
Gaithersburg, MD 20898-8126
Phone: 1-888-205-2311
Fax: 240-632-9164
Email: gardinfo@nih.gov
Internet: www.genome.gov/health

The Genetics Home Reference website (www.ghr.nlm.nih.gov) provides consumer-friendly information about genetic conditions. A service of the National Library of Medicine (NLM), NIH.

The Online Mendelian Inheritance in Man database (www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=OMIM) is a catalog of human genes and genetic disorders with references and related links. A service of the National Center for Biotechnology Information, NLM, NIH.

Entrez Gene (www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=gene) is a searchable database of genes with extensive information and related links. A service of the National Center for Biotechnology Information, NLM, NIH.

The Diabetes Research department and the Centre for Molecular Genetics at the Peninsula Medical School and Royal Devon and Exeter Hospital, Exeter, United Kingdom (www.diabetesgenes.org) provides information for patients and health care professionals about genetic forms of diabetes.

The International Society for Pediatric and Adolescent Diabetes (www.ispad.org) is an international society for health care professionals and others interested in childhood diabetes. They publish consensus guidelines; see the list of selected references on page 10.

Information About Diabetes

National Diabetes Information

Clearinghouse

1 Information Way

Bethesda, MD 20892-3560

Phone: 1-800-860-8747

Fax: 703-738-4929

Email: ndic@info.niddk.nih.gov

Internet: www.diabetes.niddk.nih.gov

National Diabetes Education Program

1 Diabetes Way

Bethesda, MD 20892-3560

Phone: 1-800-438-5383

Fax: 703-738-4929

Email: ndep@mail.nih.gov

Internet: www.ndep.nih.gov

American Diabetes Association

National Call Center

1701 North Beauregard Street

Alexandria, VA 22311-1742

Phone: 1-800-DIABETES (342-2383)

Fax: 703-549-6995

Email: AskADA@diabetes.org

Internet: www.diabetes.org

Juvenile Diabetes Research Foundation International

120 Wall Street

New York, NY 10005-4001

Phone: 1-800-533-CURE (2873)

Fax: 212-785-9595

Email: info@jdrf.org

Internet: www.jdrf.org

You may also find additional information on this topic using the following databases:

The NIDDK Reference Collection is a collection of thousands of materials produced for patients and health care professionals, including fact sheets, brochures, and audiovisual materials. Visit www.catalog.niddk.nih.gov/resources.

MedlinePlus brings together a wealth of information from the National Library of Medicine, the National Institutes of Health, and other government agencies and health-related organizations. MedlinePlus offers easy access to medical journal articles, a medical dictionary and medical encyclopedia, health information in Spanish, hospital and physician directories, drug and supplement lists, interactive patient tutorials, links to hundreds of clinical trials, and the latest health news. Visit www.medlineplus.gov.

The U.S. Government does not endorse or favor any specific commercial product or company. Trade, proprietary, or company names appearing in this document are used only because they are considered necessary in the context of the information provided. If a product is not mentioned, the omission does not mean or imply that the product is unsatisfactory.

Appendix: Characteristics of Monogenic Forms of Diabetes

Type of Diabetes	Gene or Syndrome*	Affected Protein	How Common	Usual Age of Onset	Type of Inheritance or Mutation**	Causes Intrauterine Growth Restriction?	Transient or Permanent?***	Treatment
Neonatal Diabetes Mellitus (NDM)			Rare; occurs in about one of every 100,000 to 500,000 live births					
Permanent Neonatal Diabetes Mellitus (PNDM)			50% of all cases of NDM					
PNDM	<i>KCNJ11</i>	Kir6.2	Most common type of PNDM	3 to 6 months	Autosomal dominant (10%) Spontaneous	Yes	Permanent (This gene also causes a transient form of NDM; see TNDM section on page 7)	Treated with insulin in the past but often can be treated with oral sulfonylureas
PNDM	<i>ABCC8</i>	SUR1—sulfonylurea receptor 1	Rare	1 to 3 months	Autosomal dominant (12% of NDM) Spontaneous	No	Permanent (This gene also causes a transient form of NDM; see TNDM section on page 7)	Treated with insulin in the past but often can be treated with oral sulfonylureas
PNDM	<i>GCK</i>	glucokinase	Rare	1 week	Autosomal recessive	Yes	Permanent	Insulin
PNDM	<i>IPF1</i> ; also known as <i>PDX1</i>	insulin promoter factor 1	Rare	1 week	Autosomal recessive	Yes	Permanent	Treat to replace endocrine and exocrine pancreas functions
PNDM	<i>PTF1A</i>	pancreas transcription factor 1 A	Rare	At birth	Autosomal recessive	Yes	Permanent	Treat to replace endocrine and exocrine pancreas functions
PNDM	<i>FOXP3</i> , IPEX syndrome	forkhead box P3	Rare	Sometimes present at birth	X-linked	Yes	Permanent	Insulin
PNDM	<i>EIF2AK3</i> , Wolcott-Rallison syndrome	eukaryotic translation initiation factor 2-alpha kinase 3	Rare	3 months	Autosomal recessive	Yes	Permanent	Insulin and treatment for associated conditions

Appendix: Characteristics of Monogenic Forms of Diabetes (continued)

Type of Diabetes	Gene or Syndrome*	Affected Protein	How Common	Usual Age of Onset	Type of Inheritance or Mutation**	Causes Intrauterine Growth Restriction?	Transient or Permanent?***	Treatment
Transient Neonatal Diabetes Mellitus (TNDM)			50% of all cases of NDM					
TNDM	<i>ZAC/HYMAI</i>	<i>ZAC</i> : pleomorphic adenoma gene-like 1 or <i>PLAG1</i> <i>HYMAI</i> : hydatiform mole-associated and imprinted transcript	Most common form of NDM	Birth to 3 months	Autosomal dominant Spontaneous	Yes	Transient	Initially, treat with insulin; reduce dosage as needed; when diabetes recurs, treat with diet modification and physical activity; may also require insulin
TNDM	<i>ABCC8</i>	<i>SUR1</i> —sulfonylurea receptor 1	Rare	Birth to 6 months	Autosomal dominant Spontaneous	Varies	Transient (This gene also causes a permanent form of NDM; see PNDM section on page 6)	Oral sulfonylureas
TNDM	<i>KCNJ11</i>	Kir6.2	Uncommon cause of TNDM but most common cause of PNDM	Birth to 6 months	Autosomal dominant Spontaneous	Yes	Transient (This gene also causes a permanent form of NDM; see PNDM section on page 6)	Oral sulfonylureas
TNDM	<i>HNF1β</i> (beta); also known as <i>HNF1B</i>	hepatocyte nuclear factor 1B	Rare	Birth to 6 months	Autosomal dominant (60%) Spontaneous	Yes	Transient	Insulin

Appendix: Characteristics of Monogenic Forms of Diabetes (continued)

Type of Diabetes	Gene or Syndrome*	Affected Protein	How Common	Usual Age of Onset	Type of Inheritance or Mutation**	Causes Intrauterine Growth Restriction?	Transient or Permanent?***	Treatment
Maturity-onset Diabetes of the Young (MODY)			1 to 5% of all cases of diabetes in the United States					
MODY 1	<i>HNF4A</i>	hepatocyte nuclear factor 4 α (alpha)	Rare	Adolescence or early adulthood	Autosomal dominant	No	Permanent	For most, oral sulfonylureas; some patients may need insulin
MODY 2	<i>GCK</i>	glucokinase	MODY 2 and MODY 3 account for about two-thirds of all cases of MODY MODY 2 is the second-most common form of MODY	Mild hyperglycemia may be present at birth; otherwise, early childhood	Autosomal dominant	Lower than normal birthweight can occur	Permanent	Diet modification and physical activity; medications usually not required; some patients do not require any treatment during childhood
MODY 3	<i>TCF1</i>	hepatic nuclear factor 1 α (alpha) or HNF1 α (alpha) or <i>HNF1A</i>	MODY 3 is the most common form of MODY	Adolescence or early adulthood	Autosomal dominant	No	Permanent	Initially, treat with diet modification; can be treated with oral sulfonylureas; some patients may need insulin
MODY 4	<i>IPF1</i> ; also known as <i>PDX1</i>	insulin promoter factor 1	Rare	Early adulthood; can present later	Autosomal dominant	No	Permanent	Oral sulfonylureas; some patients may need insulin

Appendix: Characteristics of Monogenic Forms of Diabetes (continued)

Type of Diabetes	Gene or Syndrome*	Affected Protein	How Common	Usual Age of Onset	Type of Inheritance or Mutation**	Causes Intrauterine Growth Restriction?	Transient or Permanent?***	Treatment
MODY 5	<i>TCF2</i>	hepatic nuclear factor 1 β (beta) or <i>HNF1B</i>	Rare	Adolescence or early adulthood	Autosomal dominant	No	Permanent	Insulin; patients also may need treatment for related conditions such as kidney failure or cysts
MODY 6	<i>NeuroD1</i> , or <i>BETA2</i>	neurogenic differentiation factor 1	Rare	In the fourth decade of life	Autosomal dominant	No	Permanent	Insulin

* Gene or Syndrome: the name of the gene with the mutation or the syndrome—a grouping of conditions that occur together and indicate a specific disease—caused by the mutated gene

** Type of Inheritance or Mutation:

- **Autosomal dominant.** Normally, every cell has two copies of each gene—one that comes from the mother and one from the father. An autosomal dominant inheritance pattern means that a mutation happens in only one copy of the gene, and a parent with a mutation can pass on a copy of their working gene or a copy of their damaged gene. In autosomal dominant inheritance, a child who has a parent with a mutation has a 50% chance of inheriting that mutation.
- **Autosomal recessive.** Normally, every cell has two copies of each gene—one that comes from the mother and one from the father. An autosomal recessive inheritance pattern means a mutation must be present in both copies of the gene in order for a person to be affected, and each parent must pass on a gene mutation for a child to be affected. If a person only has one copy of the gene mutation, that person is called a carrier. If both parents are carriers of a recessive gene mutation, each child has a 25% chance of being affected.
- **Spontaneous.** A new mutation, or change, in a gene
- **X-linked.** When a trait or a disease occurs in a person who has inherited a mutated gene on the X chromosome, one of the sex chromosomes

*** Transient or Permanent: whether the form of diabetes goes away after some time, called transient, or is permanent

Selected References

- Babenko AP, Polak M, Cavé H, Busiah K, Czernichow P, Scharfmann R, Bryan J, Aguilar-Bryan L, Vaxillaire M, Froguel P. Activating mutations in the *ABCC8* gene in neonatal diabetes mellitus. *New England Journal of Medicine*. 2006;355(5):456–466.
- Colombo C, Delvecchio M, Zecchino C, Falenza MF, Cavallo L, Barbetti F. Transient neonatal diabetes mellitus is associated with a recurrent (R201H) *KCNJ11* (Kir6.2) mutation. *Diabetologia*. 2005;48:2439–2441.
- Craig ME, Hattersley A, Donaghue K. International Society for Pediatric and Adolescent Diabetes (ISPAD) Clinical Practice Consensus Guidelines 2006–2007. Definition, epidemiology and classification. *Pediatric Diabetes*. 2006;7:343–351.
- Fajans SS, Bell GI, Polonsky KS. Molecular mechanisms and clinical pathophysiology of maturity-onset diabetes of the young. *New England Journal of Medicine*. 2001;345(13):971–980.
- Gloyn AL, Pearson ER, Antcliff JF, Proks P, Bruining GJ, Slingerland AS, Howard N, Srinivasan S, Silva JM, Molnes J, Edghill EL, Frayling TM, Temple IK, Mackay D, Shield JP, Sumnik Z, van Rhijn A, Wales JK, Clark P, Gorman S, Aisenberg J, Ellard S, Njølstad PR, Ashcroft FM, Hattersley AT. Activating mutations in the gene encoding the ATP-sensitive potassium-channel subunit Kir6.2 and permanent neonatal diabetes. *New England Journal of Medicine*. 2004;350(18):1838–1849.
- Hattersley A, Bruining J, Shield J, Njølstad P, Donaghue K. International Society for Pediatric and Adolescent Diabetes (ISPAD) Clinical Practice Consensus Guidelines 2006–2007. The diagnosis and management of monogenic diabetes in children. *Pediatric Diabetes*. 2006;7:352–360.
- Hattersley AT. Beyond the beta cell in diabetes. *Nature Genetics*. 2006;38(1):12–13.
- Hattersley AT, Ashcroft FM. Activating mutations in Kir6.2 and neonatal diabetes: new clinical syndromes, new scientific insights, and new therapy. *Diabetes*. 2005;54:2503–2513.

- Hattersley AT, Pearson, ER. Minireview: pharmacogenetics and beyond: the interaction of therapeutic response, beta-cell physiology, and genetics in diabetes. *Endocrinology*. 2006;147:2657–2663.
- National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases. The genetic landscape of diabetes. Available at: www.ncbi.nlm.nih.gov/books/bv.fcgi?call=bv.View..ShowTOC&rid=diabetes.TOC&depth=2. Posted 2004. Accessed January 2, 2007.
- Pearson ER, Flechtner I, Njølstad PR, Malecki MT, Flanagan SE, Larkin B, Ashcroft FM, Klimes I, Codner E, Iotova V, Slingerland AS, Shield J, Robert JJ, Holst JJ, Clark PM, Ellard S, Søvik O, Polak M, Hattersley AT for the Neonatal Diabetes International Collaborative Group. Switching from insulin to oral sulfonylureas in patients with diabetes due to Kir6.2 mutations. *New England Journal of Medicine*. 2006;355(5):467–477.
- Ræder H, Johansson S, Holm PI, Haldorsen IS, Mas E, Sbarra V, Neramoen I, Eide SA, Grevle L, Bjorkhaug L, Sagen JV, Aksnes L, Søvik O, Lombardo D, Molven A, Njølstad PR. Mutations in the CEL VNTR cause a syndrome of diabetes and pancreatic exocrine dysfunction. *Nature Genetics*. 2006;38(1):54–62.
- Sperling MA. ATP-sensitive potassium channels—neonatal diabetes mellitus and beyond. *New England Journal of Medicine*. 2006;355(5):507–510.
- Sperling MA. Neonatal diabetes mellitus: from understudy to center stage. *Current Opinion in Pediatrics*. 2005;17:512–518.
- Vaxillaire M, Froguel P. Genetic basis of maturity-onset diabetes of the young. *Endocrinology and Metabolism Clinics of North America*. 2006;35:371–384.

National Diabetes Information Clearinghouse

1 Information Way
Bethesda, MD 20892-3560
Phone: 1-800-860-8747
Fax: 703-738-4929
Email: ndic@info.niddk.nih.gov
Internet: www.diabetes.niddk.nih.gov

The National Diabetes Information Clearinghouse (NDIC) is a service of the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK). The NIDDK is part of the National Institutes of Health of the U.S. Department of Health and Human Services. Established in 1978, the Clearinghouse provides information about diabetes to people with diabetes and to their families, health care professionals, and the public. The NDIC answers inquiries, develops and distributes publications, and works closely with professional and patient organizations and Government agencies to coordinate resources about diabetes.

Publications produced by the Clearinghouse are carefully reviewed by both NIDDK scientists and outside experts. This publication was reviewed by Mark A. Sperling, M.D., Department of Pediatrics, Children's Hospital, University of Pittsburgh; Kenneth S. Polonsky, M.D., Department of Medicine, Washington University School of Medicine; and Concepcion R. Nierras, Ph.D., Juvenile Diabetes Research Foundation International.

This publication is not copyrighted. The Clearinghouse encourages users of this fact sheet to duplicate and distribute as many copies as desired.

This fact sheet is also available at
www.diabetes.niddk.nih.gov.



U.S. DEPARTMENT OF HEALTH
AND HUMAN SERVICES
National Institutes of Health

NIH Publication No. 07-6141
March 2007