


## Acrylonitrile

### CAS No. 107-13-1

Reasonably anticipated to be a human carcinogen  
First Listed in the *Second Annual Report on Carcinogens* (1981)



### Carcinogenicity

Acrylonitrile is *reasonably anticipated to be a human carcinogen* based on sufficient evidence of carcinogenicity in experimental animals (IARC 1979, 1982, 1987, 1999, ATSDR 1990). When administered orally (by gavage or in drinking water), acrylonitrile induced increased incidences of forestomach squamous cell papillomas, central nervous system microgliomas, mammary gland carcinomas, and Zymbal gland carcinomas in rats of both sexes. Inhalation studies demonstrated that acrylonitrile induced Zymbal gland carcinomas, glial cell tumors, forestomach papillomas and acanthomas, and central nervous system neoplasms in rats of both sexes, mammary tumors and extrahepatic angiosarcomas in female rats, and hepatocellular tumors in male rats (IARC 1999).

The potential carcinogenicity of acrylonitrile in humans has been studied in several occupational studies. An increased risk of cancer of the lung was observed in U.S. textile plant workers exposed to acrylonitrile and observed for 20 years or more (IARC 1979). Since acrylonitrile was reviewed for listing in the *Second Annual Report on Carcinogens* (1981), the International Agency for Research on Cancer (IARC) has reviewed human epidemiological studies on exposure to acrylonitrile several times (IARC 1982, 1987, 1999). An increase in cancer risk among workers exposed to acrylonitrile was suggested by studies conducted in the 1970s and 1980s (IARC 1979, 1982, 1987). Further follow-up of the U.S. textile plant workers revealed a continued excess of lung cancer and an excess of prostate cancer. Other studies of workers exposed to acrylonitrile (including textile workers and rubber workers), published in the 1980's, reported an excess of cancers at other sites including prostate, stomach, colon, brain, and the lymphatic and hematopoietic systems. IARC stated that these earlier studies were limited by inadequate exposure assessment, small sample size, insufficient length of follow-up and possible confounding (IARC 1999). In the 1990s, the results of four cohort studies (some of which included subjects from the earlier studies) conducted in the United States (two studies), United Kingdom, and the Netherlands, which involved larger number of subjects, were reported. These studies did not show an excess of risk of any type of cancer among workers exposure to acrylonitrile. However, an increased risk of lung cancer was observed in individuals exposed to the highest levels of acrylonitrile in one of the studies (IARC 1999).

### Properties

Acrylonitrile is a colorless, volatile liquid that is soluble in water and most common organic solvents such as acetone, benzene, carbon tetrachloride, ethyl acetate, and toluene. It melts at -84°C and boils at 77°C. Technical-grade acrylonitrile is more than 99% pure (IARC 1999). The technical-grade product always contains a polymerization inhibitor. Acrylonitrile is a reactive chemical that polymerizes spontaneously and can explode when exposed to flame (HSDB 2002).

### Use

Acrylonitrile is an important industrial chemical. It is used extensively in the manufacture of synthetic fibers, resins, plastics, elastomers, and rubber for a variety of consumer goods such as textiles, dinnerware, food containers, toys, luggage, automotive parts, small appliances, and telephones (SRI 1984). In 1986, about 40% of the acrylonitrile produced was used to produce acrylic and modacrylic fibers, 28% to

produce acrylonitrile-butadiene-styrene (ABS) and styrene-acrylonitrile (SAN) resins, and 15% to produce adiponitrile, an intermediate used in nylon production. The remainder was used in the production of acrylamide (10%), nitrile elastomers, barrier resins, and miscellaneous specialty chemicals (4%) (CMR1986). The manufacture of carbon fibers used for high-performance applications in the aircraft, defense, and aerospace industries is a growing specialty application of acrylonitrile. Other specialty applications include the production of fatty amines, ion exchange resins, and fatty amine amides used in cosmetics, adhesives, corrosion inhibitors, and water-treatment resins (IARC 1999). Acrylonitrile has also been used as a fumigant; however, most pesticide registrations of the chemical were canceled in 1978, and its use as a fumigant has been abandoned (ATSDR 1990).

### Production

Acrylonitrile has been produced in the United States since 1940 (IARC 1979). It has been ranked in the top 50 highest-volume chemicals by Chemical and Engineering News for the past several years (CEN 1998). U.S. production of acrylonitrile averaged 2.7 billion lb for the time period 1985 to 1987. In 1990 and 1993, 2.7 billion lb and 2.5 billion lb were produced, respectively. U.S. production of acrylonitrile increased to 3.4 billion lb in 1996 (IARC 1999). Chem Sources (2001) identified 21 suppliers of acrylonitrile in the United States in 2001.

As reported in 1990, imports of acrylonitrile had substantially decreased in the 1970s and had been considered negligible since 1984 (ATSDR 1990); for example, in 1985, only 441 lb were imported into the United States (USDOC Imports 1986). However, in 1989, the amount of acrylonitrile imported into the U.S. increased dramatically, with the reported import value greater than 400,000 lb (USDOC Imports 1990). In 2000, more than 17 million lb were imported (ITA 2001). In contrast, a sizable fraction of acrylonitrile is exported, with over 1.5 billion lb exported in 2000 (ATSDR 1990, ITA 2001). In 1989, over 942 million lb were exported (USDOC Exports 1990). This figure is similar to the 944 million lb reported for 1986 and the 943 million lb reported for 1985 (CW 1987, USDOC Exports 1986). In 1984, the value was 834 million lb (Chem. Prod. 1985).

### Exposure

The primary routes of potential human exposure to acrylonitrile are inhalation and dermal contact. Exposure to acrylonitrile may occur during its manufacture and production; greater potential for exposure exists for workers using acrylonitrile to make other products in factories where the compound is not easily contained (DPIM 1989). The National Occupational Exposure Survey (1981-1983) indicated that 51,153 total workers, including 25,320 women, were potentially exposed to acrylonitrile (NIOSH 1984). The National Occupational Hazard Survey, conducted by NIOSH from 1972 to 1974, estimated that 71,610 workers were potentially exposed to acrylonitrile in the workplace (NIOSH 1976). In a later report, NIOSH estimated that 125,000 persons have the potential for occupational exposure (NIOSH 1978). These may be overestimates because non-exposed workers within each Standard Industrial Classification may not have been excluded. Based on industrial sources, a more reasonable estimate of the number of workers exposed is 5,130 people (NCI 1985). These individuals include acrylic resin makers, synthetic organic chemists, pesticide workers, and rubber, synthetic fiber, and textile makers.

People living near chemical factories or waste sites are the likely candidates for exposure to measurable amounts of acrylonitrile in air and water. The remaining general population may be potentially exposed through consumer product usage such as acrylic carpeting or by ingestion of contaminated foods. Exposure in each case, however, is very low because of little migration of the monomer from such products (ATSDR 1990). The concentrations of acrylonitrile in consumer products are estimated to be less than 1 ppm in acrylic and modacrylic

fibers, 30 to 50 ppm in ABS copolymers, 15 ppm in SAN copolymers, and 0 to 750 ppm in nitrile rubber and latex goods (Patrianakos and Hoffman 1979). In the 1960s and 1970s, acrylonitrile was also detected in cigarette smoke, usually at levels of 1 to 2 mg per cigarette, since the chemical was used as a fumigant for stored tobacco. It is unlikely that cigarette smoking is today a major source of exposure to acrylonitrile because its use as a fumigant has been discontinued (ATSDR 1990). The presence of acrylonitrile, even as a trace contaminant, may be cause for concern. The extent or risk of any potential exposure for the general population through air emissions from production, manufacture, bulk storage, and waste disposal has not been determined. A 1977 report stated that total acrylonitrile air emissions were 2.2% of total production. The Toxic Chemical Release Inventory (TRI) estimated that 4,891,577 lb of acrylonitrile were released to the environment from 97 facilities that produced, processed, or used the chemical in the United States in 1996. Of that total, 26.5% and 73.5% were released to the air and underground injection wells, respectively (TRI96 1998). In 1999, the amount of acrylonitrile released into the environment increased to 5,466,510 lb from 117 facilities that produced, processed, or used the chemical in the United States. Of that total, 17.9% was released to the air and 81.6% to underground injection wells. Air emissions from 17 facilities, each releasing >10,000 lb, represented 92.0% of the total atmospheric release. Four facilities, each releasing >100,000 lb, accounted for 98.4% of the total underground injection release. Releases to water and land were very low with 1,068 lb and 560 lb released, respectively (TRI99 2001).

## Regulations

### DOT

Acrylonitrile is considered a hazardous material and special requirements have been set for marking, labeling, and transporting this material

### EPA

#### Clean Air Act

- NESHAP: Listed as a Hazardous Air Pollutant (HAP)
- NSPS: Manufacture of substance is subject to certain provisions for the control of Volatile Organic Compound (VOC) emissions
- Prevention of Accidental Release: Threshold Quantity (TQ) = 20,000 lb
- Urban Air Toxics Strategy: Identified as one of 33 HAPs that present the greatest threat to public health in urban areas

#### Clean Water Act

- Effluent Guidelines: Listed as a Toxic Pollutant
- Water Quality Criteria: Based on fish/shellfish and water consumption = 0.051 µg/L; based on fish/shellfish consumption only = 0.25 µg/L

#### Comprehensive Environmental Response Compensation and Liability Act

- Reportable Quantity (RQ) = 100 lb

#### Emergency Planning and Community Right-to-Know Act

- Toxics Release Inventory: Listed substance subject to reporting requirements
- Reportable Quantity (RQ) = 100 lb
- Threshold Planning Quantity (TPQ) = 10,000 lb

#### Resource Conservation and Recovery Act

- Listed Hazardous Waste: Waste codes in which listing is based wholly or partly on substance - U009, K011, K013
- Listed as a Hazardous Constituent of Waste

### FDA

Acrylonitrile copolymers and resins containing less than 30 percent acrylonitrile may be used in food packaging materials provided certain conditions in the regulation are met

### OSHA

Ceiling Concentration = 10 ppm [15 minute exposure]  
 Permissible Exposure Limit (PEL) = 2 ppm  
 "Comprehensive Standards" for occupational exposure to this substance have been developed

## Guidelines

### ACGIH

Threshold Limit Value - Time-Weighted Average Limit (TLV-TWA) = 2 ppm

### NIOSH

Ceiling Recommended Exposure Limit = 10 ppm (15 minute exposure)  
 Immediately Dangerous to Life and Health (IDLH) = 85 ppm  
 Recommended Exposure Limit (time-weighted-average workday) = 1 ppm  
 Listed as a potential occupational carcinogen

## REFERENCES

- ATSDR. 1990. Toxicological Profile for Acrylonitrile (Final Report). NTIS Accession No. PB91-180489. Atlanta, GA: Agency for Toxic Substances and Disease Registry. 140 pp.
- CEN. 1998. Production Profiles. Chem Eng News 76(26): 43.
- ChemProd. 1985. Chemical Product Synopsis: Acrylonitrile. Cortland, NY: Mansville Chemical Products Corporation.
- ChemSources. 2001. Chemical Sources International, Inc. <http://www.chemsources.com>.
- CMR. 1986. Chemical Profile: Acrylonitrile. Chem Mark Report 229(11): 58.
- CV. 1987. Chem Week 140(6): 26.
- DPIM. 1989. Acetaldehyde. In Dangerous Properties of Industrial Materials Report, vol. 9. J. C. Prager, ed. New York, NY: Van Nostrand Reinhold Company, Inc. p. 30-45.
- HSDB. 2002. Hazardous Substances Database. National Library of Medicine. <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>.
- IARC. 1979. Some Monomers, Plastics, and Synthetic Elastomers, and Acrolein. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, vol. 19. Lyon, France: International Agency for Research on Cancer. 513 pp.
- IARC. 1982. Chemicals, Industrial Processes and Industries Associated with Cancer in Humans. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, Supplement 4. Lyon, France: International Agency for Research on Cancer. 292 pp.
- IARC. 1987. Overall Evaluations of Carcinogenicity. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, Supplement 7. Lyon, France: International Agency for Research on Cancer. 440 pp.
- IARC. 1999. Re-evaluation of Some Organic Chemicals, Hydrazine, and Hydrogen Peroxide. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, vol. 71. Lyon, France: International Agency for Research on Cancer. 1589 pp.
- ITA. 2001. U.S. Trade Quick-Reference Tables: December 2002 Imports. Subheading 292610: Acrylonitrile. International Trade Administration. U.S. Department of Commerce. <http://www.ita.doc.gov/td/industry/otea/Trade-Detail/>.
- NCI. 1985. Monograph on Human Exposure to Chemicals in the Workplace: Acrylonitrile. Technical Report No. 85-192. Bethesda, MD: Department of Health and Human Services.
- NIOSH. 1976. National Occupational Hazard Survey (1972-74). Cincinnati, OH: Department of Health, Education and Welfare.
- NIOSH. 1978. A Recommended Standard for Occupational Exposure to Acrylonitrile. Cincinnati, OH: Department of Health, Education and Welfare. 14.
- NIOSH. 1984. National Occupational Exposure Survey (1981-83). Cincinnati, OH: U. S. Department of Health and Human Services. <http://www.cdc.gov/noes/noes3/empl0003.html>.
- Patrianakos, C. and D. Hoffmann. 1979. Chemical studies in tobacco smoke LXIV. On the analysis of aromatic amines in cigarette smoke. J Anal Toxicol 3: 50.
- SRI. 1984. Acrylonitrile. In Chemical Economics Handbook: Organic Chemicals. Menlo Park, CA: SRI International.
- TRI96. 1998. Toxic Chemical Release Inventory 1996. Data contained in the Toxic Chemical Release Inventory (TRI). National Library of Medicine.
- TRI99. 2001. Toxic Chemical Release Inventory 1999. Data contained in the Toxic Chemical Release Inventory (TRI). National Library of Medicine. <http://www.epa.gov/triexplorer/>.
- USDOC Exports. 1986. U.S. Exports, Schedule E, Commodity by Country, 1985. Washington, D.C.: U.S. Government Printing Office.
- USDOC Exports. 1990. U.S. Exports, Harmonized Schedule B, Commodity by Country. Washington, D.C.: U.S. Government Printing Office.
- USDOC Imports. 1986. U.S. Imports for Consumption and General Imports, TSUSA Commodity by Country of Origin. Washington, D.C.: U.S. Government Printing Office.
- USDOC Imports. 1990. U.S. Imports for Consumption, Harmonized TSUSA Commodity by Country of Origin. Washington, D.C.: U.S. Government Printing Office.