

# Mechanisms of Chloroform and Carbon Tetrachloride Toxicity in Primary Cultured Mouse Hepatocytes

by Randall J. Ruch,\* James E. Klaunig,\*† Norman E. Schultz,\* Augusta B. Askari,‡ David A. Lacher,\* Michael A. Pereira,§ and Peter J. Goldblatt\*


Mechanisms of chloroform ( $\text{CHCl}_3$ ) and carbon tetrachloride ( $\text{CCl}_4$ ) toxicity to primary cultured male B6C3F1 mouse hepatocytes were investigated. The cytotoxicity of both  $\text{CHCl}_3$  and  $\text{CCl}_4$  was dose- and duration-dependent. Maximal hepatocyte toxicity, as determined by lactate dehydrogenase leakage into the culture medium, occurred with the highest concentrations of  $\text{CHCl}_3$  (5 mM) and  $\text{CCl}_4$  (2.5 mM) used and with the longest duration of treatment (20 hr).  $\text{CCl}_4$  was approximately 16 times more toxic than  $\text{CHCl}_3$  to the hepatocytes. The toxicity of these compounds was decreased by adding the mixed function oxidase system (MFOS) inhibitor, SKF-525A (25  $\mu\text{M}$ ) to the cultures. The addition of diethyl maleate (0.25 mM), which depletes intracellular glutathione (GSH)-potentiated  $\text{CHCl}_3$  and  $\text{CCl}_4$  toxicity. The toxicity of  $\text{CHCl}_3$  and  $\text{CCl}_4$  could also be decreased by adding the antioxidants *N,N'*-diphenyl-*p*-phenylenediamine (DPPD) (25  $\mu\text{M}$ ),  $\alpha$ -tocopherol acetate (Vitamin E) (0.1 mM), or superoxide dismutase (SOD) (100 U/mL) to the cultures. These results suggest that: in mouse hepatocytes, both  $\text{CHCl}_3$  and  $\text{CCl}_4$  are metabolized to toxic components by the MFOS; GSH plays a role in detoxifying those metabolites; free radicals are produced during the metabolism of  $\text{CHCl}_3$  and  $\text{CCl}_4$ ; and free radicals may be important mediators of the toxicity of these two halomethanes.

## Introduction


Chloroform ( $\text{CHCl}_3$ ) and carbon tetrachloride ( $\text{CCl}_4$ ) are formed as by-products of sewage and drinking water chlorination and have been identified in public drinking water (1). Both compounds are toxic to mammalian liver *in vivo* and to isolated liver cells, with  $\text{CCl}_4$  being more toxic than  $\text{CHCl}_3$  (2).  $\text{CCl}_4$  is hepatocarcinogenic in both rats and mice (3), whereas  $\text{CHCl}_3$  has been shown to be hepatocarcinogenic only in the B6C3F1 strain of mouse when administered by corn oil gavage (4). Both compounds may be functioning as liver tumor promoters (5).

The metabolism of  $\text{CCl}_4$  and  $\text{CHCl}_3$  may provide clues to their relative cytotoxic potency and their capability as tumor promoters. It is becoming evident that free radicals generated during xenobiotic metabolism play a role in cellular toxicity (6), abnormal growth control (7), and tumor promotion (8). The metabolism of  $\text{CCl}_4$  and

$\text{CHCl}_3$  has been extensively studied in both rats and mice. The first step in  $\text{CCl}_4$  metabolism is a one-electron reduction and homolytic cleavage catalyzed by cytochrome P450 of the mixed function oxidase system (MFOS) to yield the trichloromethyl radical (9) [Eq. (1)].



P450 is maintained in the reduced form by reduced nicotinamide adenine dinucleotide (NADPH). The  $\cdot\text{CCl}_3$  radical rapidly reacts with molecular oxygen to form the trichloromethylperoxyl radical (10) [Eq. (2)].



The  $\cdot\text{OOCCL}_3$  radical is more electrophilic than the  $\cdot\text{CCl}_3$  radical and may be more responsible for attacks on unsaturated fatty acids, leading to lipid peroxidation (10).  $\cdot\text{CCl}_3$  may be more involved in covalent binding reactions of  $\text{CCl}_4$  (10).

$\text{CHCl}_3$  is first metabolized by a cytochrome P450-catalyzed hydroxylation to trichloromethanol ( $\text{CCl}_3\text{OH}$ ), which spontaneously dehydrochlorinates to the toxic component, phosgene ( $\text{COCl}_2$ ) (11) [Eq. (3)].



Phosgene can covalently bind to macromolecules if not detoxified by conjugation with glutathione (GSH), or

\*Department of Pathology, Medical College of Ohio, Toledo, OH 43699.

§Health Effects Research Laboratory, U.S. Environmental Protection Agency, Cincinnati, OH 45268.

†Author to whom correspondence should be addressed.

‡Department of Surgery, Medical College of Ohio, Toledo, OH 43699.

can be further metabolized to carbon dioxide ( $\text{CO}_2$ ) by reaction with water (11). It has also been suggested that free radicals are generated during  $\text{CHCl}_3$  metabolism and that they play a role in  $\text{CHCl}_3$  toxicity (12, 13). Most studies, however, have failed to detect enhanced levels of free radicals or of lipid peroxidation, although no study has used the B6C3F1 strain of mouse.

To provide evidence that the hepatotoxicity of  $\text{CHCl}_3$  and  $\text{CCl}_4$  in B6C3F1 mice might be caused by free radical production during the metabolism of these compounds, the present study was undertaken. We examined whether the modifiers of hepatocyte xenobiotic metabolism, SKF-525A and diethyl maleate (DEM), and the antioxidants,  $\alpha$ -tocopherol acetate (vitamin E),  $N,N'$ -diphenyl- $p$ -phenylenediamine (DPPD), and superoxide dismutase (SOD), could alter the toxicity of  $\text{CHCl}_3$  and  $\text{CCl}_4$  to isolated B6C3F1 mouse hepatocytes. SKF-525A is an inhibitor of the MFOS, DEM decreases intracellular GSH levels, Vitamin E and DPPD are free-radical scavengers, and SOD dismutates superoxide radicals to hydrogen peroxide.

## Materials and Methods

### Animals

Six-month-old male B6C3F1 mice were used in this study. They were bred at the Medical College of Ohio from C3H/He sires and C57Bl/6 dams (Charles River Laboratories, Inc., Wilmington, MA). Mice were kept in plastic cages (five per cage) containing corncob bedding and given water and certified Purina Lab Chow *ad libitum*.

### Chemicals

Leibovitz's L-15 medium and gentamicin were purchased from Grand Island Biological Co. (Grand Island, NY); fetal bovine serum, from Hyclone Laboratories (Logan, UT);  $\text{CHCl}_3$  and  $\text{CCl}_4$ , from Aldrich Chemical Co. (Madison, WI); and DEM,  $\alpha$ -tocopherol acetate, DPPD, SOD, dimethyl sulfoxide (DMSO), dexamethasone and glucose, from Sigma Chemical Co. (St. Louis, MO). SKF-525A was a gift from Smith, Kline and French Laboratories (Philadelphia, PA). All other reagents were purchased from Sigma Chemical Co.

### Hepatocyte Isolation and Culture

Mouse hepatocytes were isolated by two-stage *in situ* perfusion (14). Cell viability of the isolated cells was 90%–95%, as determined by exclusion of trypan blue. Hepatocytes were cultured in L-15 medium supplemented with glucose (1 mg/mL), dexamethasone (1  $\mu\text{M}$ ), fetal bovine serum (10%), and gentamicin (50  $\mu\text{g}/\text{mL}$ ). The hepatocytes were plated at a density of  $0.5 \times 10^6$  cells per 60 mm dish and incubated in a humidified 100% air incubator at 37°C.




FIGURE 1. Toxicity of chloroform ( $\text{CHCl}_3$ ) to primary cultured male B6C3F1 mouse hepatocytes after 2-, 4-, or 20-hr treatment. Each point represents the mean ( $n=3$ ) of the percent total LDH released.

### Cytotoxicity Assay

After initial plating and a 4-hr attachment period, the cultured cells were washed once and re-fed with 5 mL of fresh medium. The cultures were then treated with the test compounds. All compounds, except SOD, were dissolved in DMSO. The final concentration of DMSO in all cultures, except for untreated cultures, was 0.4%. After 2, 4, and 20 hr of treatment, aliquots of the medium were removed, filtered through Nitrex nylon mesh to remove cells, and analyzed for lactate dehydrogenase activity (LDH) on a Beckman Multistat Analyzer (Beckman Instrument Corp., Palo Alto, CA). Total LDH per culture was also determined from 0.01% Triton X-100 lysates of untreated cultures, mean of three dishes.




FIGURE 2. Toxicity of carbon tetrachloride ( $\text{CCl}_4$ ) to primary cultured male B6C3F1 mouse hepatocytes after 2-, 4-, or 20-hr treatment. Each point represents the mean ( $n=3$ ) of the percent total LDH released.

**Table 1. Effects of the metabolic inhibitors, SKF-525A and diethyl maleate, on chloroform toxicity in primary cultured B6C3F1 mouse hepatocytes.**

Treatments		Percent total LDH release at time		
CHCl <sub>3</sub> mM	Inhibitor <sup>a</sup>	2 hr	4 hr	20 hr
0.0	None	5.6 ± 1.7 <sup>b</sup>	5.7 ± 1.2	14.3 ± 2.3
0.0	SKF-525A	6.4 ± 0.9	6.8 ± 0.5	24.2 ± 1.0*
0.0	DEM	7.5 ± 1.4	7.8 ± 1.4	17.3 ± 1.3
1.0	None	16.8 ± 2.1	16.3 ± 1.9	27.0 ± 2.7
1.0	SKF-525A	12.6 ± 2.0	15.7 ± 2.4	38.9 ± 1.9*
1.0	DEM	28.3 ± 0.7*	48.8 ± 2.9*	54.4 ± 2.7*
2.5	None	25.7 ± 2.4	26.9 ± 3.9	33.3 ± 3.3
2.5	SKF-525A	15.8 ± 5.2*	19.7 ± 3.0	52.2 ± 1.8*
2.5	DEM	47.0 ± 0.6*	58.7 ± 1.8*	65.1 ± 2.0*
5.0	None	42.7 ± 5.1	44.9 ± 4.7	58.3 ± 6.0
5.0	SKF-525A	25.0 ± 0.9*	33.3 ± 0.9*	67.5 ± 2.9
5.0	DEM	57.0 ± 1.3*	65.9 ± 2.8*	73.2 ± 2.4*

<sup>a</sup> Inhibitor concentrations were 25  $\mu$ M for SKF-525A and 0.25  $\mu$ M for diethyl maleate (DEM).<sup>b</sup> Mean  $\pm$  SD ( $n=3$ ).\* Significant difference versus the corresponding CHCl<sub>3</sub>-only treatment.**Table 2. Effects of metabolic inhibitors, SKF-525A and diethyl maleate, on carbon tetrachloride toxicity in primary cultured B6C3F1 mouse hepatocytes.**

Treatments		Percent total LDH release at time		
CCl <sub>4</sub> mM	Inhibitor <sup>a</sup>	2 hr	4 hr	20 hr
0.0	None	8.0 ± 1.0 <sup>b</sup>	8.9 ± 1.5	14.8 ± 1.0
0.0	SKF-525A	7.9 ± 0.6	8.6 ± 0.4	24.7 ± 0.6*
0.0	DEM	7.5 ± 0.4	12.8 ± 2.4	16.3 ± 1.3
1.0	None	53.0 ± 4.8	49.3 ± 3.5	73.6 ± 2.2
1.0	SKF-525A	36.1 ± 6.9*	40.8 ± 8.5	80.7 ± 5.2
1.0	DEM	72.8 ± 2.7*	73.9 ± 4.8*	86.6 ± 0.9*
2.5	None	65.5 ± 1.6	66.8 ± 2.8	76.4 ± 5.0
2.5	SKF-525A	52.7 ± 7.7*	55.9 ± 5.1*	85.4 ± 4.6
2.5	DEM	74.4 ± 3.2*	75.5 ± 4.2*	88.0 ± 2.3*

<sup>a</sup> Inhibitor concentrations were 25  $\mu$ M for SKF-525A and 0.25  $\mu$ M for diethyl maleate (DEM).<sup>b</sup> Mean  $\pm$  SD ( $n=3$ ).\* Significant difference versus the corresponding CCl<sub>4</sub>-only treatment.**Table 3. Effects of the antioxidants DPPD, vitamin E, and superoxide dismutase on chloroform toxicity in primary cultured B6C3F1 mouse hepatocytes.**

Treatments		Percent total LDH release at time		
CHCl <sub>3</sub> mM	Antioxidant <sup>a</sup>	2 hr	4 hr	20 hr
0.0	None	5.6 ± 1.7 <sup>b</sup>	5.7 ± 1.2	14.3 ± 2.3
0.0	DPPD	5.1 ± 0.6	5.6 ± 0.4	11.8 ± 0.3
0.0	Vitamin E	7.3 ± 1.3	7.9 ± 0.9	16.0 ± 0.7
0.0	SOD	4.4 ± 0.3	4.7 ± 0.5	12.3 ± 1.1
1.0	None	16.8 ± 2.1	16.3 ± 1.9	27.0 ± 2.7
1.0	DPPD	9.0 ± 0.4*	5.7 ± 1.6*	14.3 ± 0.5*
1.0	Vitamin E	14.4 ± 3.5	14.8 ± 2.6	25.9 ± 3.2
1.0	SOD	9.0 ± 2.3*	13.4 ± 2.4	19.3 ± 2.6*
2.5	None	25.7 ± 2.4	26.0 ± 3.9	33.3 ± 3.3
2.5	DPPD	14.5 ± 0.9*	10.4 ± 0.2*	24.3 ± 1.9*
2.5	Vitamin E	14.0 ± 0.6*	12.5 ± 0.9*	18.6 ± 0.7*
2.5	SOD	12.1 ± 1.4*	14.3 ± 1.1*	19.7 ± 1.5*
5.0	None	42.7 ± 5.1	44.9 ± 4.7	58.3 ± 6.0
5.0	DPPD	30.3 ± 1.7*	33.2 ± 0.6*	47.5 ± 0.8*
5.0	Vitamin E	19.7 ± 2.3*	20.0 ± 2.4*	27.7 ± 2.8*
5.0	SOD	30.0 ± 4.9*	33.6 ± 2.8*	41.0 ± 2.7*

<sup>a</sup> Antioxidant concentrations were 25  $\mu$ M for DPPD, 0.1 mM for vitamin E, and 100 U/mL for SOD.<sup>b</sup> Mean  $\pm$  SD ( $n=3$ ).\* Significant difference versus the corresponding CHCl<sub>3</sub>-only treatment.

**Table 4. Effects of the antioxidants DPPD, vitamin E, and superoxide dismutase on carbon tetrachloride toxicity in primary cultured B6C3F1 mouse hepatocytes.**

Treatments		Percent total LDH release at time		
CCl <sub>4</sub> , mM	Antioxidant <sup>a</sup>	2 hr	4 hr	20 hr
0.0	None	8.0 ± 1.0 <sup>b</sup>	8.9 ± 1.5	14.8 ± 1.0
0.0	DPPD	5.7 ± 1.5	6.2 ± 1.4	12.3 ± 1.3
0.0	Vitamin E	7.7 ± 0.6	6.9 ± 0.2	13.8 ± 0.8
0.0	SOD	7.5 ± 0.6	9.9 ± 0.8	14.8 ± 1.1
1.0	None	53.0 ± 4.8	49.3 ± 3.5	73.6 ± 2.2
1.0	DPPD	28.5 ± 3.6*	30.9 ± 5.7*	47.6 ± 4.4*
1.0	Vitamin E	40.4 ± 3.6*	34.8 ± 3.9*	60.1 ± 2.3*
1.0	SOD	36.9 ± 2.2*	37.7 ± 2.5*	58.3 ± 4.6*
2.5	None	65.5 ± 1.6	66.8 ± 2.8	76.4 ± 5.0
2.5	DPPD	44.2 ± 4.6*	50.0 ± 7.0*	61.0 ± 5.9*
2.5	Vitamin E	45.2 ± 5.2*	40.6 ± 5.7*	62.7 ± 3.9*
2.5	SOD	42.9 ± 3.0*	41.9 ± 2.1*	60.0 ± 1.4*

<sup>a</sup> Antioxidant concentrations were 25 μM for DPPD, 0.1 mM for vitamin E, and 100 U/mL for SOD.

<sup>b</sup> Mean ± SD (n=3).

\* Significant difference versus the corresponding CCl<sub>4</sub>-only treatment.

LDH in unused, fresh culture medium was also determined to indicate initial LDH levels in the cultures. Cytotoxicity in a culture was expressed as the "percent total LDH released" by using the formula (15–17):

$$\%LDH = \frac{LDH \text{ in medium of treated culture} - LDH \text{ in unused medium}}{\text{mean total LDH} - LDH \text{ in untreated medium}} \times 100\%$$

Values for the percent total LDH released were determined in three replicate cultures for each treatment and sampling time, and the means ± S.D. were determined. Different treatments were compared statistically using Student's *t*-test.

## Results

The toxicity of CHCl<sub>3</sub> to isolated B6C3F1 mouse hepatocytes was dependent on dose and treatment duration (Fig. 1). Little increase in LDH leakage was evident in cultures treated with less than 1.0 mM CHCl<sub>3</sub>. CCl<sub>4</sub> toxicity was also dependent on dose and treatment duration (Fig. 2). As with CHCl<sub>3</sub>, little difference in LDH release was evident between the 2-hr and 4-hr sampling times, and LDH release was greatest after 20 hr of treatment. CCl<sub>4</sub> was approximately 16 times more toxic than CHCl<sub>3</sub> to the hepatocytes. Percent total LDH activities of 50% occurred with approximately 0.25 mM CCl<sub>4</sub> and 4.0 mM CHCl<sub>3</sub> at the 20-hr sampling time (Figs. 1 and 2).

The addition of SKF-525A (25 μM) to CHCl<sub>3</sub>- or CCl<sub>4</sub>-treated cultures could be shown to significantly reduce LDH release below that of cultures treated only with CHCl<sub>3</sub> or CCl<sub>4</sub> (Tables 1 and 2). This effect was evident only at the 2-hr and 4-hr sampling times. At 20 hr, SKF-525A was cytotoxic by itself and either increased or had no effect on LDH release in CHCl<sub>3</sub>- and CCl<sub>4</sub>-treated cultures. Adding DEM (0.25 mM) to CHCl<sub>3</sub>- or CCl<sub>4</sub>-treated cultures significantly increased LDH release above CHCl<sub>3</sub>- or CCl<sub>4</sub>-only treated cultures (Tables 1

and 2). This effect was evident at 2, 4, and 20 hr. DEM was not toxic by itself.

The antioxidants, DPPD (25 μM), vitamin E (0.1 mM), and SOD (100 U/mL), when added to CHCl<sub>3</sub>- and CCl<sub>4</sub>-treated cultures, significantly reduced hepatocyte toxicity below that of cultures treated with CHCl<sub>3</sub> or CCl<sub>4</sub> alone (Tables 3 and 4). With the high dose of CHCl<sub>3</sub> (5 mM), vitamin E best prevented cytotoxicity compared to SOD and DPPD (Table 3). No differences in antioxidant protection were evident with 2.5 mM CHCl<sub>3</sub>. DPPD provided the greatest protection against low-dose CHCl<sub>3</sub> (1 mM) toxicity. Cytotoxicity of the low CCl<sub>4</sub> dose (1.0 mM) was best prevented by DPPD, whereas no differences between the antioxidants were apparent with the high CCl<sub>4</sub> dose (2.5 mM) (Table 4).

## Discussion

The results of this study indicate that CHCl<sub>3</sub> and CCl<sub>4</sub> are cytotoxic to primary cultured B6C3F1 mouse hepatocytes. Both compounds exhibited dose- and time-dependent effects. CCl<sub>4</sub> was approximately 16 times more toxic than CHCl<sub>3</sub>. Previous studies have also indicated that CCl<sub>4</sub> is more hepatotoxic than CHCl<sub>3</sub> (2, 3).

CHCl<sub>3</sub>- and CCl<sub>4</sub>-induced cytotoxicity was decreased by the simultaneous treatment of SKF-525A, an inhibitor of mammalian MFOS. These results substantiate previous studies that indicated both CHCl<sub>3</sub> and CCl<sub>4</sub> are metabolized in hepatocytes by the MFOS to hepatotoxic metabolites (9, 11). In the case of CCl<sub>4</sub>, the initial reaction is a reduction and homolytic cleavage of CCl<sub>4</sub> to the trichloromethyl radical (·CCl<sub>3</sub>). This radical may react directly with cellular macromolecules or may react with oxygen to form the trichloromethylperoxyl radical (·OOCCL<sub>3</sub>), which may then attack lipids more readily than ·CCl<sub>3</sub> (10). In any event, the initiation of free radicals and attack on membrane lipids ultimately can lead to lipid peroxidation chain reactions and cell death (10).

CHCl<sub>3</sub> is thought to be first hydroxylated by a P-450 reaction to trichloromethanol (CCl<sub>3</sub>OH), which dehy-

drochlorinates spontaneously to phosgene ( $\text{COCl}_2$ ) (11). Phosgene is thought to covalently bind cellular macromolecules and to be the ultimate cytotoxic component (11). However, two reports (12, 13) have suggested that  $\text{CHCl}_3$  is metabolized to free radicals and that it induces lipid peroxidation.

The results of the present study suggest that free radical production is an important mechanism of both  $\text{CCl}_4$  and  $\text{CHCl}_3$  toxicity in B6C3F1 mouse hepatocytes. The antioxidants SOD, DPPD, and vitamin E all significantly reduced the toxicity of  $\text{CCl}_4$  and  $\text{CHCl}_3$  to these cells. Similarly, depletion of cellular glutathione (GSH) (with DEM), increased  $\text{CCl}_4$  and  $\text{CHCl}_3$  toxicity to the hepatocytes. GSH functions both as an antioxidant and in the conjugation of xenobiotic metabolites (18).

Thus, the results of this initial study indicate that both  $\text{CCl}_4$  and  $\text{CHCl}_3$  are metabolized in B6C3F1 mouse hepatocytes by the MFOS, that cellular GSH is important in the detoxification of  $\text{CCl}_4$  and  $\text{CHCl}_3$  metabolites and/or induced free radicals, and that the cytotoxicity of both compounds might be partly mediated by free radical production. Previous studies have demonstrated the role of free radicals and lipid peroxidation in  $\text{CCl}_4$ -mediated hepatotoxicity. However, evidence that  $\text{CHCl}_3$ -induced hepatotoxicity is mediated by free radicals has been less substantiated. The results of the present study provide additional evidence that, in mouse hepatocytes, free radical production may be an important mechanism of  $\text{CHCl}_3$ -induced toxicity.

The authors express their gratitude to Mrs. Marilyn Cline and Mrs. Rose Kenney for secretarial assistance. This study was supported by the Environmental Protection Agency (EPA) cooperative agreement CR-810483. This material has been subjected to Agency review and has been accepted for publication as an EPA document. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

## REFERENCES

1. Deinzer, M., Schaumburg, F., and Klein, B. Environmental Health Sciences Center task force review on halogenated organics in drinking water. *Environ. Health Perspect.* 24: 209-239 (1978).
2. Ugazio, G. Halogenoalkanes and liver injury. In: *Biochemical Mechanisms of Liver Injury* (T. F. Slater, Ed.), Academic Press, New York, 1978, pp. 709-743.
3. International Agency for Research on Cancer (IARC). IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Man. Vol 20, IARC, Lyon, 1979, pp. 371-399.
4. IARC. IARC Monographs on the Evaluation of the Carcinogenic Risk of Chemicals to Man. IARC, col. 20, IARC, Lyon, 1979, pp. 401-427.
5. Klaunig, J. E., Ruch, R. J., and Pereira, M. A. Carcinogenicity of chlorinated methane and ethane compounds administered in drinking water to mice. *Environ. Health Perspect.* In press.
6. Younes, M., and Siegers, C. P. Interrelation between lipid peroxidation and other hepatotoxic events. *Biochem. Pharmacol.* 33: 2001-2003 (1984).
7. Oberley, L. W., Oberley, T. D., and Buettner, G. R. Cell division in normal and transformed cells: the possible role of superoxide and hydrogen peroxide. *Med. Hypoth.* 7: 21-42 (1981).
8. Troll, W., and Wiesner, R. The role of oxygen radicals as a possible mechanism of tumor promotion. *Ann. Rev. Pharmacol. Toxicol.* 25: 509-528 (1985).
9. Brattin, W. J., Glende, E. A., and Recknagel, R. O. Pathological mechanisms in carbon tetrachloride hepatotoxicity. *J. Free Radicals in Biol. Med.* 1: 27-38 (1985).
10. Slater, T. F. Free radicals as reactive intermediates in tissue injury. In: *Biological Reactive Intermediates, Chemical Mechanisms and Biological Effects*, Vol. 2 (R. Snyder, D. V. Parke, J. Kocsis, D. J. Jollow, and G. G. Gibson, Eds.), Plenum Publishing Co., New York, 1982, pp. 575-609.
11. Pohl, L. R., Bhooshan, B., Whittaker, N. F., and Krishna, G. Phosgene: a metabolite of chloroform. *Biochem. Biophys. Res. Commun.* 79: 684-257-258. (1969).
12. Van Dyke, R. A. On the fate of chloroform. *Anesthesiology* 30: 257-258 (1969).
13. Brown, B. R., Sipes, I. G., and Sagalyn, A. M. Mechanisms of acute hepatic toxicity: chloroform, halothane, and glutathione. *Anesthesiology* 41: 554-561 (1974).
14. Klaunig, J. E., Goldblatt, P. J., Hinton, D. E., Lipsky, M. M., Chacko, J., and Trump, B. F. Mouse liver cell culture. I. Hepatocyte isolation. *In Vitro* 17: 913-925 (1981).
15. Casini, A., Giorli, M., Hyland, R. J., Serroni, A., Gilfor, D., and Farber, J. L. Mechanisms of cell injury in the killing of hepatocytes by bromobenzene. *J. Biol. Chem.* 257: 6721-6728 (1982).
16. Starke, P. E., and Farber, J. L. Endogenous defenses against the cytotoxicity of hydrogen peroxide in cultured rat hepatocytes. *J. Biol. Chem.* 260: 86-92 (1985).
17. Hayes, M. A., Roberts, E., Roomi, M. W., Safe, S. H., Farber, E., and Cameron, R. E. Comparative influences of different PB-type and 3-MC-type polychlorinated biphenyl-induced phenotypes on cytotoxic hepatotoxicity of bromobenzene and acetaminophen. *Toxicol. Appl. Pharmacol.* 76: 118-127 (1984).
18. Ketterer, B., Coles, B., and Meyer, D. J. The role of glutathione in detoxification. *Environ. Health Perspect.* 49: 59-69 (1983).