

2005Table 2-5. Coal workers' pneumoconiosis: Number of deaths, death rates (per million population), and years of potential life lost (YPLL) by state, U.S. residents age 15 and over, 1993–2002

State	No. of Deaths		Crude Death Rate		Age-Adjusted Death Rate		YPLL to Life Expectancy			
	Deaths	Rank	Rate	Rank	Rate	Rank	Total	Rank	YPLL/death	Rank
Alabama	126	11	3.64	9	3.71	11	1,211	12	9.6	28
Alaska	2	42	0.44	35	1.45	18	15	44	7.3	48
Arizona	41	23	1.14	22	1.14	22	351	26	8.6	36
Arkansas	45	21	2.25	15	2.03	15	466	20	10.4	19
California	128	10	0.51	34	0.58	34	1,677	10	13.1	7
Colorado	94	12	3.02	13	3.91	10	856	14	9.1	33
Connecticut	14	33	0.54	33	0.47	35	127	33	9.1	33
Delaware	7	36	1.18	21	1.30	20	58	38	8.2	41
District of Columbia	1	46	0.23	44	0.23	43	8	48	8.3	38
Florida	178	8	1.49	18	1.10	24	1,788	9	10.0	23
Georgia	36	25	0.60	31	0.78	30	400	24	11.1	13
Hawaii	–	–	–	–	–	–	–	–	–	–
Idaho	7	36	0.74	28	0.79	29	90	35	12.9	8
Illinois	259	6	2.76	14	2.79	14	2,656	6	10.3	20
Indiana	156	9	3.36	11	3.45	13	1,514	11	9.7	25
Iowa	39	24	1.71	16	1.33	19	308	30	7.9	44
Kansas	23	31	1.12	23	1.08	25	362	25	15.7	4
Kentucky	897	4	28.69	3	29.85	3	10,370	4	11.6	11
Louisiana	20	32	0.59	32	0.63	32	413	22	20.6	3
Maine	1	46	0.10	48	0.10	48	8	48	8.3	38
Maryland	65	17	1.60	17	1.85	16	633	19	9.7	25
Massachusetts	1	46	0.02	50	0.02	50	14	46	14.3	6
Michigan	84	14	1.09	24	1.13	23	911	13	10.8	16
Minnesota	2	42	0.05	49	0.05	49	21	43	10.6	17
Mississippi	71	16	3.33	12	3.48	12	2,224	8	31.3	2
Missouri	29	27	0.68	29	0.62	33	278	31	9.6	28
Montana	9	35	1.28	20	1.22	21	67	37	7.4	47
Nebraska	3	41	0.23	44	0.20	45	24	42	7.9	44
Nevada	4	39	0.30	40	0.40	37	31	41	7.8	46
New Hampshire	4	39	0.43	36	0.46	36	32	40	8.0	42
New Jersey	44	22	0.68	29	0.66	31	420	21	9.6	28
New Mexico	48	20	3.63	10	4.18	9	408	23	8.5	37
New York	57	18	0.40	37	0.38	39	649	18	11.4	12
North Carolina	82	15	1.38	19	1.48	17	796	16	9.7	25
North Dakota	2	42	0.39	38	0.31	40	15	44	7.3	48
Ohio	484	5	5.46	7	5.35	8	4,390	5	9.1	33
Oklahoma	28	30	1.07	25	1.02	26	311	29	11.1	13
Oregon	7	36	0.27	41	0.26	41	74	36	10.6	17
Pennsylvania	5,097	1	52.86	2	43.64	2	47,090	1	9.2	31
Rhode Island	2	42	0.25	43	0.20	45	14	46	7.2	50
South Carolina	29	27	0.95	26	1.02	26	342	27	11.8	9
South Dakota	1	46	0.17	47	0.18	47	39	39	38.7	1
Tennessee	232	7	5.36	8	5.69	7	2,381	7	10.3	20
Texas	50	19	0.33	39	0.40	37	765	17	15.3	5
Utah	89	13	5.84	6	8.02	6	819	15	9.2	31
Vermont	1	46	0.21	46	0.23	43	8	48	8.3	38
Virginia	1,068	3	19.70	4	22.89	4	12,478	3	11.7	10
Washington	34	26	0.76	27	0.84	28	336	28	9.9	24
West Virginia	2,298	2	154.62	1	137.75	1	25,248	2	11.0	15
Wisconsin	11	34	0.27	41	0.25	42	112	34	10.1	22
Wyoming	29	27	7.67	5	8.98	5	232	32	8.0	42

– indicates no deaths listed.

NOTE: Slight variations in numbers may exist when compared to previous WoRLD Surveillance Reports. See appendices for source description, methods, and ICD codes.

SOURCE: National Center for Health Statistics multiple cause-of-death data. Population estimates from U.S. Bureau of the Census.