

**NORA *construction* sector: Proportionate mortality ratio (PMR) adjusted for age, sex, and race for selected work-related respiratory conditions, U.S. residents age 15 and over, selected states and years, 1990–1999**

Condition	Number of Deaths	PMR	95% Confidence Interval	
			LCL	UCL
Asbestosis	702	2.4	2.2	2.6
Malignant Mesothelioma (all sites)*	77	1.6	1.2	1.9
Silicosis	118	1.3	1.1	1.5
All Pneumoconioses	1,078	1.0	1.0	1.1
Byssinosis	4	-	-	-
Unspecified and Other Pneumoconioses	90	0.6	0.5	0.7
Hypersensitivity Pneumonitis	4	-	-	-
Coal Workers' Pneumoconiosis	188	0.4	0.3	0.4
All other conditions within this sector	351,414			
<b>TOTAL</b>	<b>353,675</b>			

NORA - National Occupational Research Agenda

- indicates no PMR was calculated because there were fewer than 5 deaths.

\* data for malignant mesothelioma are for 1999 only.

NOTE: The sum of the individual and all other condition death totals may be greater than the total number of deaths in the selected states and years because some decedents have more than one condition listed on their death certificates. See appendices for source description, methods, ICD codes, industry and occupation codes, industrial sector groupings, and list of selected states and years.

SOURCE: National Center for Health Statistics multiple cause-of-death data.