

NORA agriculture, forestry, and fishing sector: Proportionate mortality ratio (PMR) adjusted for age, sex, and race for selected work-related respiratory conditions, U.S. residents age 15 and over, selected states and years, 1990–1999

Condition	Number of Deaths	PMR	95% Confidence Interval	
			LCL	UCL
Hypersensitivity Pneumonitis	53	7.2	5.4	9.5
Malignant Mesothelioma (all sites)*	16	0.6	0.3	0.9
Byssinosis	2	-	-	-
Unspecified and Other Pneumoconioses	27	0.2	0.1	0.3
Silicosis	14	0.2	0.1	0.3
Asbestosis	43	0.2	0.1	0.3
All Pneumoconioses	136	0.2	0.1	0.2
Coal Workers' Pneumoconiosis	50	0.1	0.1	0.2
All other conditions within this sector	254,403			
TOTAL	254,744			

NORA - National Occupational Research Agenda

- indicates no PMR was calculated because there were fewer than 5 deaths.

* data for malignant mesothelioma are for 1999 only.

NOTE: The sum of the individual and all other condition death totals may be greater than the total number of deaths in the selected states and years because some decedents have more than one condition listed on their death certificates. See appendices for source description, methods, ICD codes, industry and occupation codes, industrial sector groupings, and list of selected states and years.

SOURCE: National Center for Health Statistics multiple cause-of-death data.