

Table 1 (continued). Cancer incidence trends by joinpoint analyses* for 1973 through 1999 for the most common cancers by race, sex, and age, SEER†, 1973-1999.

| Site/Race/Sex | Age | Joinpoint Analyses* (1973-1999) | | | | | | | |
|------------------|-----------------|---------------------------------|-------|----------------|--------|----------------|---------|----------------|-------|
| | | Trend 1 | | Trend 2 | | Trend 3 | | Trend 4 | |
| | | Range of Years | APC‡ | Range of Years | APC‡ | Range of Years | APC‡ | Range of Years | APC‡ |
| Prostate | | | | | | | | | |
| All Races | All ages | 1973-88 | 2.7 § | 1988-92 | 16.2 § | 1992-95 | -11.7 § | 1995-99 | 1.4 |
| | <50 | 1973-90 | 3.0 § | 1990-93 | 33.6 § | 1993-99 | 10.9 § | | |
| | 50-64 | 1973-85 | 2.9 § | 1985-89 | 7.3 § | 1989-92 | 26.4 § | 1992-99 | 2.3 § |
| | 65-74 | 1973-88 | 3.7 § | 1988-92 | 20.1 § | 1992-95 | -11.2 § | 1995-99 | 0.6 |
| | 75+ | 1973-88 | 2.0 § | 1988-92 | 11.3 § | 1992-95 | -18.7 § | 1995-99 | -0.1 |
| | | | | | | | | | |
| White | All ages | 1973-88 | 2.8 § | 1988-92 | 16.1 § | 1992-95 | -12.8 § | 1995-99 | 1.6 |
| | <50 | 1973-89 | 3.5 § | 1989-93 | 24.5 § | 1993-99 | 9.7 § | | |
| | 50-64 | 1973-88 | 3.7 § | 1988-92 | 23.5 § | 1992-99 | 2.0 § | | |
| | 65-74 | 1973-88 | 4.0 § | 1988-92 | 20.0 § | 1992-95 | -12.3 § | 1995-99 | 0.9 |
| | 75+ | 1973-88 | 2.1 § | 1988-92 | 11.0 § | 1992-95 | -19.6 § | 1995-99 | -0.1 |
| | | | | | | | | | |
| Black | All ages | 1973-89 | 2.1 § | 1989-92 | 20.6 § | 1992-96 | -5.7 § | 1996-99 | 0.1 |
| | <50 | † | † | | | | | | |
| | 50-64 | 1973-89 | 2.2 § | 1989-93 | 26.2 § | 1993-99 | 1.4 | | |
| | 65-74 | 1973-89 | 2.4 § | 1989-92 | 24.7 § | 1992-99 | -5.0 § | | |
| | 75+ | 1973-89 | 1.9 § | 1989-92 | 14.3 § | 1992-95 | -16.3 § | 1995-99 | -1.0 |

* Joinpoint analysis of trends allowed for up to three joinpoints based on rates per 100,000 that were age-adjusted to the 2000 U.S. standard population.

† SEER (Surveillance, Epidemiology, and End Results) registries covering 10% of the U.S. population: metropolitan areas of San Francisco, Detroit, Atlanta, and Seattle-Puget Sound; and States of Connecticut, Hawaii, Iowa, Utah, and New Mexico.

‡ APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population using the joinpoint regression program.

§ APC is statistically significantly different from zero (two-sided p<.05).

† Joinpoint results are not shown if less than 10 cases in any year.