



**Table 7 supplement. Incidence rates and 95% confidence intervals of the 15 top cancers for American Indian/Alaska Native males by IHS region, CHSDA counties in selected areas in the United States, 1999-2004**

Cancer site <sup>a</sup>	Southwest			
	Rank	Rate <sup>c</sup>	95%CI	Count
All sites <sup>b</sup>		261.5	249.0 - 274.5	1,882
Prostate	1	67.0	60.4 - 74.0	418
Lung and bronchus	4	22.1	18.4 - 26.4	135
Colon and rectum	2	26.7	22.8 - 30.9	198
Kidney and renal pelvis	3	25.1	21.4 - 29.1	188
Urinary bladder	11	5.5	3.7 - 7.7	36
Non-Hodgkin lymphoma	7	10.4	8.2 - 13.1	84
Stomach	5	16.0	12.9 - 19.5	106
Oral cavity and pharynx	12	4.8	3.3 - 6.7	40
Liver and IBD	6	12.6	10.1 - 15.6	95
Leukemia	10	6.7	5.0 - 8.7	76
Pancreas	8	7.6	5.6 - 10.0	52
Esophagus	13	4.4	2.9 - 6.4	30
Myeloma	9	7.4	5.4 - 9.9	48
Melanoma of the skin	17	3.6	2.3 - 5.4	26
Larynx	d	d	d	d
Brain and ONS	16	3.8	2.6 - 5.5	39
Testis	14	4.2	3.2 - 5.6	61
Gallbladder	15	4.1	2.6 - 6.1	26

SEER = Surveillance, Epidemiology, and End Results; NPCR = National Program of Cancer Registries; NAACCR = North American Association of Central Cancer Registries; IHS = Indian Health Service; CHSDA = IHS Contract Health Services Delivery Area; CI = confidence interval; IBD = intrahepatic bile duct; ONS = other nervous system. Source: SEER and NPCR areas reported by NAACCR as meeting high-quality data standards from 1999 through 2004.

<sup>a</sup> Cancers are sorted in descending order according to sex-specific rates for AI/AN. More than 15 cancers may appear to include the top 15 cancers in every IHS region.

<sup>b</sup> All sites excludes myelodysplastic syndromes and borderline tumors.

<sup>c</sup> Rates are per 100,000 persons and were age-adjusted to the 2000 U.S. standard population (19 age groups - Census p25-1130).

<sup>d</sup> Statistic could not be calculated when fewer than 16 cases were reported.

**Years of data and registries used (30 states)**

1999-2004: Alabama, Alaska, California, Colorado, Connecticut, Florida, Idaho, Indiana, Iowa, Louisiana, Maine, Massachusetts, Montana, Nebraska, Nevada, New Mexico, New York, North Dakota, Oklahoma, Oregon, Pennsylvania, Rhode Island, South Carolina, Texas, Utah, Washington, Wyoming;

1999-2003: Arizona, Wisconsin; 2000-2004: Michigan.

Percent regional coverage of AI/AN CHSDA data to all AI/AN in region: Alaska - 100%, East - 12.2%, N. Plains - 38.1%, Pacific Coast - 55.1%, S. Plains - 65.0%, Southwest - 86.1%