

Table VII-1
CORPUS AND UTERUS, NOS CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2001
WITH UP TO THREE JOINPOINTS BY RACE AND AGE

		<u>Joinpoint Segment 1</u>		<u>Joinpoint Segment 2</u>		<u>Joinpoint Segment 3</u>		<u>Joinpoint Segment 4</u>	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	All Ages	1975-1979	-6.0 ^c	1979-1988	-1.7 ^c	1988-1998	0.6 ^c	1998-2001	-1.5
All Races	Under 50	1975-1984	-3.6 ^c	1984-2001	0.5				
All Races	50 and Over	1975-1979	-6.2 ^c	1979-1988	-1.7 ^c	1988-1998	0.6 ^c	1998-2001	-1.9
White	All Ages	1975-1979	-6.2 ^c	1979-1988	-1.5 ^c	1988-2001	0.4 ^c		
White	Under 50	1975-1983	-3.9 ^c	1983-2001	0.1				
White	50 and Over	1975-1979	-6.2 ^c	1979-1987	-1.7 ^c	1987-2001	0.3		
Black	All Ages	1975-2001	0.0						
Black	Under 50	1975-1985	-5.0 ^c	1985-2001	2.0 ^c				
Black	50 and Over	1975-2001	0.0						
<u>U.S. Cancer Mortality^b</u>									
All Races	All Ages	1975-1991	-1.6 ^c	1991-2001	-0.2				
All Races	Under 50	1975-1988	-4.0 ^c	1988-2001	0.6				
All Races	50 and Over	1975-1991	-1.4 ^c	1991-2001	-0.3 ^c				
White	All Ages	1975-1991	-1.7 ^c	1991-2001	-0.2				
White	Under 50	1975-1987	-4.0 ^c	1987-2001	0.5				
White	50 and Over	1975-1991	-1.6 ^c	1991-2001	-0.3				
Black	All Ages	1975-2001	-0.7 ^c						
Black	Under 50	1975-1997	-4.3 ^c	1997-2001	8.7				
Black	50 and Over	1975-2001	-0.5 ^c						

Joinpoint Regression Program Version 2.7, September 2003, National Cancer Institute. (<http://srab.cancer.gov/joinpoint/>).

^a The APC is the Annual Percent Change based on rates age-adjusted to the 2000 U.S. standard population by 5-year age groups. Trends are from the SEER 9 areas.

^b Trends are from the NCHS public use data file for the total US.

^c The APC is significantly different from zero (p<.05).

Table VII-2
CORPUS AND UTERUS, NOS CANCER (Invasive)

SEER INCIDENCE^a AND U.S. DEATH^b RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE

	Incidence ^a			US Death ^b		
	All Races, Females	White Females	Black Females	All Races, Females	White Females	Black Females
<u>AGE-SPECIFIC RATES, 1997-2001</u>						
AGE AT DIAGNOSIS/DEATH:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	-	-	-	-	-	-
25-29	1.2	1.2	-	0.1	-	-
30-34	2.9	2.6	-	0.2	0.2	-
35-39	6.4	6.2	4.5	0.4	0.4	0.5
40-44	12.9	13.1	6.4	0.8	0.7	1.1
45-49	24.0	24.4	13.3	1.8	1.7	1.9
50-54	47.9	51.4	24.0	3.4	3.3	4.5
55-59	68.3	73.1	37.2	6.4	6.1	9.6
60-64	88.0	93.7	71.7	10.8	9.8	20.3
65-69	96.3	103.7	80.0	15.4	13.7	32.8
70-74	98.7	104.8	83.7	20.4	18.9	40.0
75-79	97.5	103.7	68.7	24.5	23.2	42.4
80-84	94.4	100.0	72.0	29.8	28.7	46.1
85+	70.0	71.3	66.3	35.0	34.3	47.4
<u>AGE-ADJUSTED RATES, 1997-2001</u>						
AGE AT DIAGNOSIS/DEATH:						
All ages	25.2	26.6	17.8	4.1	3.9	6.9
Under 65	15.3	16.1	9.4	1.4	1.3	2.1
65 and over	93.7	99.5	75.7	23.0	21.7	40.3
All ages(IARC world std) ^c	18.2	19.2	12.8	2.5	2.3	4.3

^a SEER 9 areas. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups, unless noted.

^b NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups, unless noted.

^c Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 25 cases for the time interval.

Table VII-3
CORPUS AND UTERUS, NOS CANCER (Invasive)

SEER INCIDENCE^a AND U.S. DEATH^b RATES, AGE-ADJUSTED RATES, BY RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES</u>									
YEAR OF DIAGNOSIS:									
1975	35.4	6.5	111.3	37.3	6.8	117.1	19.9	-	62.5
1976	34.4	6.2	108.1	36.2	6.3	114.7	17.6	-	55.1
1977	31.6	5.7	99.6	33.1	5.9	104.5	20.1	-	63.4
1978	30.0	6.1	92.9	31.7	6.3	98.2	18.6	-	57.5
1979	27.9	5.3	87.0	29.2	5.3	91.7	16.8	-	50.5
1980	27.3	5.0	85.8	28.5	5.0	90.2	16.2	-	52.1
1981	27.3	5.6	84.0	28.3	5.5	88.3	17.4	-	55.5
1982	26.8	5.0	83.8	28.2	5.2	88.4	17.2	-	55.8
1983	26.6	4.8	83.4	27.8	4.9	87.8	18.8	-	59.2
1984	25.7	4.6	80.9	27.1	4.9	85.5	17.4	-	58.4
1985	25.2	4.6	79.2	26.5	4.8	83.4	17.5	-	57.5
1986	24.4	4.8	75.6	25.7	5.0	79.7	16.7	-	53.8
1987	24.7	5.0	76.4	26.1	5.3	80.7	15.5	-	51.7
1988	23.6	4.5	73.4	24.6	4.4	77.3	16.9	3.0	53.3
1989	24.4	4.5	76.6	25.5	4.8	79.7	19.4	-	65.5
1990	25.1	4.6	78.7	26.6	4.8	83.5	16.8	-	55.0
1991	24.6	4.4	77.5	25.9	4.4	82.1	17.6	3.3	55.0
1992	24.8	4.8	77.2	26.2	4.8	82.0	17.0	2.8	54.4
1993	24.3	5.2	74.5	25.7	5.1	79.5	17.0	3.6	52.3
1994	25.1	5.0	77.8	26.2	5.0	81.9	17.8	2.7	57.6
1995	25.3	4.7	79.3	26.4	4.6	83.6	18.7	2.7	60.6
1996	24.9	4.5	78.2	25.9	4.3	82.3	19.3	3.2	61.6
1997	25.8	5.0	80.3	27.3	5.0	85.6	16.3	2.4	52.8
1998	25.7	4.8	80.3	27.0	4.8	85.2	19.2	3.0	61.7
1999	25.2	5.2	77.8	26.7	5.1	83.4	17.2	2.7	55.2
2000	24.6	5.1	75.6	25.9	5.2	80.1	17.7	3.0	56.2
2001	24.7	5.0	76.4	26.1	4.9	81.7	18.5	3.7	57.4
1997-2001	25.2	5.0	78.0	26.6	5.0	83.1	17.8	3.0	56.7
<u>U.S. DEATH RATES</u>									
YEAR OF DEATH:									
1975	5.3	0.5	17.7	5.1	0.5	17.1	7.8	0.9	25.8
1976	5.3	0.5	17.9	5.1	0.5	17.2	8.1	0.9	27.0
1977	5.2	0.5	17.5	5.0	0.5	16.7	8.4	0.8	28.0
1978	5.1	0.5	17.1	4.8	0.5	16.2	8.2	0.8	27.3
1979	5.0	0.4	17.0	4.7	0.4	16.2	7.9	0.7	26.7
1980	5.1	0.4	17.4	4.8	0.4	16.5	8.4	0.9	28.1
1981	4.9	0.4	16.6	4.7	0.4	15.9	7.3	0.6	25.1
1982	4.9	0.4	16.5	4.6	0.4	15.7	7.6	0.4	26.4
1983	4.8	0.4	16.4	4.6	0.3	15.8	7.3	0.6	24.8
1984	4.7	0.4	16.0	4.4	0.3	15.2	7.7	0.7	26.1
1985	4.6	0.4	15.7	4.4	0.3	15.0	7.4	0.5	25.3
1986	4.4	0.3	15.2	4.2	0.3	14.4	7.3	0.6	24.9
1987	4.4	0.3	15.1	4.2	0.3	14.3	7.3	0.5	25.1
1988	4.3	0.3	14.9	4.1	0.3	14.0	7.4	0.5	25.6
1989	4.3	0.3	14.5	4.0	0.3	13.6	7.3	0.5	25.0
1990	4.3	0.3	14.8	4.1	0.3	13.9	7.4	0.4	25.6
1991	4.2	0.3	14.2	3.9	0.3	13.3	7.3	0.5	25.1
1992	4.2	0.3	14.4	4.0	0.3	13.6	7.0	0.4	24.1
1993	4.2	0.3	14.3	3.9	0.3	13.3	7.3	0.4	25.2
1994	4.2	0.3	14.3	3.9	0.3	13.4	7.1	0.3	24.7
1995	4.1	0.3	14.1	4.0	0.3	13.5	6.5	0.3	22.5
1996	4.1	0.3	14.1	3.9	0.3	13.2	7.2	0.4	25.0
1997	4.0	0.3	13.7	3.8	0.3	12.9	6.7	0.3	23.6
1998	4.1	0.3	14.0	3.9	0.3	13.2	6.8	0.4	23.6
1999	4.1	0.3	13.9	3.9	0.3	13.1	6.8	0.4	23.7
2000	4.1	0.4	13.9	3.8	0.3	13.0	7.2	0.4	25.1
2001	4.2	0.4	14.1	3.9	0.3	13.3	7.0	0.5	24.0
1997-2001	4.1	0.3	13.9	3.9	0.3	13.1	6.9	0.4	24.0

^a SEER 9 areas. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups, unless noted.

^b NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 25 cases for the time interval.

Table VII-4
CORPUS AND UTERUS, NOS CANCER (Invasive)

SURVIVAL RATES, BY RACE, DIAGNOSIS YEAR, STAGE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>5-YR RELATIVE SURVIVAL RATES</u>									
YEAR OF DIAGNOSIS:									
1960-1963 ^a	-	-	-	73	-	-	31	-	-
1970-1973 ^a	-	-	-	81	-	-	44	-	-
1974-1976 ^b	88.0	92.2	87.4	88.9	92.9	88.4	61.5	84.4 ^d	56.5
1977-1979 ^b	85.1	92.2	84.1	86.4	93.1	85.5	57.9	81.1 ^d	52.5
1980-1982 ^b	81.6	91.2	80.4	82.8	92.4	81.7	55.2	85.5 ^d	50.3
1983-1985 ^b	82.9	92.6	81.6	84.5	93.4	83.5	54.3	81.4 ^d	50.3
1986-1988 ^b	82.9	92.5	81.4	84.4	92.6	83.3	57.5	86.7	52.2
1989-1991 ^b	83.8	88.5	83.0	85.7	90.9	84.9	57.8	72.2 ^d	55.2
1992-1994 ^b	84.3	90.2	83.2	86.3	91.7	85.5	56.1	81.2	50.0
1995-2000 ^b	84.4 ^c	89.6 ^c	83.5 ^c	86.1 ^c	90.7 ^c	85.4 ^c	63.1	81.1	59.2
1995-2000 ^b									
ALL STAGES	84.4	89.6	83.5	86.1	90.7	85.4	63.1	81.1	59.2
LOCALIZED	95.8	96.3	95.8	96.6	97.0	96.6	85.2	92.2	83.4
REGIONAL	67.0	82.1	64.2	69.6	84.3	67.0	43.5	64.2 ^d	39.6
DISTANT	25.6	38.0	23.6	27.5	38.5 ^d	25.7	16.8	-	15.3
UNSTAGED	56.3	82.9 ^d	52.2	56.6	89.0	52.4	52.6 ^d	-	47.9 ^d

STAGE DISTRIBUTION (%) 1995-2000^b

ALL STAGES									
Number of cases	17,250	2,539	14,711	14,909	1,963	12,946	1,099	181	918
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
LOCALIZED	72	74	72	73	73	73	55	69	52
REGIONAL	16	16	16	15	16	15	22	19	22
DISTANT	8	7	8	7	7	7	16	6	18
UNSTAGED	4	4	5	4	4	4	8	7	8

5-YR RELATIVE SURVIVAL RATES, 1995-2000^b

AGE AT DIAGNOSIS:									
<45	89.5	-	-	89.9	-	-	80.3	-	-
45-54	90.8	-	-	92.0	-	-	81.2	-	-
55-64	88.5	-	-	90.9	-	-	61.2	-	-
65-74	81.9	-	-	84.2	-	-	55.4	-	-
75+	74.0	-	-	75.9	-	-	45.5 ^d	-	-
Under 65	89.5	-	-	91.2	-	-	71.1	-	-
65 and over	78.8	-	-	80.8	-	-	53.3	-	-

- ^a Rates are based on End Results data from a series of hospital registries and one population-based registry.
- ^b Rates are from the SEER 9 areas. They are based on data from population-based registries in Connecticut, New Mexico, Utah, Iowa, Hawaii, Atlanta, Detroit, Seattle-Puget Sound and San Francisco-Oakland. Rates are based on follow-up of patients into 2001.
- ^c The difference in rates between 1974-1976 and 1995-2000 is statistically significant (p<.05).
- ^d The standard error of the survival rate is between 5 and 10 percentage points.
- ^e The standard error of the survival rate is greater than 10 percentage points.
- Statistic could not be calculated.

Table VII-5
CORPUS AND UTERUS, NOS CANCER (Invasive)

SURVIVAL RATES^a

By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975- 1979	1980- 1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	
Relative Survival Rates (SEER ^b)																			
1-year	93.9	91.7	93.2	93.1	92.1	92.3	92.3	92.7	92.0	93.1	92.5	92.0	93.3	92.0	93.3	93.7	94.1	93.9	
2-year	90.3	86.7	88.2	87.1	87.5	88.4	88.0	88.2	88.1	89.8	88.7	87.4	88.8	88.5	89.4	89.8	89.6		
3-year	88.1	84.1	85.5	85.4	84.4	85.5	85.9	85.4	85.8	87.5	86.0	85.2	86.5	86.4	86.4	86.9			
4-year	86.7	82.9	83.8	84.0	83.6	84.0	84.9	84.1	84.3	86.2	84.8	83.7	85.3	85.5	85.6				
5-year	86.4	82.0	83.5	83.0	82.6	83.2	84.2	83.4	83.8	86.1	84.0	83.0	84.9	84.3					
6-year	86.0	81.7	82.9	82.6	81.8	82.9	83.8	82.3	82.7	85.4	83.2	82.0	84.3						
7-year	85.8	81.3	82.3	82.6	81.8	82.6	83.8	82.3	82.6	85.4	83.2	81.5							
8-year	85.8	80.9	82.2	82.6	81.6	82.6	82.8	82.1	82.2	85.1	82.0								
9-year	85.8	80.8	81.7	82.3	81.6	82.3	82.8	82.1	82.2	85.1									
10-year	85.8	80.7	81.7	82.3	81.6	81.9	82.8	81.5	81.8										
11-year	85.8	80.6	81.7	82.3	81.6	81.9	82.6	81.5											
12-year	85.8	80.6	81.7	82.3	81.6	81.7	81.8												
13-year	85.8	80.4	81.4	82.3	81.6	81.3													
14-year	85.8	80.3	81.0	82.3	81.5														
15-year	85.8	80.2	81.0	82.3															
16-year	85.8	79.8	80.8																
17-year	85.8	79.7																	
18-year	85.8																		
19-year	85.8																		
20-year	85.8																		

^a Survival rates are relative rates expressed as percents.
^b Rates are from the SEER 9 areas.

Table VII-6
CORPUS AND UTERUS, NOS CANCER (Invasive)

Percent Diagnosed With Cancer In 10, 20 and 30 Years and In Remaining Lifetime,
Given Cancer Free At Current Age By Race

Lifetime Risk (Percent) of Being Diagnosed (12 SEER Areas) With Cancer and
Lifetime Risk (Percent) of Dying (Total U.S.) From Cancer

1999-2001

All Races

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	2.62
10	0.00	0.01	0.06	2.65
20	0.01	0.06	0.25	2.66
30	0.05	0.24	0.76	2.66
40	0.19	0.72	1.50	2.63
50	0.54	1.34	2.03	2.49
60	0.85	1.57	1.99	2.06

Lifetime Risk of Being Diagnosed = 2.62%
Lifetime Risk of Dying = 0.50%

Whites

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	2.82
10	0.00	0.01	0.06	2.85
20	0.01	0.06	0.25	2.86
30	0.05	0.24	0.81	2.86
40	0.19	0.76	1.62	2.83
50	0.58	1.45	2.20	2.68
60	0.92	1.70	2.14	2.21

Lifetime Risk of Being Diagnosed = 2.82%
Lifetime Risk of Dying = 0.48%

Blacks

Current Age	+10 yrs	+20 yrs	+30 yrs	Eventually
0	0.00	0.00	0.01	1.74
10	0.00	0.01	0.05	1.78
20	0.01	0.05	0.15	1.78
30	0.04	0.14	0.44	1.79
40	0.11	0.41	1.03	1.78
50	0.31	0.96	1.46	1.75
60	0.71	1.26	1.52	1.57

Lifetime Risk of Being Diagnosed = 1.74%
Lifetime Risk of Dying = 0.73%

Table VII-7
CORPUS AND UTERUS, NOS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE AND U.S. DEATH RATES

By Race/Ethnicity and Sex

Incidence^a

RACE/ETHNICITY	Rate 1997-2001	Trend 1992-2001
	Rate per 100,000 persons	APC (%)
	Females	Females
All Races	24.5	-0.1
White	26.1	-0.1
White Hispanic ^b	17.3	0.3
White Non-Hispanic ^b	27.1	-0.1
Black	18.2	0.9
Asian/Pacific Islander	17.2	1.4 ^d
Amer Ind/Alaska Nat	10.2	-
Hispanic ^b	16.8	0.5

Mortality^c

RACE/ETHNICITY	Rate 1997-2001	Trend 1992-2001
	Rate per 100,000 persons	APC (%)
	Females	Females
All Races	4.1	-0.2
White	3.9	-0.2
White Hispanic ^b	3.3	0.5
White Non-Hispanic ^b	3.8	-0.1
Black	6.9	-0.1
Asian/Pacific Islander	2.1	0.4
Amer Ind/Alaska Nat	2.3	-1.8
Hispanic ^b	3.2	0.4

The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 25 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Incidence data are from the 12 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, and Alaska Native Registry).

^b Hispanic and Non-Hispanic are not mutually exclusive from Whites, Blacks, Asian Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics do not include cases from Detroit, Hawaii and Alaska Native Registry. For mortality, information is included for all states except Connecticut, Maine, Maryland, Oklahoma, New York, New Hampshire, North Dakota and Vermont.

^c Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

^d The APC is significantly different from zero (p<.05).

Table VII-8
CORPUS AND UTERUS, NOS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE RATES^a AND SEER DEATH RATES^b
By Registry, Race and Age

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>SEER INCIDENCE RATES^a, 1997-2001</u>									
REGISTRY									
12 registries	24.5	5.0	75.6	26.1	5.0	81.4	18.2	3.0	58.0
11 registries	24.5	5.0	75.7	26.1	5.0	81.4	18.2	3.0	58.0
9 registries	25.2	5.0	78.0	26.6	5.0	83.1	17.8	3.0	56.7
San Francisco-Oakland	24.9	4.4	78.6	27.5	4.3	88.1	19.1	3.4	60.2
Connecticut	28.3	5.1	89.1	29.2	5.2	92.0	17.5	-	54.1
Metropolitan Detroit	26.2	5.2	81.1	28.5	5.7	88.0	19.0	3.5	59.7
Hawaii	24.5	9.5	63.6	23.4	4.2	73.6	-	-	-
Iowa	27.8	5.6	85.8	28.0	5.7	86.4	-	-	-
New Mexico	19.0	4.8	56.1	19.3	4.4	58.1	-	-	-
Seattle-Puget Sound	26.2	4.7	82.6	27.4	4.8	86.6	17.2	-	59.0
Utah	23.4	5.0	71.3	23.5	5.0	72.1	-	-	-
Metropolitan Atlanta	20.4	3.6	64.5	22.6	4.5	70.0	16.3	2.3	52.8
San Jose-Monterey	22.9	3.8	72.8	25.2	3.8	81.3	-	-	-
Los Angeles	22.8	5.1	69.0	24.6	5.4	75.1	19.5	3.2	62.1
<u>SEER DEATH RATES^b, 1997-2001</u>									
REGISTRY									
12 registries	4.0	0.4	13.6	4.0	0.3	13.6	6.2	0.5	21.0
11 registries	4.0	0.4	13.6	4.0	0.3	13.6	6.2	0.5	21.1
9 registries	4.0	0.4	13.6	3.9	0.3	13.3	6.1	-	20.9
San Francisco-Oakland	4.1	0.3	13.8	4.3	-	14.5	5.9	-	20.7
Connecticut	3.9	-	13.2	3.8	-	13.0	5.8	-	18.7
Metropolitan Detroit	4.4	0.5	14.5	3.8	0.5	12.5	6.8	-	23.3
Hawaii	3.5	-	10.4	-	-	-	-	-	-
Iowa	4.1	-	13.9	4.1	-	13.9	-	-	-
New Mexico	3.6	-	12.0	3.6	-	12.2	-	-	-
Seattle-Puget Sound	4.0	-	13.8	4.1	-	14.0	-	-	-
Utah	4.1	-	14.2	4.1	-	14.3	-	-	-
Metropolitan Atlanta	3.9	-	13.5	3.4	-	11.8	5.4	-	18.8
San Jose-Monterey	3.9	-	13.6	4.1	-	14.3	-	-	-
Los Angeles	4.1	0.4	13.8	4.2	0.3	14.3	6.5	-	21.7

Note: San Jose-Monterey and Los Angeles are new SEER areas as of 1992 and their data are only used in tables where indicated.

^a SEER 9 areas. Incidence rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

^b NCHS public use data file for the total US. Death rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.

- Statistic could not be calculated or populations are inadequate.

Table VII-9
CORPUS AND UTERUS, NOS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 1997-2001

All Ages

State	Rate	SE	Rank	PD
TOTAL U.S.	4.1	0.02		
High Five States				
District of Columbia	7.1 ^{bc}	0.66	(01)	73.2
New Jersey	5.1 ^{bc}	0.14	(02)	24.4
Vermont	4.9 ^b	0.53	(03)	19.5
New York	4.8 ^{bc}	0.09	(04)	17.1
Pennsylvania	4.7 ^c	0.11	(05)	14.6
Low Five States				
Alaska	3.2 ^b	0.67	(47)	-22.0
Colorado	3.2 ^{bc}	0.18	(48)	-22.0
Nevada	3.0 ^{bc}	0.26	(49)	-26.8
North Dakota	3.0 ^b	0.38	(50)	-26.8
Arizona	2.9 ^{bc}	0.15	(51)	-29.3

Ages <50

State	Rate	SE	Rank	PD
TOTAL U.S.	0.3	0.01		
High Five States				
Hawaii	0.8 ^b	0.19	(01)	166.7
West Virginia	0.6 ^b	0.14	(02)	100.0
Vermont	0.6 ^b	0.23	(03)	100.0
Wyoming	0.5 ^b	0.24	(04)	66.7
South Dakota	0.5 ^b	0.19	(05)	66.7
Low Five States				
Rhode Island	0.2 ^b	0.10	(47)	-33.3
Oregon	0.2 ^b	0.06	(48)	-33.3
Alaska	0.2 ^b	0.11	(49)	-33.3
Delaware	0.1 ^b	0.10	(50)	-66.7
Idaho	0.1 ^b	0.08	(51)	-66.7

Ages 50+

State	Rate	SE	Rank	PD
TOTAL U.S.	13.9	0.08		
High Five States				
District of Columbia	24.6 ^{bc}	2.32	(01)	77.0
New Jersey	17.6 ^{bc}	0.50	(02)	26.6
New York	16.3 ^{bc}	0.32	(03)	17.3
Pennsylvania	16.1 ^{bc}	0.37	(04)	15.8
Vermont	16.1 ^b	1.80	(05)	15.8
Low Five States				
Colorado	10.9 ^{bc}	0.64	(47)	-21.6
Hawaii	10.4 ^b	1.07	(48)	-25.2
Arizona	9.9 ^{bc}	0.51	(49)	-28.8
Nevada	9.9 ^{bc}	0.90	(50)	-28.8
North Dakota	9.8 ^b	1.26	(51)	-29.5

For additional detailed cancer statistics by state, please visit the State Cancer Profiles website. <http://statecancerprofiles.cancer.gov/>

^a NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 U.S. standard population by 5-year age groups.
^b Absolute percent difference between state rate and total U.S. rate is 15% or more.
^c Difference between state rate and total U.S. rate is statistically significant (p<=.0002).
SE Standard error of the rate.
PD Percent difference between state rate and total U.S. rate.

Table VII-10
CORPUS AND UTERUS, NOS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2001
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <11 ^e	0 to <26 ^e	>=26 ^g	Complete ^h
<u>Race</u>									
All Races ^b	140,432	105,849	82,158	67,121	64,672	264,462	474,773	96,080	570,853
White ^b	127,016	97,961	77,056	63,723	61,587	242,104	441,307	86,284	527,591
Black ^b	7,805	4,561	3,045	1,892	1,630	12,930	19,237	1,871	21,108
Asian/Pacific Islander ^c	3,225	1,968	+	+	+	5,467	+	+	+
Hispanic ^d	5,965	3,852	+	+	+	10,546	+	+	+

Estimated prevalence percent^a on January 1, 2001, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	All Ages	0-9	10-19	20-29	Age Specific (Crude)						Age-Adjusted ^f
					30-39	40-49	50-59	60-69	70-79	80+	All Ages
<u>Race</u>											
All Races ^c	0.1492%	-	-	0.0025%	0.0180%	0.0802%	0.2827%	0.5864%	0.6946%	0.5967%	0.1508%
White ^c	0.1717%	-	-	0.0022%	0.0175%	0.0806%	0.3069%	0.6594%	0.7710%	0.6471%	0.1649%
Black ^c	0.0611%	-	-	0.0029%	0.0149%	0.0440%	0.1267%	0.3214%	0.3958%	0.2936%	0.0806%
Asian/Pacific Islander ^c	0.0937%	-	-	0.0031%	0.0209%	0.1054%	0.2505%	0.3098%	0.3185%	0.2788%	0.0985%
Hispanic ^d	0.0546%	-	-	0.0034%	0.0225%	0.0650%	0.1991%	0.3601%	0.3846%	0.3114%	0.0957%

^a US 2001 cancer prevalence counts are based on 2001 cancer prevalence proportions from the SEER registries and 1/1/2001 US population estimates based on the average of 2000 and 2001 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on b) SEER 9 Areas c) SEER 11 Areas d) SEER 11 Areas excluding Hawaii and Detroit.

^e Maximum limited-duration prevalence: 26 years for 1975-2001 SEER 9 data; 11 years for 1990-2001 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^f Percentages are age-adjusted to the 2000 US standard population by 5-year age groups.

^{g h} (g) Cases diagnosed 26 years or more ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h) Complete prevalence is obtained by summing 0 to <26 and >=26.

- Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

+ Not available.

SEER Incidence and US Death Rates+ Corpus Uteri Cancer, by Race




Figure VII-1

+ Source: SEER 9 areas and NCHS public use data file for the total US. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines are calculated using the Joinpoint Regression Program Version 2.7, September 2003, National Cancer Institute.

SEER Incidence and US Death Rates+ Corpus Uteri Cancer, by Age and Race




Figure VII-2

+ Source: SEER 9 areas and NCHS public use data file for the total US. Rates are age-adjusted to the 2000 US standard million population by 5-year age groups. Regression lines are calculated using the Joinpoint Regression Program Version 2.7, September 2003, National Cancer Institute.