

Table VII-1
CORPUS AND UTERUS, NOS CANCER (Invasive)

TRENDS IN SEER INCIDENCE^a AND U.S. MORTALITY^b USING THE JOINPOINT REGRESSION PROGRAM, 1975-2004
WITH UP TO THREE JOINPOINTS BY RACE AND AGE

		Joinpoint Segment 1		Joinpoint Segment 2		Joinpoint Segment 3		Joinpoint Segment 4	
		Year	APC	Year	APC	Year	APC	Year	APC
<u>SEER Cancer Incidence^a</u>									
All Races	All Ages	1975-1979	-6.0*	1979-1988	-1.7*	1988-1998	0.6*	1998-2004	-1.0*
All Races	Under 50	1975-1985	-3.3*	1985-2004	0.7*				
All Races	50 and Over	1975-1979	-6.2*	1979-1988	-1.7*	1988-1998	0.6*	1998-2004	-1.3*
White	All Ages	1975-1979	-6.1*	1979-1988	-1.6*	1988-1998	0.6*	1998-2004	-1.2*
White	Under 50	1975-1983	-4.0*	1983-2004	0.2				
White	50 and Over	1975-1979	-6.2*	1979-1988	-1.7*	1988-1998	0.7*	1998-2004	-1.5*
Black	All Ages	1975-2004	0.4*						
Black	Under 50	1975-1985	-5.4*	1985-2004	2.7*				
Black	50 and Over	1975-2004	0.3						
<u>SEER Delay-Adjusted Incidence^a</u>									
All Races	All Ages	1975-1979	-6.0*	1979-1988	-1.7*	1988-1998	0.6*	1998-2004	-0.8*
All Races	Under 50	1975-1985	-3.3*	1985-2004	0.7*				
All Races	50 and Over	1975-1979	-6.2*	1979-1988	-1.7*	1988-1998	0.6*	1998-2004	-1.2*
White	All Ages	1975-1979	-6.1*	1979-1988	-1.6*	1988-1998	0.7*	1998-2004	-1.0*
White	Under 50	1975-1983	-4.0*	1983-2004	0.2				
White	50 and Over	1975-1979	-6.2*	1979-1988	-1.7*	1988-1998	0.7*	1998-2004	-1.3*
Black	All Ages	1975-2004	0.5*						
Black	Under 50	1975-1985	-5.5*	1985-2004	2.9*				
Black	50 and Over	1975-2004	0.4*						
<u>U.S. Cancer Mortality^b</u>									
All Races	All Ages	1975-1992	-1.5*	1992-2004	0.0				
All Races	Under 50	1975-1988	-4.1*	1988-2004	0.7*				
All Races	50 and Over	1975-1992	-1.4*	1992-2004	-0.1				
White	All Ages	1975-1991	-1.7*	1991-2004	-0.2				
White	Under 50	1975-1988	-3.8*	1988-2004	0.8*				
White	50 and Over	1975-1991	-1.6*	1991-2004	-0.3*				
Black	All Ages	1975-1995	-0.8*	1995-2004	0.3				
Black	Under 50	1975-1995	-4.5*	1995-2004	3.1				
Black	50 and Over	1975-2004	-0.4*						

Joinpoint Regression Program Version 3.0, April 2005, National Cancer Institute. (<http://srab.cancer.gov/joinpoint/>).

The APC is the Annual Percent Change based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

Trends are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Trends are from the NCHS public use data file for the total US.

The APC is significantly different from zero (p<.05).

a

b

*

Table VII-2
CORPUS AND UTERUS, NOS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE^a RATES BY YEAR, RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
SEER INCIDENCE RATES									
YEAR OF DIAGNOSIS:									
1975	35.5	6.5	111.4	37.3	6.8	117.2	19.9	3.6	62.5
1976	34.4	6.2	108.1	36.2	6.3	114.7	17.6	3.3	55.1
1977	31.7	5.7	99.5	33.1	5.9	104.4	20.1	3.5	63.4
1978	30.0	6.1	92.8	31.7	6.3	98.0	18.6	3.7	57.5
1979	27.9	5.3	87.0	29.2	5.3	91.8	16.8	3.9	50.5
1980	27.3	5.0	85.8	28.5	5.0	90.2	16.2	-	52.1
1981	27.3	5.6	84.0	28.3	5.5	88.3	17.4	2.9	55.5
1982	26.8	5.0	83.8	28.2	5.2	88.5	17.1	2.5	55.2
1983	26.5	4.9	83.3	27.8	4.9	87.7	18.8	3.4	59.2
1984	25.7	4.6	80.9	27.1	4.8	85.5	17.4	-	58.4
1985	25.2	4.6	79.2	26.5	4.8	83.5	17.5	2.2	57.5
1986	24.4	4.8	75.6	25.6	5.0	79.6	16.8	2.5	54.1
1987	24.7	5.0	76.4	26.1	5.3	80.8	15.8	-	51.8
1988	23.6	4.5	73.4	24.6	4.5	77.3	16.9	3.0	53.3
1989	24.4	4.5	76.6	25.5	4.8	79.7	19.4	1.8	65.4
1990	25.1	4.6	78.6	26.5	4.8	83.4	16.9	2.3	55.3
1991	24.6	4.4	77.5	26.0	4.5	82.3	17.4	3.1	54.7
1992	24.8	4.8	77.1	26.2	4.8	82.0	16.9	2.8	54.1
1993	24.3	5.2	74.3	25.7	5.1	79.5	17.1	3.6	52.6
1994	25.1	5.0	77.8	26.3	5.0	81.9	17.7	2.5	57.6
1995	25.3	4.7	79.3	26.5	4.6	83.6	18.7	2.7	60.6
1996	24.9	4.5	78.3	26.0	4.4	82.5	19.3	3.2	61.5
1997	25.8	5.0	80.4	27.3	5.0	85.6	16.5	2.5	53.1
1998	25.7	4.8	80.3	27.0	4.7	85.3	19.4	3.0	62.3
1999	25.3	5.2	78.0	26.7	5.1	83.5	17.5	2.8	55.9
2000	24.7	5.2	75.9	26.2	5.3	80.7	17.9	3.0	57.1
2001	25.2	5.1	77.8	26.7	5.0	83.5	18.9	3.9	58.1
2002	24.7	5.0	76.3	25.1	4.5	78.9	23.1	4.7	71.2
2003	24.0	5.2	73.3	25.2	5.2	77.5	19.6	3.6	61.6
2004	24.4	5.3	74.4	25.4	5.1	78.7	19.1	3.3	60.5
1975-2004	26.2	5.0	81.8	27.6	5.1	86.7	18.1	3.0	57.7

^a SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta). Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VII-3
CORPUS AND UTERUS, NOS CANCER (Invasive)

AGE-ADJUSTED U.S. DEATH^a RATES BY YEAR, RACE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
U.S. DEATH RATES									
YEAR OF DEATH:									
1975	5.3	0.5	17.7	5.1	0.5	17.1	7.8	0.9	25.8
1976	5.3	0.5	17.9	5.1	0.5	17.2	8.1	0.9	27.0
1977	5.2	0.5	17.5	5.0	0.5	16.7	8.4	0.8	28.0
1978	5.1	0.5	17.1	4.8	0.5	16.2	8.2	0.8	27.3
1979	5.0	0.4	17.0	4.7	0.4	16.2	7.9	0.7	26.7
1980	5.1	0.4	17.4	4.8	0.4	16.5	8.4	0.9	28.1
1981	4.9	0.4	16.6	4.7	0.4	15.9	7.3	0.6	25.1
1982	4.9	0.4	16.5	4.6	0.4	15.7	7.6	0.4	26.4
1983	4.8	0.4	16.4	4.6	0.3	15.8	7.3	0.6	24.8
1984	4.7	0.4	16.0	4.4	0.3	15.2	7.7	0.7	26.1
1985	4.6	0.4	15.7	4.4	0.3	15.0	7.4	0.5	25.3
1986	4.4	0.3	15.2	4.2	0.3	14.4	7.3	0.6	24.9
1987	4.4	0.3	15.1	4.2	0.3	14.3	7.3	0.5	25.1
1988	4.3	0.3	14.9	4.1	0.3	14.0	7.4	0.5	25.6
1989	4.3	0.3	14.5	4.0	0.3	13.6	7.3	0.5	25.0
1990	4.3	0.3	14.8	4.1	0.3	13.9	7.4	0.4	25.6
1991	4.2	0.3	14.2	3.9	0.3	13.3	7.3	0.5	25.1
1992	4.2	0.3	14.4	4.0	0.3	13.6	7.0	0.4	24.1
1993	4.2	0.3	14.3	3.9	0.3	13.3	7.3	0.4	25.2
1994	4.2	0.3	14.3	3.9	0.3	13.4	7.1	0.3	24.7
1995	4.1	0.3	14.1	4.0	0.3	13.5	6.5	0.3	22.5
1996	4.1	0.3	14.1	3.9	0.3	13.2	7.2	0.4	25.0
1997	4.0	0.3	13.7	3.8	0.3	12.9	6.7	0.3	23.6
1998	4.1	0.3	14.0	3.9	0.3	13.2	6.8	0.4	23.6
1999	4.1	0.3	13.9	3.9	0.3	13.1	6.8	0.4	23.7
2000	4.1	0.4	13.9	3.8	0.3	13.0	7.2	0.4	25.1
2001	4.2	0.4	14.2	3.9	0.3	13.4	7.0	0.5	24.0
2002	4.2	0.3	14.1	3.9	0.3	13.1	7.3	0.4	25.4
2003	4.1	0.3	14.0	3.9	0.3	13.2	7.0	0.5	24.2
2004	4.1	0.4	13.9	3.9	0.4	13.0	7.0	0.5	24.0
1975-2004	4.4	0.4	15.1	4.2	0.3	14.3	7.3	0.5	25.1

^a NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).
- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VII-4
CORPUS AND UTERUS, NOS CANCER (Invasive)

SEER INCIDENCE^a AND U.S. DEATH^b RATES, AGE-ADJUSTED AND AGE-SPECIFIC RATES, BY RACE

	Incidence ^a			US Death ^b		
	All Races, Females	White Females	Black Females	All Races, Females	White Females	Black Females
<u>AGE-ADJUSTED RATES, 2000-2004</u>						
AGE AT DIAGNOSIS/DEATH:						
All ages	23.2	24.3	19.6	4.1	3.9	7.1
Under 65	14.5	15.1	10.4	1.4	1.3	2.2
65 and over	83.5	87.4	82.9	22.8	21.4	40.8
All ages(IARC world std) ^c	17.0	17.7	14.0	2.6	2.4	4.5
<u>AGE-SPECIFIC RATES, 2000-2004</u>						
AGE AT DIAGNOSIS/DEATH:						
<1	-	-	-	-	-	-
1-4	-	-	-	-	-	-
5-9	-	-	-	-	-	-
10-14	-	-	-	-	-	-
15-19	-	-	-	-	-	-
20-24	0.3	0.3	-	-	-	-
25-29	1.3	1.3	1.4	0.1	0.0	-
30-34	3.4	3.3	3.0	0.2	0.2	0.3
35-39	6.8	6.6	5.6	0.4	0.4	0.5
40-44	12.8	13.1	8.3	0.9	0.9	1.2
45-49	23.1	23.5	13.3	1.8	1.7	2.1
50-54	42.5	45.1	25.2	3.7	3.6	5.1
55-59	66.9	70.7	44.7	6.7	6.4	9.9
60-64	81.8	85.6	76.1	11.2	10.2	21.5
65-69	89.1	93.9	89.1	16.0	14.0	34.3
70-74	86.5	91.0	88.8	20.5	18.8	41.6
75-79	86.6	90.6	88.9	23.9	22.6	42.7
80-84	81.8	85.3	72.0	28.7	27.7	45.9
85+	61.4	62.9	59.2	34.0	33.6	44.2

^a SEER 17 areas. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^b NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130), unless noted.

^c Rates are per 100,000 and are age-adjusted to the IARC world standard population.

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VII-5
CORPUS AND UTERUS, NOS CANCER (Invasive)

SURVIVAL RATES, BY RACE, DIAGNOSIS YEAR, STAGE AND AGE

	All Races, Females			White Females			Black Females		
	All	<50	50+	All	<50	50+	All	<50	50+
<u>5-YR RELATIVE SURVIVAL RATES</u>									
YEAR OF DIAGNOSIS:									
1960-1963 ^a	-	-	-	73	-	-	31	-	-
1970-1973 ^a	-	-	-	81	-	-	44	-	-
1975-1977 ^b	87.6	92.7	87.0	88.7	93.3	88.2	61.0	84.4 ^f	56.1
1978-1980 ^b	83.6	91.7	82.5	84.8	93.1	83.8	55.8	82.0 ^f	50.3
1981-1983 ^b	81.9	91.3	80.6	83.3	92.3	82.2	52.5	82.8 ^f	47.2
1984-1986 ^b	83.5	93.6	82.1	85.0	93.9	83.8	57.8	83.6 ^f	54.0
1987-1989 ^b	83.4	90.2	82.3	85.0	91.0	84.1	58.3	84.6 ^f	54.1
1990-1992 ^b	84.5	89.9	83.6	86.7	92.1	85.8	54.9	72.1 ^f	51.1
1993-1995 ^b	84.0	90.1	82.9	85.7	91.6	84.7	60.1	82.0	55.2
1996-2003 ^b	84.3 ^e	89.5 ^e	83.2 ^e	86.2 ^e	90.8	85.4 ^e	61.0	76.7	57.0
<u>5-YR PERIOD SURVIVAL RATES^{cd}</u>									
YEAR OF DIAGNOSIS:									
1996-2003	83.0	89.2	81.7	84.9	90.3	83.8	60.7	78.0	56.7
<u>STAGE DISTRIBUTION (%) 1996-2003^c</u>									
All Stages									
Number of cases	46,771	7,354	39,417	39,689	5,682	34,007	3,452	568	2,884
Percent	100%	100%	100%	100%	100%	100%	100%	100%	100%
Localized	70	73	69	71	73	71	54	66	52
Regional	17	16	17	17	16	17	23	16	24
Distant	9	7	9	8	7	8	16	11	17
Unstaged	4	4	4	4	4	4	7	7	7
<u>5-YR RELATIVE SURVIVAL RATES, 1996-2003^c</u>									
AGE AT DIAGNOSIS:									
<45	90.0	-	-	90.7	-	-	82.2	-	-
45-54	88.5	-	-	89.5	-	-	70.7	-	-
55-64	86.0	-	-	88.1	-	-	62.6	-	-
65-74	80.1	-	-	82.9	-	-	52.7	-	-
75+	71.7	-	-	73.9	-	-	45.2	-	-
Under 65	87.5	-	-	89.0	-	-	68.8	-	-
65 and over	76.8	-	-	79.3	-	-	50.3	-	-
STAGE:									
All Stages	82.9	89.1	81.7	84.7	89.9	83.7	60.8	78.6	56.8
Localized	95.3	96.2	95.2	96.1	96.4	96.1	83.2	89.7	81.4
Regional	67.4	80.8	64.9	69.8	83.5	67.4	44.3	63.9 ^f	41.6
Distant	23.1	37.5	20.5	24.9	38.7	22.7	12.7	29.3 ^f	10.6
Unstaged	55.8	85.9	49.2	55.1	85.8	49.1	42.3	78.0 ^f	34.0

^a Rates are based on End Results data from a series of hospital registries and one population-based registry.

^b Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta).

^c Rates are based on follow-up of patients into 2004.

^d Rates are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey). California excluding SF/SJM/LA, Kentucky, Louisiana, and New Jersey contribute cases for diagnosis years 2000-2003. The remaining 13 SEER Areas contribute cases for the entire period 1996-2003. Rates are based on follow-up of patients into 2004.

^e Period survival provides a more up-to-date estimate of survival by piecing together the most recent conditional survival estimates from several cohorts. It is computed here using three calendar year blocks (2001-2003: 0-1 year survival), (2000-2002: 1-2 year survival), (1999-2001: 2-3 year survival), (1998-2000: 3-4 year survival), (1997-1999: 4-5 years survival).

^f The difference in rates between 1975-1977 and 1996-2003 is statistically significant ($p < .05$).

The standard error of the survival rate is between 5 and 10 percentage points.

The standard error of the survival rate is greater than 10 percentage points.

- Statistic could not be calculated due to fewer than 16 cases during the time period.

Table VII-6
CORPUS AND UTERUS, NOS CANCER (Invasive)

SURVIVAL RATES^a

By Year of Diagnosis

All Races, Females

Year of Diagnosis

	1975-		1980-																			
	1979	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	
Relative Survival Rates (SEER^b)																						
1-year	94.0	91.8	93.2	93.0	92.1	92.4	92.2	92.8	92.1	93.1	92.6	92.0	93.3	91.9	93.3	93.7	94.2	94.0	92.5	92.3	93.2	
2-year	90.4	86.8	88.3	87.1	87.5	88.4	88.0	88.3	88.3	89.8	88.8	87.5	88.8	88.5	89.4	89.9	89.8	89.8	88.6	88.4		
3-year	88.2	84.2	85.6	85.4	84.6	85.5	85.9	85.6	86.0	87.6	86.2	85.2	86.5	86.4	86.6	87.0	87.8	87.8	87.8	86.4		
4-year	86.9	83.0	83.9	84.1	83.8	83.9	84.9	84.2	84.3	86.2	84.9	83.7	85.3	85.5	85.8	85.2	86.3	86.8				
5-year	86.6	82.1	83.6	83.1	82.7	83.3	84.2	83.5	84.0	86.1	84.0	83.1	84.9	84.3	84.9	84.3	85.1					
6-year	86.2	81.7	83.1	82.7	81.8	82.9	83.7	82.4	82.9	85.3	83.4	82.1	84.4	83.6	84.1	83.7						
7-year	86.1	81.5	82.4	82.4	81.8	82.7	83.7	82.4	82.8	85.3	83.3	81.4	84.0	83.0	83.6							
8-year	86.1	81.1	82.1	82.4	81.8	82.7	82.8	82.3	82.2	85.0	82.4	81.0	83.8	82.4								
9-year	86.1	81.0	81.9	82.4	81.8	82.5	82.8	82.3	82.1	85.0	82.3	81.0	83.2									
10-year	86.1	80.9	81.9	82.3	81.8	82.0	82.8	81.8	82.1	85.0	81.8	81.0										
11-year	86.1	80.7	81.9	82.3	81.8	82.0	82.7	81.8	82.1	85.0	81.4											
12-year	86.1	80.7	81.9	82.3	81.8	82.0	82.1	81.6	81.8	84.9												
13-year	86.1	80.7	81.7	82.3	81.8	81.9	81.6	81.0	81.4													
14-year	86.1	80.6	81.4	82.3	81.8	81.7	81.4	81.0														
15-year	86.1	80.6	81.4	82.3	80.8	81.0	81.3															
16-year	86.1	80.1	81.2	82.3	80.6	81.0																
17-year	86.1	80.1	80.8	81.6	80.6																	
18-year	86.1	79.9	80.5	81.6																		
19-year	86.1	79.4	80.5																			
20-year	86.1	79.3																				

^a Survival rates are relative rates expressed as percents.

^b Rates are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).

Table VII-7

CORPUS AND UTERUS, NOS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
All Races	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	2.45 (2.42, 2.48)	0.51 (0.51, 0.52)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.06 (0.06, 0.07)	2.48 (2.45, 2.51)	0.52 (0.51, 0.53)	
	20	0.01 (0.01, 0.01)	0.06 (0.06, 0.07)	0.24 (0.24, 0.25)	2.49 (2.46, 2.52)	0.52 (0.51, 0.53)	
	30	0.05 (0.05, 0.06)	0.24 (0.23, 0.24)	0.75 (0.74, 0.77)	2.49 (2.46, 2.52)	0.52 (0.52, 0.53)	
	40	0.18 (0.18, 0.19)	0.71 (0.69, 0.72)	1.44 (1.42, 1.46)	2.46 (2.43, 2.49)	0.52 (0.52, 0.53)	
	50	0.53 (0.52, 0.54)	1.29 (1.27, 1.31)	1.92 (1.89, 1.94)	2.33 (2.30, 2.36)	0.52 (0.51, 0.53)	
	60	0.79 (0.78, 0.81)	1.46 (1.43, 1.48)	1.81 (1.78, 1.84)	1.89 (1.86, 1.92)	0.49 (0.49, 0.50)	
	70	0.75 (0.73, 0.76)	1.15 (1.12, 1.17)	- (- , -)	1.23 (1.21, 1.26)	0.41 (0.40, 0.42)	
	80	0.54 (0.52, 0.56)	- (- , -)	- (- , -)	0.66 (0.64, 0.68)	0.29 (0.28, 0.30)	
White	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	2.56 (2.53, 2.60)	0.49 (0.48, 0.50)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.06 (0.06, 0.07)	2.59 (2.55, 2.62)	0.49 (0.49, 0.50)	
	20	0.01 (0.01, 0.01)	0.06 (0.06, 0.07)	0.25 (0.24, 0.25)	2.59 (2.56, 2.63)	0.49 (0.49, 0.50)	
	30	0.05 (0.05, 0.06)	0.24 (0.23, 0.25)	0.78 (0.77, 0.80)	2.60 (2.56, 2.63)	0.50 (0.49, 0.50)	
	40	0.19 (0.18, 0.19)	0.74 (0.72, 0.75)	1.51 (1.48, 1.53)	2.56 (2.53, 2.60)	0.50 (0.49, 0.51)	
	50	0.56 (0.55, 0.57)	1.35 (1.32, 1.37)	2.01 (1.98, 2.04)	2.43 (2.39, 2.46)	0.49 (0.49, 0.50)	
	60	0.83 (0.81, 0.84)	1.52 (1.49, 1.55)	1.88 (1.85, 1.92)	1.96 (1.93, 1.99)	0.46 (0.46, 0.47)	
	70	0.78 (0.76, 0.80)	1.19 (1.17, 1.22)	- (- , -)	1.28 (1.25, 1.31)	0.39 (0.38, 0.40)	
	80	0.56 (0.54, 0.58)	- (- , -)	- (- , -)	0.67 (0.65, 0.70)	0.28 (0.27, 0.29)	
Black	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	1.95 (1.86, 2.03)	0.75 (0.73, 0.78)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	0.05 (0.04, 0.06)	1.99 (1.90, 2.08)	0.77 (0.74, 0.80)	
	20	0.01 (0.01, 0.02)	0.05 (0.04, 0.06)	0.16 (0.15, 0.18)	1.99 (1.91, 2.09)	0.77 (0.74, 0.80)	
	30	0.04 (0.04, 0.05)	0.15 (0.14, 0.17)	0.50 (0.46, 0.53)	2.00 (1.91, 2.09)	0.78 (0.75, 0.80)	
	40	0.11 (0.10, 0.13)	0.46 (0.43, 0.49)	1.16 (1.10, 1.22)	1.99 (1.90, 2.08)	0.79 (0.76, 0.81)	
	50	0.36 (0.33, 0.39)	1.09 (1.03, 1.15)	1.66 (1.58, 1.74)	1.96 (1.87, 2.05)	0.80 (0.77, 0.83)	
	60	0.79 (0.74, 0.85)	1.41 (1.33, 1.50)	1.69 (1.60, 1.78)	1.74 (1.65, 1.84)	0.79 (0.76, 0.82)	
	70	0.74 (0.68, 0.81)	1.07 (0.99, 1.16)	- (- , -)	1.14 (1.06, 1.23)	0.63 (0.60, 0.66)	
	80	0.49 (0.43, 0.56)	- (- , -)	- (- , -)	0.59 (0.51, 0.67)	0.41 (0.38, 0.44)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VII-7 - continued

CORPUS AND UTERUS, NOS CANCER (Invasive)

Risk of Being Diagnosed With Cancer in 10, 20 and 30 Years, Lifetime Risk of Being Diagnosed with Cancer, and
Lifetime Risk of Dying from Cancer Given Cancer Free At Current Age
Females, 2002-2004 By Race/Ethnicity

Race/ Ethnicity	Current Age	Risk of Being Diagnosed with Cancer				Risk of Dying from Cancer	
		+10 yrs Percent (95% C.I.)	+20 yrs Percent (95% C.I.)	+30 yrs Percent (95% C.I.)	Eventually Percent (95% C.I.)	Eventually Percent (95% C.I.)	
Asian/ Pacific	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.02)	1.77 (1.67, 1.88)	0.34 (0.30, 0.41)	
Islander	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.08 (0.07, 0.09)	1.78 (1.68, 1.90)	0.35 (0.30, 0.41)	
	20	0.01 (0.01, 0.01)	0.08 (0.07, 0.09)	0.30 (0.28, 0.33)	1.78 (1.69, 1.90)	0.35 (0.30, 0.41)	
	30	0.07 (0.06, 0.08)	0.30 (0.27, 0.32)	0.73 (0.69, 0.77)	1.78 (1.68, 1.90)	0.35 (0.30, 0.41)	
	40	0.23 (0.21, 0.25)	0.66 (0.62, 0.71)	1.13 (1.07, 1.19)	1.72 (1.62, 1.84)	0.35 (0.30, 0.41)	
	50	0.44 (0.41, 0.48)	0.91 (0.85, 0.97)	1.25 (1.17, 1.32)	1.51 (1.41, 1.62)	0.33 (0.29, 0.40)	
	60	0.48 (0.44, 0.53)	0.83 (0.77, 0.89)	1.04 (0.97, 1.13)	1.10 (1.01, 1.21)	0.30 (0.25, 0.36)	
	70	0.37 (0.33, 0.42)	0.60 (0.54, 0.67)	- (- , -)	0.66 (0.58, 0.77)	0.22 (0.17, 0.28)	
	80	0.28 (0.22, 0.34)	- (- , -)	- (- , -)	0.35 (0.27, 0.46)	0.14 (0.09, 0.21)	
American	0	0.00 (0.00, 0.04)	0.00 (0.00, 0.04)	0.02 (0.00, 0.06)	1.46 (1.14, 2.05)	0.29 (0.21, 0.45)	
Indian/ Alaska Native ^a	10	0.00 (0.00, 0.02)	0.02 (0.00, 0.05)	0.08 (0.04, 0.14)	1.48 (1.16, 2.08)	0.29 (0.21, 0.46)	
	20	0.02 (0.00, 0.05)	0.08 (0.04, 0.14)	0.23 (0.16, 0.32)	1.49 (1.17, 2.09)	0.29 (0.21, 0.46)	
	30	0.07 (0.03, 0.12)	0.21 (0.15, 0.30)	0.50 (0.38, 0.65)	1.49 (1.16, 2.10)	0.30 (0.21, 0.46)	
	40	0.15 (0.10, 0.22)	0.44 (0.33, 0.59)	0.88 (0.68, 1.13)	1.45 (1.12, 2.07)	0.30 (0.21, 0.47)	
	50	0.30 (0.21, 0.43)	0.75 (0.56, 1.00)	1.11 (0.85, 1.44)	1.33 (1.01, 1.97)	0.28 (0.20, 0.46)	
	60	0.48 (0.32, 0.69)	0.86 (0.61, 1.18)	1.03 (0.74, 1.43)	1.09 (0.77, 1.75)	0.27 (0.18, 0.46)	
	70	0.44 (0.25, 0.71)	0.64 (0.38, 1.03)	- (- , -)	0.71 (0.42, 1.43)	0.23 (0.14, 0.44)	
	80	0.27 (0.10, 0.65)	- (- , -)	- (- , -)	0.36 (0.13, 1.28)	0.13 (0.06, 0.40)	
Hispanic ^b	0	0.00 (0.00, 0.00)	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	1.97 (1.88, 2.08)	0.45 (0.41, 0.48)	
	10	0.00 (0.00, 0.00)	0.01 (0.01, 0.01)	0.07 (0.06, 0.08)	1.99 (1.89, 2.10)	0.45 (0.42, 0.49)	
	20	0.01 (0.01, 0.01)	0.07 (0.06, 0.07)	0.24 (0.22, 0.25)	1.99 (1.90, 2.10)	0.45 (0.42, 0.49)	
	30	0.06 (0.05, 0.06)	0.23 (0.21, 0.24)	0.63 (0.60, 0.66)	1.99 (1.89, 2.10)	0.45 (0.42, 0.49)	
	40	0.17 (0.16, 0.18)	0.58 (0.54, 0.61)	1.10 (1.05, 1.16)	1.95 (1.85, 2.06)	0.45 (0.42, 0.49)	
	50	0.41 (0.39, 0.44)	0.95 (0.89, 1.00)	1.44 (1.37, 1.52)	1.80 (1.71, 1.92)	0.44 (0.41, 0.48)	
	60	0.55 (0.51, 0.60)	1.07 (1.00, 1.14)	1.36 (1.27, 1.45)	1.45 (1.35, 1.55)	0.42 (0.38, 0.46)	
	70	0.56 (0.51, 0.62)	0.88 (0.81, 0.96)	- (- , -)	0.98 (0.89, 1.08)	0.35 (0.31, 0.39)	
	80	0.40 (0.34, 0.47)	- (- , -)	- (- , -)	0.52 (0.44, 0.63)	0.26 (0.22, 0.30)	

Devcan Version 6.2.0, April 2006, National Cancer Institute (<http://srab.cancer.gov/devcan/>).

Source: Incidence data are from the SEER 17 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

Mortality data are from the NCHS public use data file for the total US.

^a Underlying incidence and mortality data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^b Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Underlying incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. Underlying mortality data for Hispanics exclude deaths from Minnesota, New Hampshire and North Dakota.

- Statistic could not be calculated.

A percent or confidence interval value of 0.00 represents a value that is below 0.005.

Table VII-8
CORPUS AND UTERUS, NOS CANCER (Invasive)

SEER INCIDENCE AND U.S. MORTALITY
AGE-ADJUSTED RATES AND TRENDS^a
By Race/Ethnicity

SEER Incidence

RACE/ETHNICITY	Rate 2000-2004 ^b Rate per 100,000 persons	Trend 1995-2004 ^c APC (%)
	Females	Females
All Races	23.2	-0.8*
White	24.3	-1.0*
White Hispanic ^d	17.8	0.7
White Non-Hispanic ^d	25.3	-1.0*
Black	19.6	1.4
Asian/Pacific Islander	16.2	0.2
Amer Ind/Alaska Nat ^e	15.1	-
Hispanic ^d	17.4	0.8

U.S. Mortality^f

RACE/ETHNICITY	Rate 2000-2004 Rate per 100,000 persons	Trend 1995-2004 APC (%)
	Females	Females
All Races	4.1	0.1
White	3.9	0.0
White Hispanic ^d	3.2	-0.6
White Non-Hispanic ^d	3.9	0.0
Black	7.1	0.6
Asian/Pacific Islander	2.4	1.1
Amer Ind/Alaska Nat		
Total U.S.	2.3	-1.2
CHSDA Counties	2.9	-
Non-CHSDA Counties	1.6	-
Hispanic ^d	3.1	-0.5

- The APC is the Annual Percent Change over the time interval.

- Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Incidence data used in calculating the rates are from the 17 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).

^c Incidence data used in calculating the trends are from the 13 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).

^d Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry and Kentucky. The 2000-2004 Hispanic and Non-Hispanic death rates exclude deaths from Minnesota, New Hampshire and North Dakota. The 1995-2004 Hispanic and Non-Hispanic mortality trends exclude deaths from Maine, Minnesota, New Hampshire, North Dakota, and Oklahoma.

^e Incidence data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^f Mortality data are analyzed from a public use file provided by the National Center for Health Statistics (NCHS).

* The APC is significantly different from zero ($p < .05$).

Table VII-9
CORPUS AND UTERUS, NOS CANCER (Invasive)

AGE-ADJUSTED SEER INCIDENCE RATES^a
By Registry, Race/Ethnicity and Age

	All Races			Whites			Blacks		
	All Ages	Ages <50	Ages 50+	All Ages	Ages <50	Ages 50+	All Ages	Ages <50	Ages 50+
SEER INCIDENCE RATES^a, 2000-2004									
REGISTRY									
Atlanta & Rural Georgia	18.8	4.0	57.8	20.2	4.2	62.2	16.7	3.4	51.7
Atlanta	19.2	4.0	58.9	20.6	4.3	63.2	17.1	3.4	53.1
Rural Georgia	13.6	-	42.0	14.3	-	47.0	12.4	-	-
California	21.7	4.7	66.0	22.6	4.6	69.9	19.1	3.3	60.5
Greater Bay Area	22.8	4.5	70.6	24.4	4.0	78.0	20.0	4.5	60.5
San Francisco-Oakland	23.7	4.7	73.3	25.1	3.8	80.8	20.5	4.7	62.0
San Jose-Monterey	21.0	4.1	65.3	23.3	4.3	73.0	15.2	-	-
Los Angeles	21.8	5.3	65.2	23.4	5.4	70.7	19.8	3.6	62.4
Greater California	21.1	4.5	64.7	21.8	4.4	67.2	17.7	2.4	57.8
Connecticut	28.5	5.6	88.4	29.3	5.9	90.6	20.1	3.3	64.1
Detroit	26.4	5.3	81.7	28.0	5.7	86.6	22.4	4.5	69.3
Hawaii	24.8	9.0	66.2	23.9	4.4	74.8	-	-	-
Iowa	27.2	5.7	83.3	27.4	5.8	83.9	-	-	-
Kentucky	23.1	6.5	66.8	23.2	6.5	66.9	19.6	4.8	58.1
Louisiana	17.3	4.5	50.8	17.5	5.6	48.5	18.0	2.4	59.0
New Jersey	28.2	5.1	88.8	29.8	5.4	93.6	22.4	3.6	71.5
New Mexico	19.4	4.9	57.2	19.7	4.6	59.2	-	-	-
Seattle-Puget Sound	24.8	4.7	77.4	26.0	4.9	81.3	17.1	-	58.8
Utah	22.8	4.9	69.5	23.0	4.9	70.6	-	-	-
9 SEER Areas ^b	24.6	5.1	75.5	25.7	5.0	79.8	19.7	3.7	61.8
11 SEER Areas ^b	23.8	5.1	72.6	25.1	5.1	77.5	19.7	3.7	61.8
13 SEER Areas ^b	23.7	5.1	72.4	25.1	5.1	77.4	19.6	3.7	61.4
17 SEER Areas ^b	23.2	5.0	71.0	24.3	5.0	74.6	19.6	3.3	62.1

^a Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130)
^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.
The SEER 11 areas comprise the SEER 9 areas plus San Jose-Monterey and Los Angeles.
The SEER 13 areas comprise the SEER 11 areas plus the Alaska Native Registry and Rural Georgia.
The SEER 17 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VII-10
CORPUS AND UTERUS, NOS CANCER (Invasive)

AGE-ADJUSTED SEER DEATH RATES^a
By Registry, Race/Ethnicity and Age

	All Races			Whites			Blacks		
	All Ages	Ages <50	Ages 50+	All Ages	Ages <50	Ages 50+	All Ages	Ages <50	Ages 50+
<u>SEER DEATH RATES^a, 2000-2004</u>									
REGISTRY									
Atlanta & Rural Georgia	3.9	-	13.4	3.1	-	10.7	5.9	-	20.6
Atlanta	3.8	-	13.3	3.1	-	10.7	5.8	-	20.2
Rural Georgia	4.1	-	14.7	-	-	-	-	-	-
California	3.9	0.4	13.1	3.9	0.3	13.1	7.2	0.5	24.7
Greater Bay Area	4.0	0.3	13.5	4.1	0.2	14.1	7.0	-	24.4
San Francisco-Oakland	4.1	0.4	13.8	4.1	-	14.2	7.3	-	25.5
San Jose-Monterey	3.7	-	12.8	4.0	-	13.9	-	-	-
Los Angeles	4.2	0.4	14.4	4.2	0.4	14.1	7.7	-	26.6
Greater California	3.7	0.4	12.3	3.7	0.4	12.4	6.5	-	22.1
Connecticut	4.0	0.3	13.7	3.9	0.3	13.4	6.3	-	20.9
Detroit	4.9	0.5	16.5	4.3	0.3	14.6	7.2	1.0	23.6
Hawaii	3.8	0.8	11.8	2.7	-	8.8	-	-	-
Iowa	4.4	0.5	14.4	4.4	0.5	14.5	-	-	-
Kentucky	3.6	0.4	12.0	3.5	0.4	11.4	6.6	-	23.4
Louisiana	3.6	0.4	11.7	2.5	0.4	7.8	7.0	-	24.1
New Jersey	5.2	0.3	17.8	5.0	0.4	17.2	7.5	-	26.3
New Mexico	3.8	0.5	12.5	3.8	-	12.8	-	-	-
Seattle-Puget Sound	4.0	0.3	13.7	4.1	0.3	13.8	6.3	-	22.8
Utah	4.3	-	14.9	4.3	-	15.0	-	-	-
9 SEER Areas ^b	4.2	0.4	14.1	4.0	0.3	13.7	6.6	0.6	22.3
11 SEER Areas ^b	4.2	0.4	14.1	4.1	0.3	13.8	6.9	0.6	23.4
13 SEER Areas ^b	4.2	0.4	14.1	4.0	0.3	13.8	6.9	0.6	23.4
17 SEER Areas ^b	4.1	0.4	13.9	4.0	0.3	13.4	7.0	0.5	23.9
Total U.S.	4.1	0.4	14.0	3.9	0.3	13.1	7.1	0.4	24.5

^a NCHS public use data file for the total US. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

The SEER 11 areas comprise the SEER 9 areas plus San Jose-Monterey and Los Angeles.

The SEER 13 areas comprise the SEER 11 areas plus the Alaska Native Registry and Rural Georgia.

The SEER 17 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey

- Statistic not shown. Rate based on less than 16 cases for the time interval.

Table VII-11
CORPUS AND UTERUS, NOS CANCER (Invasive)

AVERAGE ANNUAL AGE-ADJUSTED CANCER DEATH^a RATES BY STATE, ALL RACES, 2000-2004

Females

<u>State</u>	<u>Rate</u>	<u>SE</u>	<u>Rank</u>	<u>PD</u>	<u>State</u>	<u>Rate</u>	<u>SE</u>	<u>Rank</u>	<u>PD</u>
TOTAL U.S.	4.1	0.02							
High Five States									
District of Columbia	7.4 ^b	0.67	(01)	80.5 ^c					
Vermont	5.8	0.56	(02)	41.5 ^c					
New Jersey	5.2 ^b	0.14	(03)	26.8 ^c					
Nebraska	5.1	0.32	(04)	24.4 ^c					
New York	4.8 ^b	0.09	(05)	17.1 ^c					
Low Five States									
Idaho	3.3	0.31	(47)	-19.5 ^c					
Alaska	3.3	0.61	(48)	-19.5 ^c					
Nevada	3.0 ^b	0.24	(49)	-26.8 ^c					
Rhode Island	3.0 ^b	0.30	(50)	-26.8 ^c					
Arizona	2.8 ^b	0.14	(51)	-31.7 ^c					
State	Rate	SE	Rank	PD	State	Rate	SE	Rank	PD
Alabama	3.6	0.16	(40)	-12.2	Montana	4.6	0.41	(09)	12.2
Alaska	3.3	0.61	(48)	-19.5 ^c	Nebraska	5.1	0.32	(04)	24.4 ^c
Arizona	2.8 ^b	0.14	(51)	-31.7 ^c	Nevada	3.0 ^b	0.24	(49)	-26.8 ^c
Arkansas	3.5	0.20	(45)	-14.6	New Hampshire	3.9	0.33	(31)	-4.9
California	3.9 ^b	0.07	(30)	-4.9	New Jersey	5.2 ^b	0.14	(03)	26.8 ^c
Colorado	3.5 ^b	0.18	(46)	-14.6	New Mexico	3.8	0.28	(36)	-7.3
Connecticut	4.0	0.19	(27)	-2.4	New York	4.8 ^b	0.09	(05)	17.1 ^c
Delaware	4.7	0.45	(08)	14.6	North Carolina	4.0	0.13	(26)	-2.4
District of Columbia	7.4 ^b	0.67	(01)	80.5 ^c	North Dakota	4.6	0.47	(10)	12.2
Florida	3.6 ^b	0.08	(43)	-12.2	Ohio	4.5	0.11	(13)	9.8
Georgia	3.8	0.14	(34)	-7.3	Oklahoma	3.9	0.19	(32)	-4.9
Hawaii	3.8	0.33	(35)	-7.3	Oregon	3.9	0.20	(29)	-4.9
Idaho	3.3	0.31	(47)	-19.5 ^c	Pennsylvania	4.8 ^b	0.11	(06)	17.1 ^c
Illinois	4.5	0.11	(11)	9.8	Rhode Island	3.0 ^b	0.30	(50)	-26.8 ^c
Indiana	4.8 ^b	0.16	(07)	17.1 ^c	South Carolina	3.8	0.18	(33)	-7.3
Iowa	4.4	0.22	(17)	7.3	South Dakota	4.3	0.43	(21)	4.9
Kansas	4.0	0.23	(28)	-2.4	Tennessee	3.7	0.15	(39)	-9.8
Kentucky	3.6	0.17	(42)	-12.2	Texas	3.7 ^b	0.09	(38)	-9.8
Louisiana	3.6	0.17	(44)	-12.2	Utah	4.3	0.31	(19)	4.9
Maine	4.5	0.33	(12)	9.8	Vermont	5.8	0.56	(02)	41.5 ^c
Maryland	4.5	0.17	(14)	9.8	Virginia	4.2	0.15	(23)	2.4
Massachusetts	4.4	0.15	(18)	7.3	Washington	4.1	0.16	(25)	0.0
Michigan	4.4	0.12	(15)	7.3	West Virginia	4.3	0.27	(22)	4.9
Minnesota	4.2	0.17	(24)	2.4	Wisconsin	4.4	0.17	(16)	7.3
Mississippi	3.8	0.22	(37)	-7.3	Wyoming	3.6	0.52	(41)	-12.2
Missouri	4.3	0.16	(20)	4.9					

^a NCHS public use data file. Rates are per 100,000 and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

^b Difference between state rate and total U.S. rate is statistically significant ($p \leq .0002$).

^c Absolute percent difference between state rate and total U.S. rate is 15% or more.

SE Standard error of the rate.

PD Percent difference between state rate and total U.S. rate.

Table VII-12
CORPUS AND UTERUS, NOS CANCER (Invasive)

Estimated United States Cancer Prevalence Counts^a on January 1, 2004
By Race/Ethnicity and Years Since Diagnosis

Years Since Diagnosis	0 to <5	5 to <10	10 to <15	15 to <20	20 to <25	0 to <14 ^e	0 to <29 ^e	>=29 ^g	Complete ^h
<u>Race</u>									
All Races ^b	144,321	111,240	84,444	64,539	52,023	325,042	500,188	68,219	568,407
White ^b	128,644	101,702	79,124	60,661	49,320	295,404	461,176	60,537	521,713
Black ^b	9,355	5,220	3,091	2,276	1,376	17,120	22,275	1,425	23,700
Asian/Pacific Islander ^c	3,867	2,512	+	+	+	7,637	+	+	+
Hispanic ^d	7,272	4,561	+	+	+	14,687	+	+	+

Estimated prevalence percent^a on January 1, 2004, of the SEER 11 population diagnosed in the previous 10 years
By Age at Prevalence and Race/Ethnicity

Age at Prevalence	Age Specific (Crude)									Age-Adjusted ^f All Ages
	All Ages	0-9	10-19	20-29	30-39	40-49	50-59	60-69	70-79	
<u>Race</u>										
All Races ^c	0.1522%	-	-	0.0026%	0.0207%	0.0848%	0.2844%	0.5902%	0.6642%	0.5303%
White ^c	0.1730%	-	-	0.0023%	0.0210%	0.0850%	0.3045%	0.6534%	0.7410%	0.5792%
Black ^c	0.0712%	-	-	0.0032%	0.0173%	0.0504%	0.1358%	0.3784%	0.4291%	0.2874%
Asian/Pacific Islander ^c	0.1039%	-	-	0.0028%	0.0199%	0.1070%	0.2752%	0.3438%	0.3162%	0.2382%
Hispanic ^d	0.0602%	-	-	0.0030%	0.0266%	0.0769%	0.2041%	0.3760%	0.3559%	0.2709%

^a US 2004 cancer prevalence counts are based on 2004 cancer prevalence proportions from the SEER registries and 1/1/2004 US population estimates based on the average of 2003 and 2004 population estimates from the US Bureau of the Census. Prevalence was calculated using the First Malignant Primary Only for a person.

^{b c d} Statistics based on (b) SEER 9 Areas (c) SEER 11 Areas and Rural Georgia (d) NHIA for Hispanic for SEER 11 Areas and Rural Georgia.

^e Maximum limited-duration prevalence: 29 years for 1975-2004 SEER 9 data; 14 years for 1990-2004 SEER 11 data (used to calculate prevalence for Hispanics and Asian Pacific Islanders).

^{f g h} Percentages are age-adjusted to the 2000 US Standard Population (19 age groups - Census P25-1130) by 5-year age groups.

^(g) Cases diagnosed more than 29 years ago were estimated using the completeness index method (Capocaccia et. al. 1997, Merrill et. al. 2000). (h)Complete prevalence is obtained by summing 0 to <29 and >=29.

⁻ Statistic not shown. Statistic based on fewer than 5 cases estimated alive in SEER for the time interval.

⁺ Not available.

SEER Incidence, Delay Adjusted Incidence and US Death Rates^a Corpus and Uterus, NOS Cancer, by Race




Figure VII-1

^a Source: SEER 9 areas and NCHS public use data file for the total US. Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines and APCs are calculated using the Joinpoint Regression Program Version 3.0, April 2005, National Cancer Institute.

The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

SEER Incidence, Delay Adjusted Incidence and US Death Rates^a Corpus and Uterus Cancer, by Age and Race




Figure VII-2

^a Source: SEER 9 areas and NCHS public use data file for the total US. Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103). Regression lines and APCs are calculated using the Joinpoint Regression Program Version 3.0, April 2005, National Cancer Institute.


The APC is the Annual Percent Change for the regression line segments. The APC shown on the graph is for the most recent trend.

* The APC is significantly different from zero ($p < 0.05$).

SEER Incidence and US Death Rates^a

Corpus and Uterus, NOS Cancer

Joinpoint Analyses for Whites and Blacks from 1975-2004
and for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics from 1992-2004



Source: Incidence data for whites and blacks are from the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta). Incidence data for Asian/Pacific Islanders, American Indians/Alaska Natives and Hispanics are from the SEER 13 Areas (SEER 9 Areas, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia). Mortality data are from NCHS public use data file for the total US.

^a Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1103).

Regression lines are calculated using the Joinpoint Regression Program Version 3.0, April 2005, National Cancer Institute.

^b API = Asian/Pacific Islander.

^c AI/AN = American Indian/Alaska Native. Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

^d Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry. Mortality data for Hispanics exclude cases from Connecticut, Maine, Maryland, Minnesota, New Hampshire, New York, North Dakota, Oklahoma, and Vermont.

Figure VII-3