

References

- Baskettter DA, Scholes EW. 1992. Comparison of the local lymph node assay with the guinea-pig maximization test for the detection of a range of contact allergens. *Food and Chemical Toxicology* 30(1): 65-69.
- Baskettter DA, Scholes EW, Kimber I. 1994. The performance of the local lymph node assay with chemicals identified as contact allergens in the human maximization test. *Food and Chemical Toxicology* 32(6): 543-547.
- Baskettter DA, Lea LJ, Dickens A, Briggs D, Pate I, Dearman RJ, et al. 1999a. A comparison of statistical approaches to the derivation of EC3 values from local lymph node assay dose responses. *Journal of Applied Toxicology* 19(4): 261-266.
- Baskettter DA, Lea LJ, Cooper K, Stocks J, Dickens A, Pate I, et al. 1999b. Threshold for classification as a skin sensitizer in the local lymph node assay: A statistical evaluation. *Food and Chemical Toxicology* 37(12): 1167-1174.
- Baskettter DA, Lea LJ, Cooper KJ, Ryan CA, Gerberick GF, Dearman RJ, et al. 1999c. Identification of metal allergens in the local lymph node assay. *American Journal of Contact Dermatitis* 10(4): 207-212.
- Baskettter DA, Smith Pease CK, Patlewicz GY. 2003. Contact allergy: The local lymph node assay for the prediction of hazard and risk. *Clinical and Experimental Dermatology* 28(2): 218-221.
- Baskettter DA, Kimber I. 2006. Predictive tests for irritants and allergens and their use in quantitative risk assessment. *Contact Dermatitis*, 4th Edition Frosch P J, Menne T, Lepoittevin J-P (eds): Heidelberg, Springer Verlag: 179-188.
- Bourrinet P, Puchault P, Sarrazin G, Berkovic A. 1979. Comparative investigation of allergenic products in man and in guinea-pig. *J Pharm Belg* 34(1): 21-26.
- De Jong WH, Van Och FMM, Den Hartog Jager CF, Spiekstra SW, Slob W, Vandebriel RJ, et al. 2002. Ranking of allergenic potency of rubber chemicals in a modified local lymph node assay. *Toxicological Sciences* 66(2): 226-232.
- Gad SC, Dunn BJ, Dobbs DW. 1986. Development and validation of an alternative dermal sensitization test: The mouse ear swelling test (MEST). *Toxicology and Applied Pharmacology* 84(1): 93-114.
- Gerberick GF, House RV, Fletcher ER, Ryan CA. 1992. Examination of the local lymph node assay for use in contact sensitization risk assessment. *Fundamental and Applied Toxicology* 19(3): 438-445.
- Goodwin BFJ, Crevel RWR, Johnson AW. 1981. A comparison of three guinea-pig sensitization procedures for the detection of 19 reported human contact sensitizers. *Contact Dermatitis* 7(5): 248-258.
- Griem P, Goebel C, Scheffler H. 2003. Proposal for a risk assessment methodology for skin sensitization based on sensitization potency data. *Regulatory Toxicology and Pharmacology* 38(3): 269-290.

- Hicks R, Hewitt PJ, Lam HF. 1979. An investigation of the experimental induction of hypersensitivity in the guinea pig by material containing chromium, nickel and cobalt from arc welding fumes. International Archives of Allergy and Applied Immunology 59(3): 265-272.
- ICCVAM. 1991. The Murine Local Lymph Node Assay: A Test Method for Assessing the Allergic Contact Dermatitis Potential of Chemicals/Compounds. National Institutes of Health (NIH) Publication No: 99-4494.
- Ikarashi Y, Tsuchiya T, Nakamura A. 1992. Detection of contact sensitivity of metal salts using the murine local lymph node assay. Toxicology Letters 62(1): 53-61.
- Kimber I, Basketter DA, Butler M, Gamer A, Garrigue JL, Gerberick GF, et al. 2003. Classification of contact allergens according to potency: Proposals. Food and Chemical Toxicology 41(12): 1799-1809.
- Kligman AM. 1966a. The identification of contact allergens by human assay. II. Factors influencing the induction and measurement of allergic contact dermatitis. Journal of Investigative Dermatology 47(5): 375-392.
- Kligman AM. 1966b. The identification of contact allergens by human assay. 3. The maximization test: a procedure for screening and rating contact sensitizers. Journal of Investigative Dermatology 47(5): 393-409.
- Magnusson B, Kligman AM. 1969. The identification of contact allergens by animal assay. The guinea pig maximization test. Journal of Investigative Dermatology 52(3): 268-276.
- Mandervelt C, Clottens FL, Demedts M, Nemery B. 1997. Assessment of the sensitization potential of five metal salts in the murine local lymph node assay. Toxicology 120(1): 65-73.
- Marzulli FN, Maibach HI. 1976a. Contact allergy: predictive testing in man. Contact Dermatitis 2(1): 1-17.
- Maurer T. 1985. The optimization test. Current problems in dermatology 14: 114-151.
- Menne T. 1994. Quantitative aspects of nickel dermatitis. Sensitization and eliciting threshold concentrations. Science of the Total Environment 148(2-3): 275-281.
- Ryan CA, Gerberick GF, Cruse LW, Basketter DA, Lea L, Blaikie L, et al. 2000. Activity of human contact allergens in the murine local lymph node assay. Contact Dermatitis 43(2): 95-102.
- Ryan CA, Cruse LW, Skinner RA, Dearman RJ, Kimber I, Gerberick GF. 2002. Examination of a vehicle for use with water soluble materials in the murine local lymph node assay. Food and Chemical Toxicology 40(11): 1719-1725.
- Schneider K, Akkan Z. 2004. Quantitative relationship between the local lymph node assay and human skin sensitization assays. Regulatory Toxicology and Pharmacology 39(3): 245-255.

Uter W, Fuchs T, Hausser M, Ippen H. 1995. Patch test results with serial dilutions of nickel sulfate (with and without detergent), palladium chloride, and nickel and palladium metal plates. Contact Dermatitis 32(3): 135-142.

Vandenberg JJ, Epstein WL. 1963. Experimental nickel contact sensitization in man. Journal of Investigative Dermatology 167(207-215).

van Och FMM, Vandebriel RJ, Prinsen MK, De Jong WH, Slob W, Van Loveren H. 2001. Comparison of dose-responses of contact allergens using the guinea pig maximization test and the local lymph node assay. Toxicology 167(3): 207-215.

Wahlberg JE, Boman A. 1985. Guinea pig maximization test. Current problems in dermatology 14: 59-106.