



THE IHS PRIMARY CARE PROVIDER



A journal for health professionals working with American Indians and Alaska Natives

January 2007

Volume 32 Number 1

Taking Control of Methamphetamine in Our Communities: An Opportunity of Necessity

*H. Westley Clark, MD, JD, MPH, CAS, FASAM, Director,
Center for Substance Abuse Treatment, Substance Abuse and
Mental Health Services Administration, US Department of
Health and Human Services, Rockville, Maryland*

The Substance Abuse and Mental Health Services Administration (SAMHSA) recognizes that the production and consumption of methamphetamine takes a severe toll on all Americans. However, the affect on American Indians and Alaska Natives (AI/AN) is particularly severe. Thus, it is essential that we continue our partnership with AI/AN communities in an effort to contain and neutralize the impact that methamphetamine use has in Indian Country.

In our effort to address methamphetamine use in Indian Country, SAMHSA partners with Indian Health Service (IHS) as well as other Agencies and Departments within the Federal government. We are an active member of the Office of National Drug Control Policy's Executive Tribal Law Enforcement Workgroup, which includes representatives from the IHS, Department of Justice, Department of Interior, Environmental Protection Agency, Federal Bureau of Investigation, Tribal Law Enforcement, and the Drug Enforcement Agency. The goal of this work group is to address methamphetamine use and production in Indian Country and to help educate AI/AN communities about methamphetamine. We also partner with the National Institute on Drug Abuse in an effort to foster appropriate treatment strategies, and with the Centers for Disease Control and Prevention to address the infectious disease aspects of methamphetamine use and abuse.

Methamphetamine use and abuse has created an opportunity of necessity. Small laboratories are on the decline in many areas, but they still remain a problem. In addition to laboratories, law enforcement agents recognize the need to address the problem of across-the-border smuggling and drug dealing.

While there are no magic solutions to the problem of methamphetamine use and abuse, there is the realization that the health of the AI/AN communities rests on the wisdom of those communities to work together within and outside of Indian Country with people of good will committed to the well-being of these communities. With this partnership, progress can be made and will be made.

Methamphetamine Abuse in Indian Country

Lori De Ravello, MPH, CDC Assignee to the National STD Program, volunteered to coordinate two special issues of *The IHS Provider* dedicated to the problem of methamphetamine abuse in Indian Country. The first issue, a series of articles *defining the problem, appeared in the December 2006 issue*. This issue is devoted to *working together on solutions*. Due to the enthusiastic response of the many authors who volunteered for this project, we have several additional articles that will be published this spring. We wish to recognize Ms. De Ravello's initiative and hard work that made this project possible.

In this Issue...

- 1 Taking Control of Methamphetamine in Our Communities: An Opportunity of Necessity
- 1 Methamphetamine Abuse in Indian Country
- 2 Strength Through Unity: The IHS/SAMHSA Methamphetamine Initiative
- 3 Using the "5 A's" to Address Methamphetamine Use in the Primary Care Setting
- 6 Online Methamphetamine Resources
- 7 Healing to Wellness Courts: Best Practice for Indian Country
- 9 Healing from Methamphetamine: Community Efforts, Recovery, and Healing in American Indian and Alaska Native Communities
- 11 OB/GYN Chief Clinical Consultant's Corner Digest
- 18 2007 Clinical Update on Substance Abuse and Dependency
- 19 IHS Child Health Notes
- 22 Save the Date
- 22 The 11th Annual Elders Issue
- 23 Prevention of Methamphetamine Use in Indian Country: Promising Practices
- 26 Meetings of Interest
- 27 Position Vacancies

Strength Through Unity: The IHS/SAMHSA Methamphetamine Initiative

Wilbur Woodis, Division of Behavioral Health, Indian Health Service, Rockville, Maryland

In spring 2005, IHS and the Substance Abuse and Mental Health Services Administration (SAMHSA) formed the IHS/SAMHSA Methamphetamine Initiative Workgroup. The goal of the workgroup is to help coordinate, collect, analyze, and report information concerning methamphetamine use in Indian Country to a variety of interested parties, including legislators, policy makers, program managers, and persons active in prevention and treatment.

The workgroup has monthly information-sharing calls hosted by the IHS Division of Behavioral Health. Some early methamphetamine-related efforts recognized by the workgroup include:

- SAMHSA Center for Substance Abuse Prevention (CSAP) supported the Pascua Yaqui Tribe to collaborate with the University of Arizona to address substance abuse prevention, focusing on inhalants, methamphetamine, and alcohol abuse.
- The Cherokee Nation formed a Multi-Disciplinary Task Force to address methamphetamine prevention as well as rehabilitation and education.
- SAMHSA made a three-year Target Capacity Expansion grant to the Choctaw Nation to treat methamphetamine-using women with children.
- The SAMHSA-funded Methamphetamine Treatment Project has provided the Billings Area with technical assistance since 1998, when methamphetamine first became a crisis on some of the Area reservations.
- The Matrix Institute trained I/T/U providers on the Ft. Peck and Blackfeet reservations and in Great Falls and Billings in the Matrix model of cognitive behavioral treatment for stimulant use disorders.
- The Phoenix Area developed an Intensive Outpatient Alcohol and Drug Treatment Manual that provides a step-by-step treatment curriculum for methamphetamine addicts with an accompanying Clinician's Guide to Methamphetamine.
- The Phoenix Area is conducting assessments to determine community-level stage of readiness to deal with methamphetamines and to develop strategic plans to address the problem.
- The Navajo Area produced the film, Methamphetamine on the Navajo Reservation "G" and recently the Navajo Nation Council passed

legislation prohibiting the manufacturing, distribution, sales, possession, and use of methamphetamines.

Another activity of the workgroup was to develop a plan to address problems caused by methamphetamine abuse, including aspects related to primary care, mental health, emergency departments, addictions treatment, environmental health, health education, and law enforcement. Some components are underway, while others currently lack resources to implement. As the workgroup continued to meet via conference calls, it began to identify local, regional, and national experts who could provide technical assistance at meetings, conferences, and summits. The workgroup assisted the National Native American Law Enforcement Association (NNALEA) to coordinate the methamphetamine training track at its annual conference in November 2006 and is assisting with the planning of an Aberdeen Area Methamphetamine Summit, scheduled to take place in Fargo, North Dakota in spring 2007.

The workgroup supports the ideals of the HHS "Indian Country Methamphetamine Initiative," many of which were identified through a series of tribal roundtables in 2006. One outcome of the tribal roundtables was a comprehensive plan to address methamphetamine abuse issues across Indian Country. Tribal officials identified many areas of need and concern, including domestic violence, the environmental impact of manufacturing methamphetamine, suicide, increased risk for sexually transmitted diseases, the burden on law enforcement and courts, and drug endangered children.

For information on the IHS/SAMHSA Methamphetamine Initiative Workgroup, visit our website at: <http://www.ihs.gov/MedicalPrograms/Behavioral/index.cfm?module=BH&option=Meth>.



Using the “5 A’s” to Address Methamphetamine Use in the Primary Care Setting

Kathleen Masis MD, CAPT, USPHS (ret), Billings, Montana

Methamphetamine (meth) abuse and dependence, and related behaviors (e.g., injection drug use, interpersonal violence, child abuse and neglect, self-destructive behaviors) in our service populations are creating a public health dilemma for many Indian Health Service/Tribal/Urban primary care clinics. What are evidence-based strategies to address prevention and intervention in our practices?

Other articles in this series on methamphetamine have demonstrated the prevalence of problems from its use among American Indians and Alaska Natives. This article will present a rationale and method for delivering preventive services to a targeted population at risk.

The U.S. Preventive Services Task Force and the American Academy of Family Physicians have issued evidence-based recommendations on the level of intervention indicated for risk behaviors, including tobacco abuse, sedentary lifestyle, unhealthy diet, and alcohol abuse. There is insufficient evidence to recommend for or against screening for illicit drug use, other than alcohol and tobacco, among adolescent patients.¹ The same holds for “screening” adults.

Longitudinal studies have shown that substance abuse treatment results in reduced use of drugs and costs to society.² As primary care providers, our role is to assist our patients to move towards positive behavior change. Unfortunately, methamphetamine users often demonstrate “more than the usual denial.”³

A model recommended for intervention with tobacco users who are willing to quit is the “5 A’s.”⁴ The “5 A’s,” which are applied in order, are Ask, Advise, Assess, Assist, and Arrange. Appropriate types of visits to initiate the Five A’s for methamphetamine use might be Prenatal, Well Woman, New Patient, Adolescent, Behavioral Health (anxiety, depression, insomnia, chronic pain,) STD testing and treatment, Dental, and visits by incarcerated patients.

Ask: Ask If The Patient Is Using

The task is to identify patients currently using methamphetamine.

Provider: “I can see that your rash is bothering you a lot. In order to help you, I need to ask some other questions about your health. (Do a targeted ROS.) Are you taking any medications? When was your last

alcohol use? Methamphetamine, “crystal” or (local terminology)? How do you use it, smoking, or injecting?”

We are familiar with the distinctive and/or common indicators of chronic methamphetamine use: missing teeth, dermatitis around the mouth; muscle cramping with dehydration and abnormal electrolytes, rhabdomyolysis; anorexia and extreme weight loss; pulmonary, cardiovascular, and cerebrovascular disorders, including cognitive deficits; hypertensive crisis; choreoathetosis; and psychiatric and behavioral disorders such as insomnia, paranoia, anxiety, perceptual disturbances (such as bugs under the skin,) delusions and hallucinations, depression, and risk of suicide.⁵

Primary care practitioners are starting to recognize that many methamphetamine users are seen in primary care settings at an earlier stage before these manifestations are apparent. Brief interventions may be more effective at this stage.

Strategies include the following:

- The manner and attitude of the professional doing the asking is as important (even more important) than the specific question(s) or “tool” that is used. A non-judgmental attitude is crucial.
- For patients who are at risk, but deny the behavior, the least intrusive intervention would be indicated; that is general information that targets all community members. The objective is not to “close the door” between the health care provider and the patient.

Example: If the patient denies meth use: “A lot of our patients are using meth. I’ve seen lots of health problems such as itching and rashes, caused by meth, and so I tell all my patients about quitting meth.”

Advise: Give Clear, Strong, and Personalized Behavior Change Advice, Including Information about Personal Health Harms of Meth and Benefits to Quitting

It is important to personalize your advice by including specifics of the patient’s health concerns, including past experiences, family, or social situations. “Education is often the first phase of treatment.”³ Provide feedback to the patient on the information you collected in the first step. For example:

Provider: “You told me that you have used meth for a year, and injected meth only one time, which was last week. You have never shared needles with anyone else . . . You

have symptoms of anxiety and depression, which are affecting your work and family life . . .

“I am concerned about your risks for STD, HIV, and worsening mental state if you continue to use . . .

“As your health care provider, I can tell you that the most important thing you can do for your mental and physical health is to stop using meth and to stop using it now . . .

“Because a lack of awareness or even memory of what a person does while they are high on meth is common, and they end up sharing a dirty needle or having unprotected sex . . .

“Our clinic is interested in helping people get off meth. We have several ways to help.”

Often even a positive drug screen result does not result in admission of use by the patient. Nevertheless, the core message to convey to the patient is: meth is dangerous to your health, it is possible to stop, and your family doctor can help. In fact, “he or she can be your biggest ally.”⁶

Assess: Determine Willingness to Make a Quit Attempt

For example:

Provider: “Are you interested in quitting meth?”

Patient: “I’ve been depressed for as long as I can remember. When I tried meth, I felt normal for the first time. But I have to have clean urines to get custody of my kids back.”

Assist: Assist the Patient in Quitting

The assistance we provide should be tailored to the patient’s perceived needs. Returning to our patient example, above (a composite of actual patient visits):

Provider: “You want to get your depression under control, and to get custody of your kids . . . I am willing to work with you on both depression and meth issues. I suggest we start by assessing both problems and coming up with a plan for each.”

The health care provider may be more helpful and credible if he/she can describe the common course of withdrawal from meth, (See Table 1) and provide the patient with strategies to deal with them in health-promoting ways (See Table 2). Medical detoxification is seldom needed; but monitoring by a supportive friend, family member, or non-medical detox program is warranted: suicide is the most lethal complication.

It is important to have direct knowledge of community resources when making a referral. Encourage the patient to enlist support from non-using family and friends. No

Table 1. Course of Methamphetamine Withdrawal

<i>1, 2, and 3 are common</i>	
1. Early crash	Dysphoria, agitation, craving. Onset 12-24 hrs after last use. “Tweaking” can appear now and/or later, and includes irritability, jittery state, quivery speech, brisk, jerky movements. Thinking seems scattered, subject to paranoid delusions, aggression.
2. Middle crash	Protracted sleep, 24-36 hrs. Fatigue, depression, anhedonia. Decreased mental and physical activity. Desire for sleep, with insomnia (users commonly take alcohol, benzos, opiates to deal with these symptoms).
3. Late crash	Awake and hungry
4. Protracted Fatigue <i>less common</i>	Symptoms opposite to meth intoxication. Withdrawal, loss of physical and mental energy. Depression, anhedonia, limited interest in surroundings. May increase in intensity over 12-96 hours following crash, or wax and wane over several weeks. Severe, persistent depression can result in suicidal behavior. Anhedonia and dysphoria last 6-18 weeks. Periods of intense drug craving, “triggers.”
Source: CSAT, 1999 TIP #33	

Table 2. Strategies for Quitting Methamphetamines

Some problems to expect:	What to do about it:
1. Mood is low; depression	Exercise Cultural/spiritual activities Fun (clean and sober friendships) Good nutrition Counseling Mental health assessment
<i>In case of suicidal thoughts/acts</i>	<i>911; emergency room if needed</i>
2. Memory loss	Reminders of appointments Visual (signs) or picture reminders
3. Poor judgment	Addictions treatment AA/NA
4. Drug craving	Learn how to avoid triggers Stay away from drug-using friends Schedule clean and sober activities

medication has been developed to block the effects of methamphetamine. However, there are options to consider at different stages of treatment. Antipsychotics are used for hallucinations in the immediate withdrawal period, and when they persist. Antidepressants have not been shown to be useful during withdrawal, but may reduce prolonged depression.³

Insomnia, anxiety, and pruritis can be treated symptomatically. Drug diversion is an issue that limits the choices and amounts of outpatient pharmacotherapy. Immunizations should be offered/updated.

Arrange: Arrange Follow-up

A follow-up visit was arranged for one week later.

Patient: “I had an assessment over at (Substance Abuse Program.) I was supposed to start IOP (Intensive

Outpatient Treatment) but I kept missing my appointments....”

Provider: “It sounds like you’ve already taken some steps in the right direction. So, when was your last use . . . ?”

Follow up visits address patient success, identify problems encountered, and offer encouragement. Consider referral to more intensive treatment if needed. The primary care provider can offer continuity of care. Many methamphetamine users have unrecognized health problems. Depression, STDs, Hepatitis C, and HIV should be considered.

Drug addiction is a chronic condition, with similar relapse rates as type 2 diabetes, hypertension, and asthma.⁷ As in tobacco cessation, knowledge of what to expect and methods to address the relapse factors aid the former user in recovery.

For patients unwilling to quit, and for those who relapse, motivational interventions are indicated. The provider should remain empathic, avoid arguing, and promote patient autonomy and self-efficacy.⁴

This description of the use of clinical intervention strategies with the methamphetamine user in a primary care setting should not minimize the difficulties involved in addressing this destructive addiction. The objective is for primary care providers to realize that we can apply the tools we already use for other risk behaviors, knowing that there is evidence for the benefit of brief interventions. One primary care medicine author stated, “Until we know more about which brief interventions are effective in patients with drug use problems, extrapolating findings from studies of brief interventions with problem drinkers is reasonable.”⁸

References

1. Stephens MB. Preventive Health Counseling for Adolescents. *American Family Physician*. 2006;74(7):1151-1159.
2. Dennis M, Scott CD, Funk R. An experimental evaluation of recovery management checkups (RMC) for people with chronic substance abuse disorders. *Evaluation and Program Planning* 2003;26(3):339-352.
3. California Society of Addiction Medicine (CSAM). *Recommendations for Improving California’s Response to Methamphetamine*. Available at: www.csam-asam.org. 2006. Accessed 11/3/06.
4. Fiore MC. Treating tobacco use and dependence. *AHCPR Supported Clinical Practice Guidelines*. Rockville MD: US Dept HHS, PHS, 2000. Available at: <http://www.ncbi.nlm.nih.gov/books/bv.fcgi?rid=hstat2.section.7741>. Accessed 11/6/06.
5. Rawson R. ed. 1999. *Treatment of Stimulant Use Disorders*. Rockville MD: US Dept HHS, PHS, SAMHSA. CSAT. Treatment Improvement Protocol #33.
6. American Academy of Family Physicians. *Drug Abuse: how to break the habit*. Available at: <http://familydoctor.org/357.xml>. November 2005. Accessed 11/9/06.
7. McClellan T, et al. Drug dependence, a chronic medical illness: implications for treatment, insurance, and outcomes evaluation. *JAMA*. 2000;284:1689-1695.
8. Vinson DC. Editorial: Screening and Intervening for Patients with Substance use Disorders. *American Family Physician*. April 2003. Available at: www.aafp.org/afp/20030401/editorials.html. Accessed 11/9/06.



Online Methamphetamine Resources

Lori de Ravello, MPH, CDC Assignee to the IHS National STD Program, Albuquerque, New Mexico

Prevention Programs

Project Venture; National Indian Youth Leadership Project: http://niyjp.org/programs/project_venture

Environmental Health Issues

Drug Lab Clean Up Program: <http://www.oregon.gov/DHS/ph/druglab/index.shtml>

MN DOH Meth Lab Clean Up: <http://www.health.state.mn.us/divs/eh/meth/lab/labcleanup.html>

OK Guidelines for Cleaning Up Former Meth Labs: <http://www.deq.state.ok.us/LPDnew/MethLabs/meth.htm>

WA State Dept of Health Clandestine Drug Lab Program: <http://www.doh.wa.gov/ehp/ts/CDL/default.htm>

WY DOH Cleaning Up Hazardous Chemicals at Meth Labs: <http://wdh.state.wy.us/epiid/methcleanup.asp>

Campaigns

End Meth Campaign — Partnership for a Drug Free America: <http://www.drugfree.org/endmeth>

Just Think Twice/Got Meth Campaign — Drug Enforcement Agency: <http://www.justthinktwice.com/gotmeth/home.html>

Meth Awareness Project: <http://www.methawarenessproject.org/>

Mothers Against Methamphetamine: <http://www.mamasite.net/>

Local Meth Efforts

Drug Free Northwest Arkansas: <http://www.drugfreenwa.com/>

Methamphetamine Awareness Project — Community Partnership of the Ozarks, Springfield, MO: <http://www.nometh.org/>

Meth Awareness and Prevention Project of South Dakota: <http://www.mappsdsd.org/>

Meth Education and Drug Awareness — Wright Co., MN: <http://www.meada.org/>

Meth Free Montana: <http://www.methfreemt.org/>

Methamphetamine Strike Force — County of San Diego: <http://www.no2meth.org/>

Rural Meth Education Project — Minot State University, Minot, ND: http://www.minotstateu.edu/rcjc/project_rmep.shtml

Stop Drugs — California Attorney General's Office: <http://www.stopdrugs.org/>

Drug Courts

Defining Drug Courts — Bureau of Justice Assistance: <http://www.ojp.usdoj.gov/BJA/grant/DrugCourts/DefiningDC.pdf>

Tribal Wellness to Healing Courts – Bureau of Justice Assistance: <http://www.ncjrs.gov/pdffiles1/bja/188154.pdf>

Substance Abuse Treatment

Counselor's Treatment Manual: Matrix Intensive Outpatient Treatment for People with Stimulant Use Disorders – SAMHSA: <https://store.health.org/catalog/productDetails.aspx?ProductID=17441>

The Matrix Institute on Addictions: <http://www.matrixinstitute.org/index.html>

The Methamphetamine Treatment Project - UCLA - Integrated Substance Abuse Programs (ISAP) and The Matrix Institute: <http://www.methamphetamine.org/>

Treatment for Stimulant Use Disorders – SAMHSA: <http://ncadi.samhsa.gov/govpubs/bkd289/>

Other Resources

Community Crisis: Public Health's Role in the Methamphetamine Epidemic “Partnering to Confront the Issue” (archived webcast): <http://www.mchcom.com/archivedWebcastDetailNewInterface.asp?aeid=400>

Meth on the Rez – NDN News: <http://www.ndnnews.com/Meth%20on%20the%20Rez.htm>

Methamphetamine Tools and Resources – Tribal Resource Center: <http://www.tribalresourcecenter.org/legal/details.asp?83>

National Congress of American Indians Meth Initiative: http://www.ncai.org/Meth_in_Indian_Country_Initiat.192.0.html



Healing to Wellness Courts: Best Practice for Indian Country

Honorable Amy Lovell, Pueblo of Zia Chief Judge; President and CEO, Lovell, Perales and Associates, Inc., Albuquerque, New Mexico


Healing to Wellness Courts evolved out of the Drug Court Movement from the 1980s, which was a response to the judicial and treatment systems' failure to effectively address substance abuse and its related activity. It was realized that the only way to combat the overwhelming issues of drug abuse, dependency, and addiction would be to create a marriage between the two systems intertwined with support from ancillary entities. States call their efforts Drug Courts, whereas tribes prefer Healing to Wellness Courts. Just as the name implies, a Healing to Wellness Court focuses on wellness through healing: healing of individuals, healing of families, healing of communities.

By combining indigenous forms of knowledge with modern techniques and evidence-based practices, Healing to Wellness Court teams have successfully paved pathways to recovery for substance abusing offenders and their families while also improving service delivery and achieving maximum use of limited resources. Criminal justice systems have not been effective in their punitive approaches, and most treatment systems have not been successful in keeping clients engaged long enough to effectively treat what ails them. Healing to Wellness Courts require a seamless integration between judicial and treatment systems, providing the necessary legal pressure on offenders to keep them in treatment for an adequate period of time. Results have been rather successful in holding substance abusing offenders accountable, providing them with a support network that encourages recovery through appropriate incentives and sanctions, and giving them the tools to recognize their own strengths, including resiliency to make better decisions providing for rehabilitation and reducing recidivism. In short, a Healing to Wellness Court that is effectively planned and implemented mandates culturally competent policies, procedures, and processes that make the most of traditional, holistic approaches to promote restorative justice practices resulting in healthier members, stronger families, and wealthier communities.

As tribes exercise their sovereignty, they increasingly utilize their own respective traditions, culture, and customs to do so. While their efforts to heal their communities may vary accordingly, they are becoming successful beyond measure. Similarly, no two Healing to Wellness Courts look alike, nor should they be expected to. Each community is afforded the

opportunity to empower their service providers and those served through traditional means such as talking circles, peace keeping courts, and other holistic approaches that nurture and heal the spirit, body, and mind, with the focus of each court varying as well. Originally, target populations for Healing to Wellness Courts were adult offenders, juvenile offenders, or family dependency and permanency planning. Today, they have evolved as special problem solving courts for identified challenges like methamphetamine courts, DUI courts, domestic violence courts, and grade courts to name a few. This concept works well for indigenous communities, regardless of the type of court system — tribal, contemporary, or combination — because it provides for integrated treatment systems of care that address substance abuse, mental health, medical screenings and treatment, employment, housing, academics, and other family issues such as child care and transportation. Most of our clients present with a multitude of problems that only a team approach can address with the participant and his/her family or significant others without overwhelming them. Team members vary according to tribal infrastructure and resources.

It is important that identification of all stakeholders, community strengths, assets, and resources occur. Inevitably, as tribal teams begin the planning stages, one of the first challenges noted is lack of funding. This is nothing new. Historically, tribal communities have been under-funded in every social discipline, including behavioral health, law enforcement, and judicial services among others.¹ By uniting all stakeholders, a shared vision and mission can be created and a plan of how each stakeholder can contribute and what role he/she will have in moving towards attainment of the goals and objectives defined by the vision and mission statements can be



executed. In doing so, existing funding streams are examined, restructured, and re-deployed, ensuring that those resources are maximized and gaps in service minimized. Planning and implementing in this manner provides a greater opportunity for sustainability of the Healing to Wellness Court program.

Historically, each system within tribal communities has operated autonomously. In doing so, we have often delegated the responsibility of healing to any one person or agency, and more often than not, *outside* of our communities. This activity has proven, time and time again, to be one of the most costly and detrimental acts that we have done as indigenous people, and social ills have continuously escalated over the past three decades and claimed so many lives of our relatives. Yet, we continue to do things in the same manner over and over, expecting different results. Healing to Wellness Courts mandate a team approach that requires training across disciplines. This breaks down barriers, such as lack of communication, territorialism, and professional jealousy, and promotes nothing short of professional respect and courtesy.

Healing is a journey, and participants are made aware that the team is not going to “heal” them, but rather assist them to become directors of their own healing. Because the journey is long, it is traveled in phases. Phase one usually focuses on detoxing, while phases two through four are stabilization, maintenance, and relapse prevention, respectively. Each phase brings about more responsibility for healing on the participant and his/her family, with decreasing need for judicial supervision. Treatment embodies the following principles, as identified by the National Institute of Drug Addiction:

1. No single treatment is appropriate for all.
2. Treatment needs to be readily available.
3. Effective treatment attends to the multiple needs of the individual.
4. Treatment plans must be assessed and modified continually to meet changing needs.
5. Remaining in treatment for an adequate period of time is critical for treatment effectiveness.
6. Counseling and other behavioral therapies are critical components of effective treatment.
7. Medications are an important element of treatment for many patients.
8. Co-existing disorders should be treated in an integrated way.
9. Medical detox is only the first stage of treatment.
10. Treatment does not need to be voluntary to be effective.
11. Possible drug use during treatment must be monitored continuously.
12. Treatment programs should assess for STDs, HIV/AIDS, hepatitis B and C, tuberculosis, and other infectious diseases, and help clients modify at-risk behaviors.
13. Recovery can be a long-term process and frequently requires multiple episodes of treatment.

Teams are required to continuously monitor, update, and implement individualized treatment plans that are relevant to clients and their needs. Completion of Healing to Wellness Court programs creates productive and healthy individuals who demonstrate good citizenry. And, since the wealth of a community should be calculated by the health of its members, the concept of healing needs to continue to be in the forefront of our financial portfolios.

As Indian people, we proudly boast that we have preserved our culture, our tradition, and our customs, and that we have survived acts of genocide aimed to desecrate the Indian race. Healing to Wellness Courts provide a road that is paved with indigenous forms of knowledge, guided by compassionate people, to be traversed by our relatives who are seeking a better way of life: a road that detours each of us from suffering silently with a feeling of despair and helplessness knowing that higher rates of alcoholism, addiction, and suicide plague our people and a road that has a light of hope shining brightly at the end of the tunnel. It suffices to say that Healing to Wellness Courts have evolved to become a mainstay for many tribal justice systems and a best practice for Indian Country.

Reference

1. US Commission on Civil Rights. July 2003. A Quiet Crisis: Federal Funding and Unmet Needs in Indian Country. Available at <http://www.usccr.gov/pubs/na0703/na0731.pdf>. Accessed 11/27/2006.



Healing from Methamphetamine: Community Efforts, Recovery, and Healing in American Indian and Alaska Native Communities

Don Coyhis, President and Founder, White Bison, Inc., and Richard Simonelli, Communication and Media Specialist, White Bison, Inc., Colorado Springs, Colorado

White Bison, Inc. convened a conference in Denver, Colorado in April 2006 to gather people working to stop the destructive effects of methamphetamine in American Indian and Alaska Native communities nationwide.¹ Participants were those in substance abuse/misuse recovery leadership positions, as well as grassroots community members from across the country. The Wellbriety Council of Elders was also present to share their wisdom and the unique perspectives that elders bring to any gathering. Entitled *Taking a Stand Against Meth: Recovery is Possible*, the goal of the conference was to provide hope and solutions for dealing with methamphetamine in Indian Country.

The Wellbriety Movement understands that prevention, treatment, intervention, and recovery are four interconnected aspects of substance abuse healing efforts in Indian Country. It's not possible or advisable to separate them, as might be done in some non-Native substance abuse recovery efforts. They are unified by the healing journey that arises when all four are acting in concert. Consequently, keynote addresses, workshops, and the grassroots Discovery Circles² that took place in Denver often connected these four directions of addictions recovery. The conference teaching and learning experience focused itself naturally into four primary areas: 1) knowledge about the drug methamphetamine; 2) what's working in Native American communities; 3) personal healing and recovery success stories; and 4) vision.

Knowledge about Methamphetamine

The methamphetamine epidemic is at the phase where education about its effects is vital. To drive this point home, Sissy Falcon said in a workshop entitled *Rez Watch*, "We need to become educated, educated, educated, educated, educated. Knowledge gained by feeding your spirit, knowledge about meth, changes the attitude in our hearts. We need this kind of learning for community change. We need our people educated and then we are going to see behavior change in our community." Wellbriety elder Theda Newbreast said in a workshop entitled *Women and Meth*, "We need to know meth. Know your language, know your culture, know your ceremony,

and know the facts about how the drug methamphetamine works." Then, to emphasize the need for education, she quipped, "We need to have a tee shirt that says, 'Just say 'Know' to Meth!'"

Presenters emphasized that the effects of methamphetamine are different from those of alcohol and other drugs that most recovery programs are familiar with.^{1,3} Because methamphetamine addicts a person so quickly, attacking centers in the brain, many of the cognitive-based treatment approaches that work for alcohol need to be modified for methamphetamine addicts. Group meetings based on reasoning, intuition, and the thought process, as well as methods such as reading, watching films, and other logical approaches probably won't work well for the methamphetamine sufferer. The 28-day treatment approach for alcoholics needs to be increased in duration and changed in curriculum for methamphetamine addicts.

What's Working?

Rez Watch is one of the most hopeful new approaches to deal with methamphetamine in reservation communities. Modeled after Meth Watch, which originated with non-Native communities in Kansas, Rez Watch empowers residents to take their communities back from methamphetamine dealers, users, and the houses where the drug is manufactured. It's called Rez Watch because reservations have their own cultural and logistical needs that might not be satisfied by the mainstream Neighborhood Watch or Meth Watch models. Rez Watch utilizes the anonymous reporting of possible meth-related activity through the *neighborhood activity log*, which can be collected at stores and tribal buildings on the reservation. "We need to stand for our people and we need to take back our reservations," said Sissy Falcon in the Rez Watch workshop. "They do not belong to the drug dealers."

What's also working is the integration of culture-specific healing programs into the non-Native approaches that some treatment centers are utilizing across the nation. For example, the Kokopelli Treatment Center, in Silver City, New Mexico, has had great success with the Matrix Model, created by the Matrix Institute of Los Angeles. The Matrix Model emphasizes personal structure and scheduling as well as a visual and repetitive approach for methamphetamine users. A primary component of the Matrix program at the Kokopelli

Center is recovery information tailored to those recovering from methamphetamine and addressing the fact that their cognitive ability is often impacted in early recovery.

The original Matrix program lacks a Native cultural approach, which can be added by using some of the White Bison Wellbriety movement resources. “The White Bison Firestarters Wellbriety Circle can be integrated into a Matrix approach to introduce Native culture and spirituality,” says Sue Lyons of the Kokopelli treatment Center. “The Matrix model has recognized that they were missing a Native American component,” she says. “Because of the damage to the left brain and analytical thinking in meth users, what we are finding is that there is tremendous success if we access the right brain with storytelling, drumming, prayer and ceremonies.”

Personal Recovery Stories

David Parnell stunned the conference with his own personal methamphetamine abuse recovery story. Not a word was heard as he presented slides and comments covering his own descent into meth, a suicide attempt, and then — through the Creator’s pity — his subsequent recovery. His presentation, which he makes available to communities everywhere, is exactly the prevention education package that some might need to realize what a bad choice methamphetamine use is.³ Wellbriety elder Ozzie Williamson said during a follow-up elders panel, “The session we’ve just heard is probably one of the best I have ever seen in the thirty-some years that I’ve spent in this field. The message I would have is we need to get more people like David Parnell into our communities to get our young people to understand that this can happen and will happen to them if they try using or continue to use methamphetamine or other drugs. I think it was really a blessing to have him here with us this morning.”

Four former methamphetamine users now associated with the Native American Rehabilitation Association (NARA), based in Portland, Oregon, also shared their recovery stories from methamphetamine, demonstrating the fact that people can and do recover. Each emphasized that re-embracing their own cultural traditions and ways was and is an essential part of healing from methamphetamine. One said, “I went to NARA for 3 months. There I found my self-identity, who I was, and the opportunity to teach the culture and tradition that was taught to me.” Another revealed, “I went to NARA and met the NARA family. It really helped me in my recovery. Now we’re building a traditional canoe with others in recovery. That’s part of the healing process. We got together with social services and the drug and alcohol department we have in Nisqually to back our traditional canoes in our area. I brought back the White Bison talking circle way to Nisqually, as well as other things I learned at NARA. Now White Bison is on Nisqually and it’s taking off over there like a wildfire.”

Vision

Many visionary teachings were shared during the conference in Denver. Don Coyhis, Founder and President of White Bison, Inc., encouraged participants to think in terms of healing from methamphetamine, not waging war, as is often promoted by some mainstream efforts.⁴ Healing, not war, is more in line with the Indian way of thinking.

Dr. Eduardo Duran, a psychologist, author, and drug and alcohol therapist with the Miwok/Maidu community in northern California, revealed that he sometimes asks his clients in recovery what they think the spirit of methamphetamine is trying to tell them. He asks them why they think it has come to them and what its unique message might be for them. This question often awakens the visionary response, to which many clients will respond positively.

Don McCloud, a grassroots participant from the Northwest, shared a vision that emphasizes a return to tradition and culture in healing from alcohol and drug problems. “We’re the survivors,” he says, “It’s about our culture and our spirituality. If we don’t find our way we’re just going to be like the dominant society. If we don’t find our spirituality we will become the dominant society.” The Wellbriety Movement teaches that the community is the treatment center. It is also the best place for prevention, intervention, recovery, and healing to begin and to flourish.

A document summarizing results from the five grassroots Discovery Circles that took place at the conference is available on the White Bison, Inc. website.² White Bison, Inc. is making available the DVDs and audio CDs of many of the keynote addresses, workshop sessions, and grassroots Discovery Circles from the April, 2006 conference, at cost, to readers of this IHS journal article. Contact White Bison, Inc. at (877) 871-1495, or e-mail info@whitebison.org to receive a set.

References

1. White Bison, Inc. (5/12/2006). Taking a Stand Against Meth: Recovery is Possible. *Wellbriety! Online Magazine*. 7(3).
2. White Bison, Inc. (6/5/2006). Discovery Circles: Mind Mapping Community Solutions. *Wellbriety! Online Magazine*. 7(5).
3. Coyhis D, Simonelli R. Meth in Indian Country. *Winds of Change*. Summer 2006;21(3):36-40.
4. Parnell D. *Drug Education and Methamphetamine Seminars*. (Contact info: phone (731) 469-9105, e-mail: david_parnell@bellsouth.net; website: www.facingthedragon.org).
5. Coyhis D. (10/1/2006). Declare Healing, Not War. *Wellbriety! Online Magazine*. 7(10).

Editor's Note: The following is a digest of the monthly Obstetrics and Gynecology Clinical Consultant's Newsletter (Volume 4, No. 12, December 2006) available on the Internet at <http://www.ihs.gov/MedicalPrograms/MCH/M/OBGYN01.cfm>. We wanted to make our readers aware of this resource, and encourage those who are interested to use it on a regular basis. You may also subscribe to a listserv to receive reminders about this service. If you have any questions, please contact Dr. Neil Murphy, Chief Clinical Consultant in Obstetrics and Gynecology, at nmurphy@scf.cc.

OB/GYN Chief Clinical Consultant's Corner Digest

Abstract of the Month

Organizational silence threatens patient safety

Organizational silence refers to the tendency for people to do or say very little when confronted with significant problems or issues in their organization or industry. This paper focuses on some of the less obvious factors contributing to organizational silence that can serve as threats to patient safety. Converging areas of research from the cognitive, social, and organizational sciences and the study of socio-technical systems help to identify some of the underlying factors that serve to shape and sustain organizational silence. These factors have been organized under three levels of analysis:

1. individual factors, including the availability heuristic, self-serving bias, and the status quo trap;
2. social factors, including conformity, diffusion of responsibility, and microclimates of distrust;
3. organizational factors, including unchallenged beliefs, the good provider fallacy, and neglect of the interdependencies.

Finally, a new role for health care leaders and managers is envisioned. It is one that places high value on understanding system complexity and does not take comfort in organizational silence.

Henriksen K, Dayton E. Organizational silence and hidden threats to patient safety. *Health Serv Res.* 2006 Aug;41(4 Pt 2):1539-54. Reprints (AHRQ Publication No. 06-R060) are available from the AHRQ Publications Clearinghouse, <http://www.ahrq.gov/research/order.htm#clear>

OB/GYN CCC Editorial comment:

Value dissent and multiple perspectives as signs of organizational health

Henriksen and Dayton, of the Agency for Healthcare Research and Quality (AHRQ), describe the individual, social, and organizational factors that contribute to organizational silence and can threaten patient safety. They cite several individual factors that contribute to clinician silence. For example, the availability heuristic suggests that if relatively infrequent events that harm patients go unreported and are not openly discussed, clinicians don't believe these events are a problem at their hospital. A second factor is self-serving bias. People tend to view themselves as "above average" in their chosen field of work and so "why do things differently?" Successes are attributable to their

own abilities but failures are blamed on situational factors. Finally, members of all organizations display a strong tendency to perpetuate the status quo and not speak up or rock the boat.

Several social factors also underlie clinician silence. There is great pressure to conform in order to gain acceptance and work harmoniously with coworkers. Diffusion of responsibility is also a problem. In clinical settings, individual roles and responsibilities are often assumed rather than clearly spelled out. Under these conditions of diffused responsibility, components of care that should be attended to are often missed. Also, managers who seek blame and attribute error to the individual failings of careless or incompetent staff create a microclimate of distrust. Finally, three areas of organizational vulnerability that warrant closer attention are unchallenged beliefs, the perceived qualities of the good worker who "works around" problems rather than focusing on the contributory factors to the problem, and lack of understanding of the interdependence of complex clinical systems.

The authors recommend that health care leaders and managers value dissent and multiple perspectives as signs of organizational health, and question agreement, consensus, and unity when they are too readily achieved.

Another successful example is the 100,000 Lives Campaign, which is an initiative to engage US hospitals in a commitment to implement changes in care proven to improve patient care and prevent avoidable deaths. The Institute for Healthcare Improvement estimates that the lives saved as of June 14, 2006 was 122,300.

To that end, the National Indian Health MCH and Women's Health meeting, August 15 - 17, 2007 in Albuquerque will highlight speakers from the Institute for Healthcare Improvement and others who have evaluated and treated various health care systems. The meeting has individual facility program review as well as many hours of CME/CEUs. Your facility should send a team of staff to this meeting; you and 2 - 3 colleagues from different disciplines should start planning now. For more information, go to <http://www.ihs.gov/MedicalPrograms/MCH/F/CN01.cfm#Aug07>.

From Your Colleagues

Judy Thierry, HQE

Menstruation in girls and adolescents: using the menstrual cycle as a vital sign

Evaluation of the menstrual cycle is a viable tool to assess

healthy development of teen girls' menstrual patterns. Young patients and their parents often are unsure about what represents normal menstrual patterns, and clinicians also may be unsure about normal ranges for menstrual cycle length and amount and duration of flow through adolescence. "It is important to be able to educate young patients and their parents regarding what to expect of a first period and about the range for normal cycle length of subsequent menses," the authors point out. "It is equally important for clinicians to have an understanding of bleeding patterns in girls and adolescents, the ability to differentiate between normal and abnormal menstruation, and the skill to know how to evaluate young patients' conditions appropriately."

American Academy of Pediatrics Committee on Adolescence; American College of Obstetricians and Gynecologists Committee on Adolescent Health Care; Diaz A, Laufer MR, Breech LL. Menstruation in girls and adolescents: using the menstrual cycle as a vital sign *Pediatrics*. 2006 Nov;118(5):2245-50.

Obstetrics

Shoulder dystocia: Only 43% participants could achieve delivery before training

Conclusion: This study verifies the need for shoulder dystocia training; before training only 43% participants could achieve delivery. All training with mannequins improved the management of simulated shoulder dystocia. Training on a high-fidelity mannequin, including force perception teaching, offered additional training benefits. LEVEL OF EVIDENCE: I

Crofts JF, et al Training for Shoulder Dystocia: A Trial of Simulation Using Low-Fidelity and High-Fidelity Mannequins. *Obstet Gynecol*. 2006 Dec;108(6):1477-1485.

OB/GYN CCC Editorial Comment

Regular simulation practice: win x3

The Crofts JF, et al randomized trial confirms previous recommendations that regular drills should be part of ongoing health care expectations. Regular drills and medical emergency preparedness improve patient care, as well as satisfy JCAHO evaluations. In addition, this process is endorsed in this month's ACOG Committee Opinion (see below).

The National Indian Health MCH and Women's Health meeting, August 15 - 17, 2007 in Albuquerque will highlight speakers from the Institute for Healthcare Improvement and others who have developed simulation processes. Phoenix Indian Medical Center offers a best practice example of a successful model in Indian Country. Contact Judy Whitecrane for more information Judy.Whitecrane@ihs.gov. The Advanced Life Support in Obstetrics Course is another great resource. Go to http://www.aafp.org/online/en/home/cme/aafpcourses/clinical_courses/also.html

Medical emergency preparedness, ACOG Committee

Abstract: Patient care emergencies may occur at any time in a hospital or an outpatient setting. To respond to these emergencies, it is important that obstetrician/gynecologists

prepare themselves by assessing potential emergencies that might occur, creating plans that include establishing early warning systems, designating specialized first responders, conducting emergency drills, and debriefing staff after actual events to identify what went well and what are opportunities for improvement. Having such systems in place may reduce or prevent the severity of medical emergencies.

Medical emergency preparedness. ACOG Committee Opinion No. 353. American College of Obstetricians and Gynecologists. *Obstet Gynecol*. 2006;108:1597-99.

Majority of women with GDM are not tested for glucose intolerance after delivery

Conclusion: Although persistent abnormal glucose tolerance was common in our cohort, less than half of the women were tested for it. Our data suggest that to increase rates of postpartum glucose testing, improved attendance at the postpartum visit with greater attention to testing and better continuity between antenatal and postpartum care are required. LEVEL OF EVIDENCE: II-2.

Russell MA, et al. Rates of postpartum glucose testing after gestational diabetes mellitus. *Obstet Gynecol*. 2006 Dec;108(6):1456-62.

OB/GYN CCC Editorial Comment

GDM women are not tested for glucose intolerance after delivery

Although Russell et al are not from Indian Country, the same result would be found here, despite our enormous burden of diabetes related disease. Our Indian health GDM guidelines recommend a 2-hour oral glucose tolerance test at six weeks post partum and a fasting blood glucose every three years thereafter. How many of your GDM patients actually get that follow-up? Go to http://www.ihs.gov/MedicalPrograms/MCH/w/Documents/DM_Preg102504_000.doc

Gynecology

HPV vaccine is effective: Why do we not provide it to most AI/AN?

Conclusion: Based on the data obtained in this study, widely-implemented prophylactic HPV vaccination could make an important contribution to the reduction of the risk for cervical cancer and could also prevent about half the vulvar carcinomas in younger women and about two thirds of the intraepithelial lesions in the lower genital tract. LEVEL OF EVIDENCE: II-3.

Hampl M, et al. Effect of human papillomavirus vaccines on vulvar, vaginal, and anal intraepithelial lesions and vulvar cancer. *Obstet Gynecol*. 2006 Dec;108(6):1361-1368.

OB/GYN CCC Editorial comment:

Honestly, are you actively giving out HPV to all your female patients between 9 - 26 years old?

I bet you are not, but you should be. Hampl M, et al is additional evidence that the HPV vaccine is clearly beneficial.

Yet HPV vaccine is not widely available to most AI/AN women 9 - 26 years old. How can this disparity be resolved? On November 1, 2006, the Centers for Disease Control and Prevention added the new human papillomavirus (HPV) vaccine to the federal Vaccines for Children program, which provides free vaccines to children from families with low incomes or who are uninsured. The articles in the Winter 2006 issue of the Guttmacher Policy Review discuss related policy issues. In "Achieving Universal Vaccination Against Cervical Cancer in The United States," the authors discuss the case for universal vaccination, the role of school-entry requirements, financing challenges, the potential role of family planning clinics in an HPV vaccine campaign, and solutions to various challenges presented by the HPV vaccine.

Child Health

Why do disparities in infant mortality continue to persist between AI/AN and white infants?

Objectives: To describe changes in infant mortality rates, including birthweight-specific rates and rates by age at death and cause.

Methods: We analyzed US linked birth/infant-death data for 1989 - 1991 and 1998 - 2000 for American Indians/Alaska Native (AI/AN) and White singleton infants at > or = 20 weeks' gestation born to US residents. We calculated birthweight-specific infant mortality rates (deaths in each birthweight category per 1000 live births in that category), and overall and cause-specific infant mortality rates (deaths per 100,000 live births) in infancy (0 - 364 days) and in the neonatal (0 - 27 days) and postneonatal (28 - 364 days) periods.

Results: Birthweight-specific infant mortality rates declined among AI/AN and White infants across all birthweight categories, but AI/AN infants generally had higher birthweight-specific infant mortality rates. Infant mortality rates declined for both groups, yet in 1998 - 2000, AI/AN infants were still 1.7 times more likely to die than White infants. Most of the disparity was because of elevated post-neonatal mortality, especially from sudden infant death syndrome, accidents, and pneumonia and influenza.

Conclusions: Although birthweight-specific infant mortality rates declined among both AI/AN and White infants, disparities persist. Preventable causes of infant mortality identified in this analysis should be targeted to reduce excess deaths among AIAN communities.

Tomashek KM, et al. Infant mortality trends and differences between American Indian/Alaska Native infants and White infants in the United States, 1989 - 1991 and 1998 - 2000. *American Journal of Public Health*. December 2006, Vol 96, No. 12.

Editorial Comment

Judy Thierry, MCH Coordinator, HQE

Kay Tomashek and her CDC colleagues present AI/AN infant mortality trends and differences emphasizing birthweight-specific infant mortality rates. This national picture uses three year aggregate data drawn from the NCHS

Vital Statistics 1989 - 1991 and again 1998 - 2000).¹ Given the contribution of SIDS to elevated post neonatal AI/AN infant mortality, it is important to consider several key studies and national interventions that occurred in the early to mid 1990s between these two data sets. Key contributions include New Zealand and other international studies on SIDS; further study of SIDS in Seattle King County by Spiers and Guntheroth on the supine sleep position²; roll out of the AAP and NICHD guidelines on infant sleep position and the "Back to Sleep" campaign; and the prospective case control study conducted by the IHS, NICHD, and CDC (1994 - 1997) entitled the "Aberdeen Area Tribal Chairman's Health Board (AATCHB) Infant Mortality study," "Mi Cinca kin towani ewaktonji kte sni," "I will never forget my child."

The APHA December 2006 publication of the analysis demonstrates downward trends in AI/AN birthweight-specific infant mortality rates (albeit not enough) and helps us to further clarify the populations at risk in this complex issue, one that will require further elaboration using the triple risk factor model.³ A list of key partners includes tribal epidemiology centers (local surveillance) linked with perinatal infant and child mortality review teams (local review), community health representative and public health nurse outreach (culturally-based outreach); including the Healthy Start Project model of home interventions and case management and other intensive maternal support programs. Tomashek discusses perinatal tobacco exposure. It is cited among the 33 references that are essential reading when discussing AI/AN birth and infant death data. Maternal and family risk factors will be further elucidated with an AI/AN specific point-in-time PRAMS soon to be underway. Funding of comprehensive campaigns to address tobacco exposure such as the AATCHB Smoke Free Homes Campaign should be priority. Timely access to care, quality of care, care of the maternal/fetal unit through a planned and regionalized and risk stratified manner remain fundamental to infant survival and maternal wellbeing.

References

1. Single years have too few deaths to report any significance. In some reports you will note 5 and 6 year aggregates to obtain sufficient numbers in the 'cells'. For this analysis 3 year aggregates were used.
2. Guntheroth, W. G., Spiers, P. S., Sleeping prone and the risk of sudden infant death syndrome, *JAMA*, Vol. 267 No. 17, May 6, 1992 cited December 4, 2006 http://jama.ama-assn.org/cgi/content/abstract/267/17/2359?ijkey=b3b5d2208410c6629f489a51c5f84ab6b9c0dbca&keytype=tf_ipsecsha
3. A triple-risk model for the pathogenesis of SIDS - intersection of three overlapping factors: (1) a vulnerable infant; (2) a critical developmental period in homeostatic control, and (3) an exogenous stressor(s). cited December 4 http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Abstract&list_uids=80

38282 Filiano JJ, Kinney HC. A perspective on neuropathologic findings in victims of the sudden infant death syndrome: the triple-risk model. *Biol Neonate*. 1994;65 (3-4):194-7.

Chronic disease and Illness

Firearm safety in homes with adolescents

Approximately one-third of US households with children and adolescents contain firearms. Despite recommendations to keep firearms stored unloaded and locked, a significant number of households store them loaded or unlocked, substantially increasing the risk that children or adolescents will accidentally or intentionally use a firearm to cause injury. Parents tend to assume that older children will act more responsibly, and studies have evaluated safe firearm storage according to the age of the children in the home. However, no studies have addressed individually the issues of storing firearms unloaded and of storing them in a locked place. Johnson and associates evaluated these individual safety issues in households with children or adolescents.

The authors conclude that parents of adolescents are less likely to store their firearms safely compared with parents of younger children. They add that these results are worrying because a significant number of firearm injuries occur in the adolescent age group. The authors suggest that firearm prevention programs focus on parents with adolescent children to improve safety practices.

Johnson RM, et al. Are household firearms stored less safely in homes with adolescents? Analysis of a national random sample of parents. *Arch Pediatr Adolesc Med*. August 2006;160:788-92.

Features

ACOG

Treatment with selective serotonin reuptake inhibitors during pregnancy

Abstract: Depression is a common condition among women of reproductive age, and selective serotonin reuptake inhibitors (SSRIs) are frequently used for the treatment of depression. However, recent reports regarding SSRI use during pregnancy have raised concerns about fetal cardiac defects, newborn persistent pulmonary hypertension, and other negative effects. The potential risks associated with SSRI use throughout pregnancy must be considered in the context of the risk of relapse of depression if maintenance treatment is discontinued. The American College of Obstetricians and Gynecologists' Committee on Obstetric Practice recommends that treatment with all SSRIs or selective norepinephrine reuptake inhibitors or both during pregnancy be individualized and paroxetine use among pregnant women or women planning to become pregnant be avoided, if possible.

Treatment with selective serotonin reuptake inhibitors during pregnancy. ACOG Committee Opinion No. 354. American College of Obstetricians and Gynecologists. *Obstet Gynecol*. 2006;108:1601-3. *Obstet Gynecol*. 2006;108:1597-99.

Breastfeeding

Suzan Murphy, PIMC

Flu season and breastfeeding, CDR Julie Warren, RPh, Pharmacist, PIMC, Chair, IHS MCH Breastfeeding Web Page Medication Section

When a breastfeeding mom gets an upper respiratory infection, there are many medications that can help and are safe to use. General guidelines are as follows:

- ✓ Keep breastfeeding. The baby has already been exposed. A breastfeeding mom's immune system will make antibodies that fight the infection, protecting both the mom and her baby.
- ✓ Take the medicine right after nursing or before baby's longest sleep time.
- ✓ Watch baby for effects from the medicines that you take.
- ✓ Don't choose medicines that have a variety of ingredients.
- ✓ Use "regular strength" instead of "extra strength," "maximum strength," or "long acting."
- ✓ Follow the directions on the label. Don't take more than what is recommended.
- ✓ Take the lowest dose recommended.

If mom has:

A fever, headache, or aches all over, try:

- Acetaminophen (Tylenol[®] and many other brands)
- Ibuprofen (Advil[®], Motrin[®], etc.)
- Naproxen (Aleve[®], etc.)
- Do not use aspirin.

A stuffy nose, use:

- Best: sodium chloride nasal spray
- Phenylephrine nasal spray (Neo-Synephrine[®], etc.)
- Oxymetazoline nasal spray (Afrin[®] and others)
- Pseudoephedrine oral tablets (Sudafed[®] and many other brands). Moms may notice a decrease in breast milk production if they take Sudafed for extended periods.

Sneezing, hay fever symptoms, and her allergies are acting up, consider:

- Diphenhydramine (Benadryl[®] and many other brands)
- Brompheniramine + pseudoephedrine (Bromfed[®], Rondec[®] syrup, etc.)
- Triprolidine + pseudoephedrine (Actifed[®] and other brands)
- Chlorpheniramine (Coricidin[®] and many other brands)
- Dexbrompheniramine + pseudoephedrine (Drixoral[®] and others)
- Loratadine (Claritin[®], Alavert[®], others)
- Cromolyn sodium nasal spray (Nasalacrom[®]).

A sore throat, even after a cup of hot tea, use:

- Warm to hot salt water gargles (don't swallow it!)
- Throat sprays (Cepacol[®] Maximum Strength Sore Throat Spray, others)
- Throat lozenges (Sucrets[®] Regular Strength, Halls[®] Mentho-Lyptus Drops)

- Don't use phenol and hexylresorcinol.
- A cough, try:
- Guaifenesin with or without Dextromethorphan (Robitussin[®], Robitussin DM and other brands with the same ingredients).

For more information, go to <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?LACT>. The link is also listed on the IHS MCH Breastfeeding web page, in the Medication section. Go to <http://www.ihs.gov/MedicalPrograms/MCH/M/bfMeds.cfm>.

International Health Update Claire Wendland, Madison, WI A nurse, a doctor, and an epidemiologist were standing by the river . . .

Most of us have heard this anecdote: a nurse, a doctor, and an epidemiologist are standing at a river's edge when they notice body after body floating by. The doctor and nurse jump in, fish out everyone they can, and begin resuscitating the victims. The epidemiologist runs upstream instead, hollering over her shoulder, "I'm going to see who's pushing them in!"

In recent years, scholars from public health and related fields have increasingly proposed upstream or "structural" interventions into serious problems of public health, as opposed to the traditional education or behavior change interventions. The word "structural" in this sense refers to the social, political, and economic structures that make individuals more vulnerable to disease and violence. The logic of such proposals is that, for instance, it makes little sense to combat diabetes by teaching individuals about healthy eating in a poor rural community if the only place to buy food for miles around is a gas station convenience store specializing in Cheetos, and subsidized corn syrup production means that soda is cheaper than clean water.

Although much epidemiologic and social science literature explores the structural determinants of poor health, few structural intervention trials have been conducted. In fact, controversy over whether such trials are worthwhile (or ethical) is substantial. A recent trial of microfinance initiatives and their effects on intimate partner violence (IPV) and HIV seroconversion rates provides us a rare opportunity to examine the effects of a structural intervention, although with mixed results.

Paul Pronyk and colleagues from the University of the Witwatersrand noted that poverty, lack of economic opportunities, and gender inequalities combine in rural South Africa to allow high levels of both HIV infection and IPV in women. Projects addressing violence and HIV through education alone have met with little success. Would improved economic opportunities for women do better? Pronyk's team randomized eight villages in Limpopo province to intervention – establishment of a microfinance program combined with a participatory empowerment curriculum – or a comparison group. Over 400 of the poorest women in intervention villages received one or more small loans averaging \$165 to support business initiatives; as a loan condition, they also attended training sessions on gender empowerment, relationships,

communication, HIV, and domestic violence. The researchers assessed the impact of the intervention not only on the women themselves, but also on young people living in loan recipients' households and on randomly selected villagers. In the intervention villages, reports of intimate partner violence declined dramatically (adjusted RR 0.45, 95% CI 0.23-0.91). Intervention villagers also reported improved household communication, especially on matters of sex and sexuality, and improvement in the total value of household assets – although not food security or other measures of wealth. Several other attitudinal measures of empowerment trended toward positive change, but none met criteria for statistical significance. In addition, young people in intervention villages showed no difference in HIV seroconversion and rates of unprotected sex with someone other than a spouse. (Loan recipients themselves were not asked these questions. At a mean age of 41, the authors imply they were considered too old to discuss such matters!)

Although the study did not demonstrate the effectiveness of microfinance for HIV prevention, it is the first to show that microfinance is effective in reducing intimate partner violence. (Research in South Asia demonstrated initial increases in IPV with the initiation of microfinance, perhaps related to threats to male control of household resources, followed by a later decline.) It also demonstrates that a relatively small structural intervention can have relatively quick effects at the community level.

Pronyk PM, Hargreaves JR, Kim JC et al. Effect of a structural intervention for the prevention of intimate-partner violence and HIV in rural South Africa: a cluster randomized trial. *Lancet*. 368:1973-83, 2006.

Medical Mystery Tour A boy has been born in Chile with a fetus in his stomach

Santiago, Chile (Nov. 24). A boy has been born with a fetus in his stomach in what doctors said was a rare case of "fetus in fetu" in which one twin becomes trapped inside another during pregnancy and continues to grow inside it. Doctors carried out a scan on the boy's mother shortly before she gave birth on November 15 in the southern city of Temuco and noticed the 4-inch-long fetus inside the boy's abdomen. It had limbs and a partially developed spinal cord but no head and stood no chance of survival, doctors said. After the birth, doctors operated and removed the fetus from the boy's stomach. The boy, who has not been named, was recovering at Temuco's Hernan Henriquez hospital. It's very rare," said Maria Angelica Belmar, head of the hospital's neonatal wing, speaking of fetus in fetu cases. It occurs in only one in every 500,000 live births," she told Reuters, adding that the number of cases recorded worldwide was fewer than 90. Before you explain the embryology of this case to us, please answer this one simple question: Which reputable medical resource was this story taken from? National Enquirer or Reuters?

Stay tuned till next month to find out.

Midwives Corner

Lisa Allee, CNM, Chinle

Start 'em young for future success and maybe no one will be left behind

This month I digress. I have no peer juried article; I don't even have a URL link. This month's topic is in honor of my mom. I visited her for Thanksgiving and we read an article in the Sunday *New York Times Magazine* called "Still Left Behind" about the No Child Left Behind Act. My mom worked for 20+ years as a reading specialist in a junior high school and has continued her work for literacy in retirement. She has always admonished me and my sisters to read to our children every day starting at birth, and for years we had the grandma-books-of-the-month delivery service. She was excited about this article because besides restating the previously known fact that school success is proportional to income, the author presented research delving into why this is. Some researchers have found a link between a child's school success and his/her vocabulary at age three — middle to upper income children often have 1000+ words at that age versus lower income children who have one-half to one-quarter that amount — and that the number of words a child has at age three is directly related to the mother's/parents' vocabulary. Another researcher found that not only did the parents' vocabulary matter, but the way the parents speak to the children plays a very important role. Parents of successful children (mid to upper income generally) used a higher proportion of encouragements, while parents of less successful children (generally lower income) used a much higher proportion of discouragements when speaking to their children. So, I was profoundly struck by the possibility that we as midwives, nurse practitioners, obstetricians, and pediatricians, could have an influence on parents and, thus, their children by pointing out three rules:

1. Follow Grandma Allee's rule of reading to your child every day starting at birth.
2. Improve your vocabulary and use your new words with your child.
3. Make sure at the end of every day you have said more encouraging things to your child than discouraging things.

I did this with a couple expecting their second child the other day. It took about a minute, and they said thank you for the information! It can fit into a busy clinic. If you want to read the article it is "Still Left Behind" in the November 26, 2006 Sunday *New York Times Magazine*.

Navajo News

Tomekia Strickland, Chinle

GYN spotlight: endometrial ablation

Pre-menopausal dysfunctional uterine bleeding unrelated to malignancy continues to be a significant problem for women wrought with social embarrassment, disruption of daily activities, and morbidity associated with anemia. Not only is it a challenging condition for the patient, but dysfunctional uterine bleeding usually requires lengthy and frequent outpatient visits

for appropriate evaluation and management. Many times, patients have suffered for years with the condition and often present discouraged after a series of failed hormonal regimens. Hysterectomy, the only procedure that is 100% effective in eliminating abnormal uterine bleeding, is often less acceptable to Native American women than other populations, both for cultural reasons and because of a general reluctance to undergo major surgery. Thus endometrial ablation has risen as an ideal treatment option for women who have completed child bearing, failed conservative management, and desire uterine conservation.

The Department of Gynecology at Chinle Service Unit is now offering endometrial ablation to appropriate candidates, as are some other IHS sites. There are several global endometrial ablation techniques that have become available nationally over the past few years. *Global endometrial ablation* refers to a series of FDA approved newer generation technologies that do not require an operative hysteroscope. These include Thermachoice (hot liquid filled balloon), hydrothermal ablation (circulating hot water), Novasure (bipolar desiccation), Her Option (cryoablation), and microwave ablation. This is in contrast to the *standard technique* which uses monopolar energy via a rollerball, roller barrel, or resectoscope requiring operative hysteroscopy. There is also increased risk of uterine perforation and fluid overload with the standard techniques. We have started using the Novasure system, which is a global ablative technique that utilizes a three dimensional bipolar gold mesh that when inserted conforms to the shape of the uterine cavity. The average ablation time for Novasure® is 90 seconds. It also has the advantage of not requiring hormonal pretreatment to thin the endometrial lining. When used correctly, the global ablative techniques are considered safe, effective, fast, simple to perform, painless, and cost effective to both physician and patient. Many of these procedures can also be performed as office-based procedures.

Like the standard technique, global ablation techniques are considered successful not so much according to amenorrhea rates, but by reduction in menstrual flow. Hypomenorrhea correlates with high rates of subjective patient satisfaction, usually greater than 80 - 90%. The amenorrhea rates for some of the devices are as follows: Thermachoice, 14% at 12 - 24 months; Microwave 38% at 3 years; and Novasure 51% at 1 year.

In conclusion, global endometrial ablation will most likely continue to become an increasingly popular and primary minimally invasive surgical treatment option for women who have completed childbearing and continue to suffer for abnormal uterine bleeding despite medical therapy. Like all medical and surgical interventions, care must be taken to evaluate each patient carefully and individualize their treatment plan accordingly. "Endometrial Ablation" by UpToDate, www.uptodate.com, provides a detailed discussion on the indications, contraindications, and safety profiles for each ablative procedure. If you would like more information about our exciting but still new experience with Novasure, please feel free to contact me at tomekia.strickland@ihs.gov.

Nurses Corner

Sandra Haldane, HQE

Nurses less satisfied than physicians or nurse managers: perceptions of teamwork on L/D

Caregiver role influences perceptions of teamwork. Overall, physicians and nurse managers were much more satisfied than nurses with the collaboration they experienced. For example, anesthesiologists had higher scale scores than certified registered nurse anesthetists for five of the six teamwork climate items. Most (80 %) L&D staff felt it was easy for personnel in their unit to ask questions. However, only 55 % found it easy to speak up if they perceived a problem with patient care, and only half felt that conflicts were appropriately resolved. The study was supported in part by the Agency for Healthcare Research and Quality (HS11544).

Oklahoma Perspective

Greggory Woitte, Hastings Indian Medical Center

Reduction in teen pregnancies

The preliminary numbers from 2005 from the CDC show a 2 % reduction in teenage pregnancies, down to its lowest recorded level in 65 years. The biggest decline was in the ages 15 - 17 years group. Here in Oklahoma, we were the 8th highest state in the nation for teen births ages 15 - 19 in 2002. Like in all other states, we as women's health providers have to work hard at encouraging young women to delay sexual activity as well as to take steps to prevent becoming pregnant. ACOG recently released a statement that a 13 month supply of OCPs showed a greater likelihood of continuation and use and would be very beneficial in the continued reduction of teen pregnancies. In fact, it is estimated that for every dollar invested in teen pregnancy prevention programs, at least \$2.65 was saved in direct medical and social service costs (The National Campaign to Prevent Teen Pregnancy. Not Just Another Single Issue: Teen Pregnancy's Link to Other Social Issues, 2002).

NCHS Health E Stats – Births-Preliminary Data for 2005
<http://www.cdc.gov/nchs/products/pubs/pubd/hestats/prelimbirths05/prelimbirths05.htm>

ACOG Statement 13 month Supply of OCPS leads to more consistent use http://www.acog.com/from_home/publications/press_releases/nr11-01-06-2.cfm

Perinatology Picks

George Gilson, MFM, ANMC

Amniocentesis procedure-related loss risk 1 in 1600, not prior 1 in 200

Results: The spontaneous fetal loss rate less than 24 weeks of gestation in the study group was 1.0% and was not statistically different from the background 0.94% rate seen in the control group (P=.74, 95% confidence interval -0.26%, 0.49%). The procedure-related loss rate after amniocentesis was 0.06% (1.0% minus the background rate of 0.94%). Women undergoing amniocentesis were 1.1 times more likely to have a spontaneous loss (95% confidence interval 0.7-1.5).

Conclusion: The procedure-related fetal loss rate after midtrimester amniocentesis performed on patients in a

contemporary prospective clinical trial was 0.06%. There was no significant difference in loss rates between those undergoing amniocentesis and those not undergoing amniocentesis. LEVEL OF EVIDENCE: II-2.

Eddleman KA et al. Pregnancy loss rates after midtrimester amniocentesis. *Obstet Gynecol.* 2006; 108(5):1067-72

STD Corner

Lori de Ravello, National IHS STD Program

Less than half of parents infected with HIV tell their children about the diagnosis

Parents are reluctant to disclose their HIV infection to their children, primarily because they fear the emotional impact. As a result, fewer than half (44 percent) of children are aware of their parent's HIV infection, according to a new study supported in part by the Agency for Healthcare Research and Quality (HS08578 and T32 HS00046). Parents did not disclose their HIV status to their children primarily due to worry about the emotional consequences of disclosure for the child (67 percent), worry that the child would tell other people (36 percent), and not knowing how to tell their child (28 percent). Many parents also feared that their children would reject them or lose respect for them. Certain parents were less likely to disclose their HIV infection than others. These included those who contracted HIV through heterosexual intercourse (rather than homosexual intercourse or intravenous drug use), those with higher CD4 cell counts (indicative of greater disease progression), those who were more socially isolated, and those with younger children. According to the parents, 11 percent of children who were aware of their parent's HIV infection worried they could catch HIV from their parent, 5 percent had experienced other children not wanting to play with them, and 9 percent had been teased or beaten up.

Corona R, et al. Do children know their parent's HIV status? Parental reports of child awareness in a nationally representative sample. *Ambulatory Pediatrics.* May 2006;6(3):138-144.

Barbara Stillwater

Alaska State Diabetes Program

Three years later, participants in the Diabetes Prevention Study still benefiting

Lifestyle intervention has lasting benefits in those at risk of diabetes. The effects of lifestyle intervention on diabetes risk do not disappear after active counseling has stopped, a new follow-up of the Finnish Diabetes Prevention Study shows. Three years after the end of the study, those in the intervention group still had a reduced incidence of type 2 diabetes compared with the control

Interpretation: Lifestyle intervention in people at high risk for type 2 diabetes resulted in sustained lifestyle changes and a reduction in diabetes incidence, which remained after the individual lifestyle counseling was stopped.

Lindstrom J, et al. Sustained reduction in the incidence of type 2 diabetes by lifestyle intervention: follow-up of the Finnish Diabetes Prevention Study. *Lancet.* 2006 Nov 11;368(9548):1673-9.

2007 Clinical Update on Substance Abuse and Dependency Pre-Registration

March 20-23, 2007 – Phoenix, Arizona (Pre-Registration Due February 23, 2007)

Wingate Inn Phoenix, 2520 North Central Avenue, Phoenix, AZ, 85004; (602) 716-9900.

June 26-29, 2007 – Portland, Oregon (Pre-Registration Due May 18, 2007)

Red Lion Hotel on the River, 909 North Hayden Island Drive, Portland, OR, 97217; (503) 283-4466.

PLEASE PRINT:

Name: _____ Job Title: _____

Work Facility: _____ Work Telephone: _____

Work Address: _____ Work Fax No.: _____

City/State/Zip: _____ E-mail Address: _____

Employment Category:	<input type="checkbox"/> Civil Service	<input type="checkbox"/> Commissioned Corps	<input type="checkbox"/> Direct IHS		
	<input type="checkbox"/> *Contracting Tribe	<input type="checkbox"/> *Compacting Tribe	<input type="checkbox"/> Urban Program		
	<input type="checkbox"/> Other: _____				
Professional Category:	<input type="checkbox"/> Physician	<input type="checkbox"/> PA	<input type="checkbox"/> Nurse	<input type="checkbox"/> Pharmacist	<input type="checkbox"/> Other: _____

Check Preference for Treatment Experience: Male Female

Check here if you plan to attend Friday's **OBOT** Course: Yes

Why do you want to attend the CUSAD training? In 150 words or less, write a brief summary and include the following information:

1. What clinical area are you currently working in?
2. How will you utilize the training in your current position?
3. Have you had previous chemical dependency training and what type(s)?

Supervisor's Name (print): _____ Supervisor's Approval: _____
Signature

If selected, participants are expected to attend all three days of the training session.

***Note:** There is a \$350 tuition fee for those employees of P.L. 93-638 contracted or compacted tribal facility that have taken tribal shares of the ASAPB and/or the CSC. Please make checks payable to **IHS Clinical Support Center**.

Fax by the deadline date to: Cheryl Begay, IHS Clinical Support Center, Fax (602) 364-7788.
Or register on-line at <http://www.ihs.gov/MedicalPrograms/ClinicalSupportCenter/>.

This is a page for sharing “what works” as seen in the published literature, as well as what is being done at sites that care for American Indian/Alaskan Native children. If you have any suggestions, comments, or questions, please contact Steve Holve, MD, Chief Clinical Consultant in Pediatrics at sholve@tcimc.ihs.gov.

IHS Child Health Notes

Quote of the month

“A radical is a man with both feet planted firmly in the air. A conservative is a man with two perfectly good legs, however, who has never learned to walk forward.”

Franklin Roosevelt

Article of Interest

Effect of needle size on immunogenicity and reactogenicity of vaccines in infants: randomized controlled trial. *BMJ*. 2006 Sep 16;333(7568):571. <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?CMD=search&DB=pubmed>

Editorial Comment

A cliché is that sometimes “less is more.” In this case the British demonstrate that “more is less” when it comes to needle size and vaccinations. Infants receiving intramuscular

vaccinations with a long needle (25mm) had fewer local reactions, and especially had fewer severe local reactions, when compared with infants vaccinated with a short (16mm) needle. The width of the needle did not affect the reactogenicity and immune response did not differ between groups. While we often choose the smallest needle in an effort to decrease pain, this study suggests that a longer needle may be the better choice to minimize local reactions after vaccinations.

Infectious Disease Updates. Roselyn Singleton, MD, MPH

Introducing the 2007 Child and Adolescent Immunization Schedule. Here’s a simplified version of the 2007 Child and Adolescent Immunization Schedule – although there’s nothing simple about this schedule! There are some caveats on each vaccine in the footnotes.

	Birth	2m	4m	6m	12m	15m	2yrs	4-6yrs	11-18yrs
Hep B ¹	√	√		√					
DTaP ²		√	√	√	√			√	√Tdap ¹⁰
IPV ³		√	√		√			√	
Hib		PedvaxHib®	PedvaxHib®		PedvaxHib®				
PCV ⁷		√	√	√	√				
Rotavirus ⁵		√	√	√					
MMR ⁶					√			√	
Varicella ⁶					√			√	
Hep A ⁷					√		√		
Influenza ⁸				Yearly, 6-59 month olds-----					
HPV ⁹									Gardasil®
MeningCV4 ¹¹									Menactra®

1. **HepB:** First Hep B soon after birth. You may use Pediarix® (DTaP-IPV-HepB) at 2,4, 6 months or Comvax® (HepB-PedvaxHIB) at 2, 4, and 12 months for doses 2, 3, and 4 of Hep B in infants. Final dose must be \geq 24 weeks of age, at least 8 weeks after dose #2.
2. **DTaP:** min. age 42 days. Pediarix® can be used for doses 1 - 3 of DTaP. Don't use Pediarix® in children \geq 7 yrs or for DTaP 4 or 5. DTaP 4 can be given at 12 months if 6 months since dose 3. Don't need DTaP5 if DTaP4 given after age 4 yrs.
3. **IPV:** min. 42 days. Pediarix® can be used for doses 1 - 3 of IPV. Single antigen IPV, not Pediarix®, must be used for dose 4. Do not need IPV4 if IPV3 given after age 4.
4. **Pneumo-conj.** (Prevnar®): min. age 42 days. Give at 2, 4, 6 months with booster \geq 12 months, at least 2 months from previous dose.
5. **Rotavirus** (RotaTeq®): min. 42 days. Give dose 1 at 6 - 12 weeks. Don't start series if \geq 13 weeks. All doses of RotaTeq® must be given by 32 weeks (8 months) of age.
6. **MMR and Varicella:** 12 months. 2nd MMR and Varicella routine at 4 - 6 yr, but can be given \geq 4 wks (3 mo for Var – but don't need to repeat if \geq 4 weeks) after dose 1. Varicella dose 2 can be given to Vaccine for Children eligible children through age 18.
7. **Hep A:** min. 12 months. Can be given as early as 12 months, but is forecast at 15 months. Two doses of Hep A vaccine are needed at least 6 calendar months apart.
8. **Influenza:** min. age 6 mos. 2 doses first year, 1 dose every year after for children $<$ 9 years. Routinely immunize 6 - 59 month olds.
9. **HPV** (Gardasil®): min age 9 yrs. Many states are providing for 11 - 12 year old girls. 3 dose series at "0", 1, and 6 months.
10. **Tdap** (Adacel®): min. age 11 yrs. all 11 - 18 year olds \geq 5 years since last Td/DTaP.
11. **MeningCV4** (Menactra®): 1 dose for 11 - 12 and 15 year olds and college freshmen in dorms as well as rare conditions such as asplenia and terminal complement deficiencies.

Recent Literature on American Indian/Alaskan Native Health

Doug Esposito, MD

I would like to direct the reader's attention to several excellent articles in the December issue of the *American Journal of Public Health*. This entire issue is devoted to "Embracing a Common Destiny: Health for All." It is impossible not to be consumed by the subjects of health disparities and inequities in access to health care when considering issues facing Native American and First Nations populations in North America. As regular readers of this column know, the thrust of my comments have centered on these topics since I began writing for Dr. Holve nearly one and a half years ago. Imagine my excitement whenever I come across an entire journal issue devoted to Health for All!


Health and health care for the 21st century: for all the people. *Am J Public Health*. 2006 Dec;96(12):2090-2. Epub 2006 Oct 31. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=17077403&query_hl=1&itool=pubmed_docsum

The December issue of the *American Journal of Public Health* begins with this excellent editorial by the much celebrated former Surgeon General of the United States, Dr. C. Everett Koop. Just to set the tone, here is a wonderful quote by this dedicated and outspoken public health advocate and leader: "Our capability to prevent and treat disease seems to exceed our willingness to apply our interventions." Truer (and sadder) words have not been written on the subject. We certainly have the technological and economic means to advance the health status of every American to a point far

beyond the current level. The only thing lacking is the resolve. Dr. Koop further elaborates, "I am concerned, however, that the difference between the "haves" and the "have-nots" will worsen. It is in vogue to blame problems on the victims of disease and disability, on laziness, on immigration, on those who hold opposing political points of view, or even — and perhaps most incredibly in this most wealthy of all nations — on lack of resources. Consider just two telling facts: 80% of people without health insurance are from working families and nearly 20% of them are children. Can we blame their lack of access to health care on them?"

The persistence of American Indian health disparities. *Am J Public Health*. 2006 Dec;96(12):2122-34. Epub 2006 Oct 31. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=17077399&query_hl=1&itool=pubmed_DocSum

This paper chronicles the 500-year history of health disparities for American Indian populations. It explores the diverse explanations proposed over time that have been used to rationalize, and sometimes address, the presence and persistence of these disparities. Beginning with theories of providence of God and working through to more contemporary ideas emphasizing environment, behavior, biology, and socioeconomics, the author takes the reader on a wonderful post-Columbian journey, presenting a novel, interesting, and important approach to answering some key contemporary questions. "Do American Indians have intrinsic susceptibilities to every disease for which disparities have existed? Or does the history of disparity after disparity suggest that social and



economic conditions have played a more powerful role in generating Indian vulnerability to disease?" Yet again, the issue of biology vs. society rears its ugly head (please see the "Additional Reading" section below).

In reading this article, I wonder how far we really have come since the late 1800s. "Medical campaigns, for example, suffered from inadequate funding. Commissioner of Indian Affairs T. J. Morgan compared the salaries paid to government physicians in the Army, Navy, and IHS and divided these sums by the populations served. He then calculated a crude estimate of how the government valued people: \$21.91 per soldier, \$48.10 per sailor, and \$1.25 per Indian." Now, doesn't this sound frighteningly similar to contemporary estimates of disparities in per capita health expenditures? As reported in a study done by the United States Commission on Civil Rights in April 2004: "HHS estimates the FY 2003 annual per capita health care spending for the general population at \$5,065. In contrast, IHS spent \$1,914 per eligible user, or 38 percent of that spent by the general population." What rightfully stirs the ire of Native Americans, though, is this: the U.S. spends \$3,803 per year per federal prisoner, almost twice as much as it spends for an American Indian or Alaska Native.

I believe it is absolutely critical for our society to acknowledge and come to grips with the true etiology of the health disparities endured by minority populations. Society needs to take responsibility for the existence of these deplorable realities. The fallacy that there exists a real biologic cause, conveniently reflected by skin color and body habitus alone, only serves to divert attention away from the true problem; that being disparities in wealth and power, or more simply, racism and unequal access and opportunity. But please, don't take my word for it, go to the article and deliberate for yourselves . . . then let's all work together as a society on a solution.

Additional Reading

On the issue of genomics, please see the March 2006 issue of the *IHS Child Health Notes* on Race, Genetics, and the Biologic Versus Social Determinants of Health and Health Disparities: <http://www.ihs.gov/MedicalPrograms/MCH/M/documents/ICHN306.doc>

Broken Promises: Evaluating the Native American Health Care System. US Commission on Civil Rights, April 2004. <http://www.usccr.gov/pubs/nahealth/nabroken.pdf>

Infant mortality trends and differences between American Indian/Alaska Native infants and white infants in the United

States, 1989-1991 and 1998-2000. *Am J Public Health*. 2006 Dec;96(12):2222-7. Epub 2006 Oct 31.

http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=17077400&query_hl=1&itool=pubmed_DocSum

In another article highlighting the "Health for All" theme, this study offers a detailed analysis of measurable disparities in infant and neonatal mortality suffered by Native Americans. In the October issue of the *IHS Child Health Notes*, I discussed an *MMWR QuickStats* showing that infant mortality for American Indian/Alaska Native mothers ranks second highest in the US, surpassed only by the rate for non-Hispanic blacks. The paper appearing in the December issue of the *American Journal of Public Health* is a much more detailed and relevant analysis.

The authors analyzed linked birth/infant-death data in order to generate birthweight-specific and cause-specific neonatal and infant mortality rates for white and AI/AN infants born at ≥ 20 weeks' gestation in the U.S. in 1989 - 1991 compared with 1998 - 2000. Although some definite and significant gains have been made over the period, presumably preventable disparities still do exist. Of note is that decreases in disparities between AI/AN and White infants were identified in some measures while increases were found in others.

On a slightly more cheery note, the authors found that AI/AN infant mortality overall was only 1.7 times that of White infants. This compares favorably with historical data. Infant mortality rates once were several-fold higher for Native Americans than for the general US population. However, although real progress has been made, it has been impossibly (and perhaps unforgivably) slow. As a result of the disparity in infant mortality, countless and needless infant deaths have occurred over the centuries, exacting a terrible toll on the well being of the afflicted populations through chronic, unrelenting emotional trauma. This situation will certainly have a lasting negative impact that can be expected to persist perhaps generations beyond the time the disparity is finally eliminated.

The authors offer a very thoughtful analysis of the troubles and the achievements in infant mortality over a ten-year period. This article is well worth a look by any practitioner delivering care to AI/AN infants or by anyone seeking to assist in the reduction and eventual elimination of health disparities for this population.

Additional Reading

October 2006 issue of the *IHS Child Health Notes*: <http://www.ihs.gov/MedicalPrograms/MCH/M/documents/ICHN1006.doc>

Locums Tenens and Job Opportunities

If you have a short or long term opportunity in an IHS, tribal or urban facility that you'd like for us to publicize (i.e., AAP website or complimentary ad on Ped Jobs, the official AAP on-line job board), please forward the information to indianhealth@aap.org or complete the on-line locum tenens form at <http://www.aap.org/nach/locumtenens.htm>.

Save the Date

Title: "Making Data Count: Measuring Diabetes and Obesity in Indian Health System
Sponsor: IHS Division of Diabetes Treatment and Prevention
When: July 24-26, 2006
Where: Westin La Paloma Resort
Tucson, Arizona

Who Should Attend:

Anyone who works with data addressing diabetes and obesity in Indian health care systems. The conference will provide plenary sessions and concurrent workshops with both clinical and technical applications. All users, new and experienced, will benefit from this Conference.

Program Overview:

An innovative conference that will provide opportunities for individuals and programs working in Indian health care systems to

- review current and advanced data systems and analysis tools relevant to diabetes and obesity data
- learn the impact of cost and quality issues

- network and share common issues and best practices for capturing, reporting, using, and improving data to help improve the lives of American Indian and Alaska Native people

Program Objectives:


- Identify current status of diabetes and obesity data-related issues
- Identify currently available and advanced diabetes and obesity related data systems and analysis tools in IHS, Tribal, and Urban (I/T/U) health care settings
- Discuss data quality issues and the impact on required data reporting, cost accounting and program evaluation
- Network and share ideas on diabetes and obesity related data issues and best practices in unique settings
- Explore opportunities for education, training, and career development in data management and program analysis in I/T/U systems

Registration: coming soon

The 12th Annual Elders Issue

The May 2007 issue of THE IHS PROVIDER, to be published on the occasion of National Older Americans Month, will be the eleventh annual issue dedicated to our elders. Indian Health Service, tribal, and Urban Program professionals are encouraged to submit articles for this issue on elders and

their health and health care. We are also interested in articles written by Indian elders themselves giving their perspective on health and health care issues. Inquiries or submissions can be addressed to the attention of the editor at the address on the back page of this issue.



Prevention of Methamphetamine Use in Indian Country: Promising Practices

Craig Love, PhD, Senior Study Director, Westat, Rockville, Maryland; and Maria Barrera, EdD, Division of Systems Development, Center for Substance Abuse Prevention, Rockville, Maryland

Prevention is Prevention is Prevention

Methamphetamine abuse is only one of many issues that American Indians and Alaska Natives (AI/AN) face. The “real” issues associated with methamphetamine abuse are also associated with the abuse of other substances, mental and physical health problems, and other barriers to a generally healthy lifestyle. Thus, it is important to understand the context of the growing methamphetamine problem in Indian Country.

Key Issues in Indian Country

AI/AN communities possess a rich cultural history; the traditional perception is that an individual’s identity reflects the collective identity of the community.^{1,2} In these communities, when the collective identity feels pain, an individual’s sense of personal efficacy, self-esteem, interpersonal satisfaction, and competence also suffers. Many AI/AN have experienced inter-generational and cultural traumas that persist today and are so apparent in the pervasive social problems that affect the lives of its members living in reservation, rural, and urban settings.

The creation of reservations and federal policies that “tried to take the Indian identity out of the Indian” left many AI/AN feeling misplaced.³ Anger, rage, and resentment are prominent among AI/AN youth.⁴ AI/AN inherit community stressors, and their family lives are marked by the effects of poverty, broken homes, lack of supervision, teenage pregnancy, domestic violence, child abuse, and neglect. The result is often reflected in disabilities, incarceration, death, suicide, sexual abuse, rape, unemployment, school failure, and loss of traditional ways.

There are other risk factors that affect AI/AN that require special focus. First, five million Americans have used methamphetamine in the United States, thereby creating a highly profitable enterprise for gangs affecting Indian Country today.⁴ Second, the frontier nature of many reservations lends itself to the clandestine labs used to produce methamphetamine.⁵ Also, there is insufficient law enforcement coverage in Indian Country to deal with rising gang activity. Third, residues and discharged chemicals from the methamphetamine labs are highly toxic substances that penetrate buildings and the soil, hurting not only our people, but also mother earth.⁴ Fourth, youth and adults on

reservations credit “boredom” or “nothing for kids to do” as major contributors to their eventual involvement with substances. Fifth, the victims of methamphetamine abuse include family members who are humiliated, traumatized, and physically threatened by their addicted relatives. Finally, elders and tribal officials in many communities cite the lack of education and prevention efforts. They see a need for AI/AN to reconnect to the traditional ways in order to gain the personal tools needed to solve the methamphetamine problem in Indian Country.

Not Much Evidence Yet

In the past few years, SAMHSA has funded several prevention programs that focused on the prevention of methamphetamine use. Most of the methamphetamine intervention programs are still in process and the data are not yet available. Until more evidence is available, promising practices for AI/AN must address some important features of the modern Native community.⁶ One of the complicating factors in conducting evaluations and making generalizations in research on AI/AN is the very diversity within our racial group. Over 61% live in urban settings; most, but not all, are very poor; the level of assimilation within and among the various tribes stretches across a broad continuum; and the nature of substance abuse varies between tribes (e.g., males versus females or both; style of abuse, such as bingeing; etc.). There are many variations among and within AI/AN tribes, including languages, customs, and values; some tribes have no particular problems with substance abuse.⁷ One important common feature is the role of the extended family (including elders and clan mothers and fathers), which can provide positive support for healing.



The Importance of Culture

Our understanding and belief is that any prevention intervention in Indian Country needs to be holistic and culturally appropriate. However, achieving cultural appropriateness can be challenging, depending on an individual's level of exposure to traditional ways or ceremonies. For many AI/AN, who are deeply invested in their culture and traditional ways, a traditional approach is an absolute necessity. Other AI/AN may be curious and interested in learning traditional ways, although it may not be a critical need.⁸ A further complication in AI/AN prevention programs, especially in urban settings, is that Indians from many different tribal groups may participate in the programs. This mix of AI/AN tribes makes it difficult to provide anything other than generic ceremonies in a pan-Indian approach; the lack of tribal specificity may not be as effective.

Depending on the cultural disposition of the participants, methamphetamine prevention programs for AI/AN need to focus on reconnecting them with their traditional spiritual side and culture. For most AI/AN, culture will provide a sense of place in the world and a positive spiritual perspective. There are a variety of traditional prevention programs being used in Indian Country that may help methamphetamine users change their way of life and gain that spiritual connection.

Model Programs in Indian Country

To-date, the effectiveness of methamphetamine prevention programs for AI/AN have not been rigorously evaluated. The programs discussed below are presented as promising practice.

The Medicine Wheel and other culturally based activities. AI/AN work toward living a healthy life by integrating a well-balanced lifestyle that is physically, emotionally, and spiritually stable. This integrated approach is reflected in the teaching of the Medicine Wheel Model, which has four phases: 1) embracing the individual's balance and wellness; 2) letting the individual know that addiction is an evil spirit in the body and easy to overcome; 3) helping the individual understand the reasons why they have an addiction; and 4) using recovery as a time for healing, in which the addict slowly overcomes the problem by understanding who they are as AI/AN. Other traditional AI/AN approaches include healing sweat lodge ceremonies and the use of talking circles in the course of treatment.

Project Venture and Service Learning. The National Indian Youth Leadership Project (NIYLP) "Project Venture" has been in existence and developing since 1986 and has been designated as a SAMHSA Model program.⁹ It is based on traditional pre-Columbian learning methods used across the North American continent. Youth are engaged in various outdoor activities and participate in service learning, which includes search and rescue, survival and teamwork, and other community support activities. Although McClellan Hall

developed this program in Gallup, New Mexico with mostly Dine' children, he has subsequently introduced Project Venture to many different communities and helped them adapt the program to their own cultures. According to the NIYLP, there is clear evidence that the program is effective in reducing alcohol use and halting the growth in alcohol and marijuana consumption among the AI/AN youth served in the program. In an evaluation, Project Venture participants had statistically superior outcomes compared to a control group.

Native HOPE. Another promising approach is Native Helping Our People Endure (Native HOPE), a suicide prevention program that incorporates community members, leaders, and medicine men in its implementation. The concept is to provide strong community ties among the providers of prevention service and the youth the programs serve. The project clarifies and strengthens the role of individuals in the community and helps them lead by example. Training manuals are available to help individuals fulfill their roles as facilitators.¹⁰

Next Steps

Evidence-based practices for the prevention of methamphetamine abuse among AI/AN are not yet available. It is clear that there needs to be a systematic evaluation of currently funded programs to ascertain those that are the most effective on a long-and short-term basis. As the promising practices are empirically established, their application to additional tribes and communities also needs systematic evaluation. Our intent here is to present program strategies that make sense given what we do know about Indian Country.

References

1. Beauvais F, Laboueff S. Drug and alcohol abuse intervention in American Indian communities. *The International Journal of the Addictions*. 1985;20:139-171.
2. Deloria E. *Speak of Indians*. 1944. New York: Friendship Press.
3. Baca K. A losing battle: meth hits reservations. *Colorlines Magazine: Race, Action, Culture*. Summer 2005;8(2).
4. Robinson CJ. Methamphetamine Use and Sales Among Gang Members: The Cross-Over Effect. 2005. Accessed 9/25/2006 from www.ngcrc.com/methuse.html.
5. Frosch D.). A Navajo town deals with the ravages of the latest addictive drug, crystal meth. *High Times*. November 9, 2004. Accessed 9/27/2006 from www.inthesetimes.com/site/main/article/1394/.
6. Moran JR, Reaman JA. Critical issues for substance abuse prevention targeting American Indian youth. *The Journal of Primary Prevention*. Spring, 2002;22(3):201-233.

-
7. May PA. Substance abuse and American Indians: Prevalence and susceptibility. *International Journal on the Addictions*. 1982;17:1185-1209.
 8. Roosa MW, Dumka LE, Gonzales NA, Knight GP. Cultural/ethnic issues and the prevention scientist in the 21st century. *Prevention and Treatment*. 2002;5:1-13.
 9. National Indian Youth Leadership Project. Accessed 9/29/2006 from http://niylp.org/programs/project_venture.
 10. Small C, Bighorn E. *Native HOPE (Helping Our People Endure): Training of Facilitator's Manual*. Portland, OR. One Sky Center. 2006.



This publication was made possible by from the Division of Systems Development, Center for Substance Abuse Prevention (CSAP), Substance Abuse and Mental Health Services Administration (SAMHSA), US Department of Health and Human Services. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Division, CSAP, SAMHSA, or the US Department of Health and Human Services.

Letter to the Editor

To the Editor:

Dr. Steve Holve's paper in the October, 2006 PCP, "Fluoride Varnish Applied at Well Child Care Visits Can Reduce Early Childhood Caries," gave good reasons for the use of fluoride varnish by staff at well child visits. The study documented a decrease in the severity of ECC (Early Childhood Caries). There was not a reduction of ECC itself, which would mean more caries free children. Parents/guardians should be reminded that they are the key person to keep a child caries free. They should brush the child's teeth every day (six years old or younger), preferably before bed. Research has shown that children in this age range do not have the skills or language to wield a toothbrush effectively.

Dr. Andrew Casterline
Salt River Clinic Dental Department
Scottsdale, Arizona

MEETINGS OF INTEREST □

Available EHR Courses

EHR is the Indian Health Service's Electronic Health Record software that is based on the Resource and Patient Management System (RPMS) clinical information system. For more information about any of these courses described below, please visit the EHR website at http://www.ihs.gov/CIO/EHR/index.cfm?module=rpms_ehr_training. To see registration information for any of these courses, go to <http://www.ihs.gov/Cio/RPMS/index.cfm?module=Training&option=index>.

Clinical Update on Substance Abuse and Dependency (Formerly known as the Primary Care Provider Training on Chemical Dependency)

March 20 - 22, 2007; Phoenix, Arizona

June 26 - 28, 2007; Portland, Oregon

This three-day intensive workshop includes both didactic and experiential training. The curriculum is updated annually with the most current nursing, addiction medicine, and prevention information. This training is available to Indian health providers (physicians, physician assistants, nurses, and advanced practice nurses). Enrollment is limited to 30 providers (preferably 2 - 3 person teams from the same facility representing the various disciplines targeted). The conference site for the first conference is the Wingate Inn Phoenix, 2520 North Central Avenue, Phoenix, Arizona, 85004. Make your hotel room reservations by March 5, 2007 by calling (602) 716-9900. Be sure to ask for the "Indian Health Service" group rate. Conference room rates are \$141.00 per night plus tax (single). Check-in is 3 pm and check-out is 12 noon. Reservation requests received after the cut-off date will be at prevailing rates based on availability. The conference site for the June offering in Portland, Oregon will be the Red Lion Hotel on the River. For more information or to register, contact Cheryl Begay at (602) 364-7777 or e-mail Cheryl.Begay@ihs.gov. To register on-line, go to the CSC website at <http://www.ihs.gov/MedicalPrograms/ClinicalSupportCenter/>.

Office Based Opioid Treatment Course

March 23, 2007; Phoenix, Arizona

June 29, 2007; Portland, Oregon

The IHS invites all physicians and nurses to register for its upcoming Office Based Opioid Treatment (OBOT) Course to be held Friday, March 23, 2007 in Phoenix, Arizona and Friday, June 29, 2007 in Portland, Oregon. The course faculty features the top clinicians and researchers in the field. This new treatment modality reduces the regulatory burden on physicians who choose to practice opioid addiction therapy. It is open to all physicians and nurses, including federal, state, and military. For more information, contact Dr. Anthony Dekker at (602) 263-1200 or anthony.dekker@ihs.gov

19th Annual IHS Research Conference

Dates to be Announced (Late May or Early June); Phoenix, Arizona

The 19th Annual IHS National Research Conference,

"Multiple Perspectives on AI/AN Research Policy," will enhance our ability to ensure benefits of research to Native communities and peoples. The conference will also examine in depth the impact of Research Policies on the research activity in American Indian and Alaska Native (AI/AN) communities. This three-day research conference will bring together many stakeholders in AI/AN research activities, including clinicians, health administrators, educators, consumers, researchers, and community and tribal government leaders across the nation.

The hotel site for the conference has not been determined yet.

The conference is sponsored by the Indian Health Service and the IHS Clinical Support Center (the accredited sponsor). Individuals who want to present their research will need to prepare an abstract and fax, mail, or e-mail it (in Word format) to Teresa Sappier or Dr. Alan Trachtenberg by March 1, 2007 (see Call for Papers in a future issue of *The Provider* for specific instructions). If mailing, include a diskette with the abstract in Word. E-mail addresses: Teresa.sappier@ihs.gov or alan.trachtenberg@ihs.gov.

For more information, contact Dr. Phillip Smith, Conference Coordinator, 801 Thompson Avenue, TMP 450, Rockville, MD 20852; telephone (301) 443-4700; fax (301) 443-0114; or e-mail psmith@hqe.ihs.gov.

7th Annual Advances in Indian Health

May 1 - 4, 2007; Albuquerque, New Mexico

The 7th Annual Advances in Indian Health Conference is offered for primary care physicians, nurses, and physician assistants who work with American Indian and Alaskan Native populations at federal, tribal and urban sites. Medical students and residents who are interested in serving these populations are also welcome.

Both new and experienced attendees will learn about advances in clinical care specifically relevant to Native American Indian populations with an emphasis on southwestern tribes. Opportunities to learn from experienced clinicians who are experts in native health will be emphasized. Indian Health Service Chief Clinical Consultants and disease control program directors will be available for consultation and program development.

The conference format includes three and a half days of lectures and case discussion workshops. In early spring, the brochure will be posted on the UNM CME website at <http://hsc.unm.edu/cme>. For additional information, please contact Kathy Breckenridge, University of New Mexico Office of Continuing Medical Education, at (505) 272-3942, or e-mail the UNM CME Office to request a brochure at CMEWeb@salud.unm.edu.

If you would like to review a sample program, you can find it on the National Council of Chief Clinical Consultant's website at <http://www.ihs.gov/NonMedicalPrograms/NC4/nc4-fpAdvances.asp>.

POSITION VACANCIES □

Editor's note: As a service to our readers, THE IHS PROVIDER will publish notices of clinical positions available. Indian health program employers should send brief announcements on an organizational letterhead to: Editor, THE IHS PROVIDER, The IHS Clinical Support Center, Two Renaissance Square, Suite 780, 40 North Central Avenue, Phoenix, Arizona 85004. Submissions will be run for two months, but may be renewed as many times as necessary. Tribal organizations that have taken their tribal "shares" of the CSC budget will need to reimburse CSC for the expense of this service. The Indian Health Service assumes no responsibility for the accuracy of the information in such announcements.

**Medical Director/Primary Care Physician
Cherokee Elder Care Program of All-Inclusive Care for the Elderly (PACE)
Cherokee Nation Comprehensive Care Agency;
Tahlequah, Oklahoma**

In preserving seniors with their independence, Cherokee Nation Comprehensive Care Agency is pleased to announce an opening for a Medical Director/Primary Care Physician. This position would be responsible for the development, implementation, and oversight of the program's primary care services and all clinical aspects of care and related administrative tasks. Minimum requirements include graduate of an accredited school of medicine, geriatrics training preferred, license to practice medicine in Oklahoma, and at least 2 – 3 years clinical, supervisory and administrative experience.

Interested applicants may e-mail resumes to resumes@cnhhs.org or contact the Personnel Office at (918) 456-5051, extension 223. For more information please contact Cherokee Nation Home Health Services at www.cherokeehomecare.org.

**Physician
Wilma P. Mankiller Health Center; Stilwell, Oklahoma**

The Wilma P. Mankiller Health Center has an immediate opening for a full-time family medicine or internal medicine physician. This 37,300 sq. foot facility is one of six rural ambulatory clinics operated by the Cherokee Nation. Other services offered at this facility include dental, radiology, public health nursing, an in-depth diabetes program, physical therapy, pharmacy, and laboratory. Cherokee Nation offers competitive salaries, excellent benefits, loan repayment options, no weekends, no call, and relocation expenses are available.

If interested in this exciting opportunity, please submit a completed Cherokee Nation application along with copies of degrees and/or certificates to Cherokee Nation Health Administration Office, Attn: Kathy Kilpatrick or Angie Cone,

PO Box 948, Tahlequah, OK 74465; telephone (918) 453-5000; fax (918) 458-6174; or e-mail kathy-kilpatrick@cherokee.org or angie-cone@cherokee.org. We would also like to extend an invitation to come and visit our clinic – we feel confident you'll love our charming Southern hospitality!

For more detailed information regarding job listings and for an application, log onto our website at www.cherokee.org. Applicants with Indian preference must submit a copy of their Certificate Degree of Indian Blood (CDIB) along with their application. All applicants will be required to pass a pre-employment drug screen and complete a background check.

**Physician
AMO Salina Health Center; Salina, Oklahoma**

The AMO Salina Health Center has an immediate opening for a full-time family medicine physician or pediatrician. This facility, located in Salina, is one of six rural ambulatory clinics operated by the Cherokee Nation. Other services offered at this facility include radiology, public health nursing, in-depth diabetes program, pharmacy, and laboratory. Cherokee Nation offers competitive salaries, excellent benefits, loan repayment options, no weekends, no call, and relocation expenses are available.

If interested in this exciting opportunity, please submit a completed Cherokee Nation application along with copies of degrees and/or certificates to Cherokee Nation Health Administration Office, Attn: Kathy Kilpatrick or Angie Cone, PO Box 948, Tahlequah, Oklahoma 74465; telephone (918) 453-5000; fax (918) 458-6174; or e-mail kathy-kilpatrick@cherokee.org or angie-cone@cherokee.org. We would also like to extend an invitation to come and visit our clinic – we feel confident you'll love our charming southern hospitality!

For more detailed information regarding job listings or for an application, log onto our website at www.cherokee.org. Applicants with Indian preference must submit a copy of their Certificate Degree of Indian Blood (CDIB) along with their application. All applicants will be required to pass a pre-employment drug screen and complete a background check.

**Full Time General Dentist
Colville Nation Community Health Centers,
Confederated Tribes of the Colville Reservation,
Inchelium and Keller, Washington**

The Colville Nation Community Health Centers (CNCHC) are currently seeking a full-time general dentist. The CNCHC is a Community Health Center facility with about 10,400 patient encounters per year, including medical, dental,

and optometry services. The CNCHC is owned and operated by the Confederated Tribes of the Colville Reservation under a PL-638 contract with the Indian Health Service and also a Public Health 330 grant. CNCHC offers competitive salaries, excellent benefits, potential loan repayment options, no weekends, no evening call, and relocation expenses are available. The hours of service are 7:30 am to 4:00 pm, Monday through Friday.

For detailed information, applicants may contact Mr. Philip Arnold, Clinic Manager or Anna Sandvig, Office Assistant, Human Resources at (509) 722-7610 or (509) 722-7686 between 7:30 am or 4:00 pm; or may send their applications to P.O. Box 290, Inchelium, Washington 99138-0290; or may send applications by e-mail to philip.arnold@colvilletribes.com or anna.sandvig@colvilletribes.com.

All applicants must be licensed in the state of Washington or eligible for licensure in the state of Washington. All applicants will be required to pass a pre-employment drug screen and complete a background check.

Family Practice Physician Ilanka Community Health Center; Cordova, Alaska

This is an outpatient position at a new community health center with some inpatient and emergency room duties at the Cordova Community Medical Center (a ten-bed Hospital with on-site long-term care beds). Cordova is a jewel. It is a small, beautiful community situated in southeast Prince William Sound between the Gulf of Alaska, the Sound, and the Copper River Delta. This is a great job that offers a very friendly and appreciative patient population; a satisfying practice with a nice mix of outpatient, ER and inpatient medicine; lots of prenatal care and general pediatrics (no deliveries; the hospital is not currently doing deliveries); a current staff of one full-time physician, one locum tenens physician, and four part-time physician assistants; an extremely flexible schedule (the four part-time physician assistants make it very easy for the docs to leave town whenever they need to; a very friendly town; good schools; most of town in easy walking distance to the clinic and hospital (car ownership optional!); tremendous access to outdoor sports and activities, including excellent hiking, cross country skiing, alpine skiing (small municipal ski area-single chair lift), world class kayaking, heli-skiing, ice skating, fishing and hunting; and Copper River Salmon! There is easy access to Anchorage by 45 minute flight or to the road systems at Valdez, Whittier, or Seward by beautiful ferry trip. Salary and benefits are competitive. Please contact Dr. Murray Buttner at (907) 424-3622 (work), or (907) 424-3651 (home); e-mail murray@ilanka.org. Check out the slideshow of Cordova photos at www.cordovaalaska.net.

Primary Care Physician Zuni Comprehensive Community Health Center; Zuni, New Mexico

The Zuni Comprehensive Community Health Center (Zuni-Ramah Service Unit) has openings for full-time primary care physicians starting in the spring of 2007. This is a family medicine model hospital and clinic providing the full range of primary care — including outpatient continuity clinics, urgent care, emergency care, inpatient (pediatrics and adults) and obstetrics — with community outreach, in a highly collaborative atmosphere. For a small community hospital, we care for a surprisingly broad range of medical issues. Our professional staff includes 14 physicians, two PAs, two CNMs, a podiatrist, dentists, a psychiatrist, a psychologist, optometrists, physical therapists, and pharmacists. Our patient population consists of Zunis, Navajos, and others living in the surrounding area.

Zuni Pueblo is one of the oldest continuously inhabited Native American villages in the US, estimated to be at least 800 - 900 years old. It is located in the northwestern region of New Mexico, along the Arizona border. It is high desert, ranging from 6000 - 7000 feet in elevation and is surrounded by beautiful sandstone mesas and canyons with scattered sage, juniper, and pinon pine trees. Half of our medical staff has been with us for more than seven years, reflecting the high job and lifestyle satisfaction we enjoy in this community.

For more information, contact John Bettler, MD at (505) 782-7453 (voice mail), (505) 782-4431 (to page) or by e-mail at jbettler@abq.ihs.gov. CVs can be faxed to (505) 782-4502, attn. John Bettler.

Laboratory Manager Winslow Indian Health Care Center; Winslow, Arizona

The Winslow Indian Health Care Center (WIHCC) in northern Arizona is currently looking for a laboratory manager for our COLA-accredited, high complexity medical laboratory. The lab is staffed with four technologists, two technicians, a phlebotomist/aide, and a secretary/phlebotomist. Lab staff provides services for approximately 50,000 outpatient primary care visits yearly. WIHCC offers a broad scope of ambulatory care clinics from 8 am until 5 pm Monday through Friday, as well as an urgent care center open daily, including weekends, until 11 pm. WIHCC also operates two field clinics five days a week on the Navajo Reservation, at Leupp and Dilkon, where only point-of-care lab testing is done on-site by clinical staff.

Winslow offers an awesome mix of professional, cultural, and recreational opportunities. We are located just seven miles from the breathtaking beauty of Navajoland and its people, and 50 miles from Flagstaff – a university town with extensive downhill and cross-country skiing, where many of our employees choose to live. Attractive salary and benefits, as well as a team oriented, supportive work environment are key to our mission to recruit and retain high quality professional staff.

WIHCC became an ISDA 638 contracted site in 2002, and has experienced steady growth and enhancement of programs and opportunities since the transition from a direct IHS program. Please contact Frank Armao, MD, Clinical Director, if you are interested in pursuing an opportunity here. He can be reached by e-mail at frank.armao@wihcc.org; telephone (928) 289-6233.

**Nursing Supervisor
Seattle Indian Health Board; Seattle, Washington**

The Seattle Indian Health Board, celebrating 36 years of service in Washington State, is a multi-service community health center for medical, dental, mental health, substance abuse, and community education services. Relocate to our Evergreen State that offers a wide assortment of attractions: the high desert, the rain forest, alpine wilderness, the inland waters of Puget Sound, pristine islands, famous mountains, and rivers large or intimate.

We are currently recruiting for a Nursing Supervisor. This position will participate as a member of the medical team to provide nursing services to clients. Works closely with providers to expedite care of patients. Assess patients' needs, plans nursing care and provides nursing care, health education, counseling, triage, and referral as indicated. Manage the flow of patients through the clinic, delegating tasks to other team members as needed. Oversight of nursing services and some special projects. Ensure competency, coverage, quality, and appropriateness of nursing services. Medical clinic management of triage, phone consultations, infection control policies and procedures, employee health, and daily operations, in collaboration with other clinical and administrative staff.

BSN required; graduation from an accredited program in registered nursing. Experience in ambulatory health care and community health nursing preferred. Experience in telephone consultation desirable. Supervisory experience required. Registered Nurse with current licensure in Washington State. Current CPR certification or must be obtained within 90 days of hire on employee's personal time. Washington State Driver's License. Familiarity with health and social issues facing American Indians/Alaska Natives and a desire to promote the delivery of appropriate health services to this population.

Please visit our website at www.sihb.org; or call our job line (206) 324-9360, ext 1123. You may also send application materials to PO Box 3360, Seattle, Washington 98114; or e-mail cindyp@sihb.org.

**Behavioral Health Consultant
Bemidji Area IHS; Bemidji, Minnesota**

Permanent full time federal position with regional office to serve IHS, tribal, and urban health care programs as lead of behavioral health program. Serves as the expert on mental health, substance abuse, and program integration. Qualifications: psychiatrist, psychologist, or psychiatric nurse practitioner. Salary dependent on qualifications (GS 13 – 15).

Indian Preference applies. Open until filled. Please contact the Aberdeen Are IHS Personnel Office for application at (605) 226-7399. For more information, go to http://www.ihs.gov/jobs/careerdevelop/jobs_index.asp.

**Family Practice Physician
Winslow Indian Health Care Center; Winslow, Arizona**

The Winslow Indian Health Care Center (WIHCC) in northern Arizona is currently looking for a family practice physician who is interested in a broad scope of practice, preferably including obstetrics. We have a staff of 12 physicians, including a surgeon, and eight Family Nurse Practitioners. We offer comprehensive ambulatory and urgent/emergent care to patients at our health center in Winslow, where we are currently completing construction on a state-of-the-art, seven bed urgent care center. WIHCC also operates two field clinics five days a week on the Navajo Reservation, at Leupp and Dilkon. Our physicians provide inpatient care at the local community hospital, the Little Colorado Medical Center, where obstetrical back-up is readily available. Winslow offers an awesome mix of professional, cultural, and recreational opportunities. We are located just seven miles from the breathtaking beauty of Navajoland and its people, and 50 miles from Flagstaff, a university town with extensive downhill and cross-country skiing, where several of our staff choose to live.

WIHCC became an ISDA 638 contracted site in 2002, and has experienced steady growth and enhancement of programs and opportunities since the transition from a direct IHS program. Please contact Frank Armao, MD, Clinical Director, if you are interested in pursuing an opportunity here. He can be reached at frank.armao@wihcc.org; telephone (928) 289-6233.

**Physician
Redbird Smith Health Center; Sallisaw, Oklahoma**

The Redbird Smith Health Center has an immediate opening for a full-time family medicine or internal medicine physician. This facility, located in Sallisaw, is one of six rural ambulatory clinics operated by the Cherokee Nation. Other services offered at this facility include dental, radiology, public health nursing, in-depth diabetes program, pharmacy, and laboratory. Cherokee Nation offers competitive salaries, excellent benefits, loan repayment options, no weekends, no call, and relocation expenses are available.

If interested in this exciting opportunity, please submit a completed Cherokee Nation application along with copies of degrees and/or certificates to Cherokee Nation Health Administration Office, Attn: Kathy Kilpatrick or Angie Cone, PO Box 948, Tahlequah, Oklahoma 74465; telephone (918) 453-5000; fax (918) 458-6174; or e-mail kathy-kilpatrick@cherokee.org or angie-cone@cherokee.org. We would also like to extend an invitation to come and visit our clinic – we feel confident you'll love our charming southern hospitality!

For more detailed information regarding job listings or for an application, log onto our website at www.cherokee.org. Applicants with Indian preference must submit a copy of their Certificate Degree of Indian Blood (CDIB) along with their application. All applicants will be required to pass a pre-employment drug screen and complete a background check.

**Internal Medicine Physician
Phoenix Indian Medical Center; Phoenix, Arizona**

The Internal Medicine Department is recruiting for a clinic internist (BC/BE) at the Phoenix Indian Medical Center; this position is available now. The internists in clinic carry a panel of continuity patients; there is no call. Hospital rounding occurs approximately one weekend/month with compensatory time off during the week. PIMC is one of the largest IHS sites with over 100 providers, 55 hospital/6 ICU beds, and multi-disciplinary/multi-specialty support. Electronic Health Record is being implemented currently. The IHS has a great benefits package for both Civil Service and Commissioned Corps physicians. The Phoenix metropolitan area offers great cultural, sports, educational, and family oriented opportunities.

For more information please contact/send CV: Amy Light MD, Chief of Medicine, Phoenix Indian Medical Center, 4212 N. 16th Street, Phoenix, Arizona 85016. Telephone (602) 263-1537; fax (602) 263-1593; e-mail amy.light@ihs.gov.

**Internal Medicine/Hospitalist
Phoenix Indian Medical Center; Phoenix, Arizona**

The Internal Medicine Department is recruiting for a hospitalist (BC/BE) at the Phoenix Indian Medical Center; the position is available now. This seven member hospitalist group provides both general medical and ICU care. There is a very reasonable schedule with 40 - 45 hour weeks. PIMC is one of the largest IHS sites with over 100 providers and 60 beds, including six ICU beds. Electronic Health Record is being implemented. The IHS has a great benefits package for both Civil Service and Commissioned Corps physicians. The Phoenix metropolitan area offers a variety of cultural, sports, educational, and family oriented opportunities.

For more information, please contact/send CV to Amy Light MD, Chief of Medicine, Phoenix Indian Medical Center, 4212 N. 16th Street, Phoenix, Arizona 85016. Telephone (602) 263-1537; fax (602) 263-1593; e-mail amy.light@ihs.gov.

**Family Practice Physician
Dzilth-Na-O-Dith-Hle Indian Health Service Health Center; Bloomfield, New Mexico**

The Dzilth-Na-O-Dith-Hle Indian Health Service Health Center is recruiting for a family practice physician for a full-time position. The medical department is staffed with three physicians and one physician assistant. This ambulatory care clinic is primary care-oriented with outpatient, dental, optometry, mental health, laboratory, radiology, and other services. The referral facility is the Northern Navajo Medical

Center in Shiprock, New Mexico, located approximately 70 miles away.

The Dzilth-Na-O-Dith-Hle Indian Health Service Health Center is located on the Navajo Reservation in the "four corners" area of New Mexico that borders Colorado, Arizona, and Utah. This area is renown for its outdoor activities including hiking, biking, skiing, kayaking, fishing, and many others. We are located approximately 40 miles southeast of Farmington, New Mexico, from where many of the staff commute. For more information about this area, go to www.farmingtonnm.org.

Either a Commissioned Corps or Civil Service position is available. For more information, please contact Lisa Mancao-Hatch, MD, at Dzilth-Na-O-Dith-Hle Indian Health Service Health Center, 6 RD 7586, Bloomfield, New Mexico 87413; telephone (505) 632-1801, extension 36732; or e-mail Lisa.Mancao-Hatch@ihs.gov.

**Family Practice Physician
Tanana Chiefs Conference, Chief Andrew Isaac Health Center; Fairbanks, Alaska**

We are seeking a board certified family practice physician preferably with obstetrics for a full-time position. We will have openings in the summer of 2006 and 2007.

The facility is a multispecialty clinic providing services in obstetric/gynecology, internal medicine, and family practice. It also includes dental, optometry, pharmacy, behavioral health, community health aides, and other services. Our referral region includes 43 villages in interior Alaska covering an area the size of Texas. Fairbanks has an outstanding school system and university. We offer a very competitive salary with a great benefits package. Commissioned Corps positions are also available. Contact Jim Kohler at (907) 459-3806 or james.kohler@tananachiefs.org.

**ARNP with Prescriptive Authority
Seattle Indian Health Board; Seattle, Washington**

The Seattle Indian Health Board, celebrating 36 of service in Washington State, is a multi-service community health center for medical, dental, mental health, substance abuse, and community education services. Relocate to our Evergreen State that offers a wide assortment of attractions: the high desert, the rain forest, alpine wilderness, the inland waters of Puget Sound, pristine islands, famous mountains, and rivers large or intimate.

We are currently recruiting for an ARNP with prescriptive authority. The ARNP functions in the medical clinic receiving referrals from primary care providers, pharmacy, and mental health specialists. The ARNP conducts assessments, prescribes any needed psychiatric medications, and monitors the patient follow-up, in collaboration with the PCP and MH Specialist. Current licensure in Washington State as a Clinical Nurse Specialist Psychiatric/Mental Health ARNP with prescriptive authority. Two years experience in community clinic mental health setting.

We also prefer familiarity with health and social issues facing American Indians/Alaska Natives and a desire to promote the delivery of appropriate health services to this population. Please visit our website at www.sihb.org; call our jobline (206) 324-9360, ext. 2200; or call the Human Resources office at (206) 324-9360, ext. 1123. You may also send application materials to PO Box 3360, Seattle, Washington 98114; or e-mail cindyp@sihb.org.

**Family Medicine Physician/Residency Site Director
Seattle Indian Health Board; Seattle, Washington**

The Seattle Indian Health Board, celebrating 36 of service in Washington State, is a multi-service community health center for medical, dental, mental health, substance abuse, and community education services. Relocate to our Evergreen State that offers a wide assortment of attractions: the high desert, the rain forest, alpine wilderness, the inland waters of Puget Sound, pristine islands, famous mountains, and rivers large or intimate.

We are currently recruiting for a family medicine physician/residency site director. This would be a board-certified family practitioner to serve as site director for our American Indian/Alaska Native Family Medical Residency Program. The Seattle Indian Health Board is a state-of-art community health center with on-site laboratory, pharmacy, nutrition, mental health, outpatient chemical dependency treatment, dental, community education, and case management services. The successful candidate will join the staff of this nationally-recognized program for urban Indians. Washington State MD license in good standing. Board Eligible or Board Certified in Family Medicine, or other Board Certification as approved by the Executive Director. Family Medicine required, Indian health required. Clinical management experience preferred. Knowledge of licensure, regulatory and accreditation standards required.

We also prefer familiarity with health and social issues facing American Indians/Alaska Natives and a desire to promote the delivery of appropriate health services to this population. Please visit our website at www.sihb.org; call our jobline (206) 324-9360, ext. 2200; or call the Human Resources office at (206) 324-9360, ext. 1123. You may also send application materials to PO Box 3360, Seattle, Washington 98114; or e-mail cindyp@sihb.org.





Change of Address or Request for New Subscription Form

Name _____ Job Title _____

Address _____

City/State/Zip _____

Worksite: IHS Tribal Urban Indian Other

Service Unit (if applicable) _____ Social Security Number _____

Check one: New Subscription Change of address

If change of address, please include old address, below, or attach address label.

Old Address _____



THE IHS PRIMARY CARE PROVIDER



A journal for health professionals working with American Indians and Alaska Natives

THE IHS PROVIDER is published monthly by the Indian Health Service Clinical Support Center (CSC). Telephone: (602) 364-7777; fax: (602) 364-7788; e-mail: the.provider@phx.ihs.gov. Previous issues of THE PROVIDER (beginning with the December 1994 issue) can be found on the CSC Internet home page (<http://www.ihs.gov/PublicInfo/Publications/HealthProvider/Provider.asp>).

Wesley J. Picciotti, MPADirector, CSC
John F. Saari, MDEditor
E.Y. Hooper, MD, MPH.....Contributing Editor
Cheryl Begay.....Production Assistant
Theodora R. Bradley, RN, MPHNursing Consultant
Erma J. Casuse, CDADental Assisting Training Coordinator
Edward J. Stein, PharmDPharmacy Consultant

Opinions expressed in articles are those of the authors and do not necessarily reflect those of the Indian Health Service or the Editors.

Circulation: The PROVIDER (ISSN 1063-4398) is distributed to more than 6,000 health care providers working for the IHS and tribal health programs, to medical schools throughout the country, and to health professionals working with or interested in American Indian and Alaska Native health care. If you would like to receive a copy, send your name, address, professional title, and place of employment to the address listed below.

Publication of articles: Manuscripts, comments, and letters to the editor are welcome. Items submitted for publication should be no longer than 3000 words in length, typed, double-spaced, and conform to manuscript standards. PC-compatible word processor files are preferred. Manuscripts may be received via e-mail.

Authors should submit at least one hard copy with each electronic copy. References should be included. All manuscripts are subject to editorial and peer review. Responsibility for obtaining permission from appropriate tribal authorities and Area Publications Committees to publish manuscripts rests with the author. For those who would like more information, a packet entitled "Information for Authors" is available by contacting the CSC at the address below or on our website at www.csc.ihs.gov.

Dept. of Health and Human Services
Indian Health Service
Clinical Support Center
Two Renaissance Square, Suite 780
40 North Central Avenue
Phoenix, Arizona 85004

PRESORTED STANDARD
POSTAGE AND FEES PAID
U.S. DEPT. OF HEALTH & HUMAN
SERVICES
PHOENIX, AZ
PERMIT NO. 5691

CHANGE SERVICE REQUESTED

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE \$300